

References

- 1 - Richard, S., Snell, M.R.C.S., L.R.C.P., M.B., B.S., M.D., Clinical Anatomy, Lippincott Williams & Wilkins., ISBN978-1-60913-446-4(2012).
- 2- Dugdale,C., David., "Female urinary tract". Medline Plus Medical Encyclopedia., (16 September 2011).
- 3- Simon, S., Neil, A., Alistair,C., John, C., 1,2,3Nephrology and hypertension., (2014).
- 4- López, M., Hoppe, B., History, Epidemiology and Regional Diversities of Urolithiasis, *PediatrNephrol.*, 25,49 - 59(2010).
- 5- Eknovan, G., History of urolithiasis, *Clin Rev Bone Min Metab.*, 2,177 – 185(2004).
- 6- Coe, FL., Evan, A., Worcester, E., Kidney stone disease, *J Clin Invest.*, 115,2598-2608 (2005).
- 7- Curhan, C., Rimm, B., Willett, C., Stampfer, J., Regional variation in nephrolithiasis incidence and prevalence among United Statesman, *J Urology.*, 151,838 – 841(2004).
- 8- Singh, I., Renal geology (quantitative renal stone analysis) by‘Fourier transform infrared spectroscopy’, *Int UrologyNephrol.*, 40,595 – 602(2008).
- 9- Collins, C., Edward, A., Short Course in Medical Terminology. Lippincott Williams & Wilkins., ISBN 0781747678(2005).
- 10- Worcester, EM., Parks, JH., Evan, AP., Coe, FL., Renal function in patients with nephrolithiasis, *J Urology.*,176(2):600-603(2006).
- 11- Stamatelou, KK., Francis, ME., Jones, CA., Nyberg, LM., Curhan, GC., Time trends in reported prevalence of kidney stones in the United States: 1976-1994. *Kidney Int.*, 63(5),1817-1823(2003).

- 12- Raju, K., and Rakkappan, C., "Determination of Chemical Compositions on Adult Kidney Stones, Annamalai University., India(2008).
- 13- Hodgkinson, A., combined qualitative and quantitative procedure for the chemical analysis of urinary calculi, *J. Clin. Path.*, **24**,147–151(1971).
- 14- Larsson, L.B., Sorbo, HG., Tiselius, S., Ohman, A., method for quantitative wet chemical analysis of urinary calculi, *Clin. Chem. Acta* .,140, 9–20(1984).
- 15- Fuss, M., Simon, J., Verbeelen, D., Weiser, M., Van Landuyt, P., Elskens, J., Chemical analysis of renal stones from 377 Belgian patients by using qualitative or quantitative methods, *Eur. Urology.*, 4, 90–93(1978).
- 16- Westbury, E.J., Some observations on the quantitative analysis of over 1000 calculi, *Br. J. Urology.*, 46, 215(1974).
- 17- Takasaki, E., An observation on the composition and recurrence of urinary calculi, *Urology. Int.*, 30, 228(1975).
- 18- Otnes, B., Urinary stone analysis, methods, materials and value, *Scand. J. Urology. Nephrol.*, (Suppl. 71) 1–109(1983).
- 19- Mandel, NS., Mandel, GS., Urinary tract stone disease in the United States veteran population: II. Geographical variation in composition. *J Urology.*, 142,1516–1521(1989).
- 20- Mandel, N., Mandel, I., Fryjoff, K., Rejniak, T., Mandel, G., Conversion of calcium oxalate to calcium phosphate with recurrent stone episodes. *J Urology.*,169,2026-2029(2003).
- 21- Worcester, J.H., E.M., Coe, FL., Evan, AP., Lingeman, J.E., Clinical implications of abundant calcium phosphate in routinely analyzed kidney stones. *Kidney Int* .,66, 777-785(2004).
- 22- Moe, O.W., *Kidney stones: pathophysiology and medical management*Lancet., 367, pp. 333–344(2006).

- 23- Mente, A., Honey, R.J., McLaughlin, J.R., Bull, S.B., Logan, A.G., Ethnic differences in relative risk of idiopathic calcium nephrolithiasis in North America. *J Urology.*, 178(5):1992-1997(2007).
- 24- Singhal, P.C., Jacobson, A.L., Mandin, H., Hyne, J.B., Calcium dynamics in idiopathic calcium stone formers. *Biochem Med.*, 29(1),122-133(1983).
- 25- Schlieper, G., Westenfeld, R., Brandenburg, V., Ketteler, M. Inhibitors of calcification in blood and urine. *Semin Dial.* 20(2):113-121(2007).
- 26- Ettinger, B. Recurrence of nephrolithiasis. A six-year prospective study. *Am J Med.*, 67,245–248 (1979).
- 27- Coe, FL., Favus, M.J., Crockett ,T., Strauss AI: Effect of low-calcium diet on urine calcium excretion, parathyroid function and serum 1.25(OH) 2D3 levels in patients with idiopathic hypercalciuria and normal subjects. *Am J Med.*, 72,25–32(1982).
- 28- Barcelo, P., Wuhl, O., Servitge, E., Rousaud, A., Pak, C.Y.C., Randomized double-blind study of potassium citrate in idiopathic hypocitraturic calcium nephrolithiasis. *J Urology.*, 150, 1761–1764(1993).
- 29- Pak, C.Y.C., Fuller, C., Idiopathic hypocitraturic calcium oxalate nephrolithiasis successfully treated with potassium citrate. *Ann Intern Med.*, 104, 33–37(1986).
- 30- Gillen,DL., Worcester, E.M., Coe, F.L., Decreased renal function among adults with a history of nephrolithiasis: a study of NHANES III. *Kidney Int.*, 67(2):685-690(2005).
- 31- Curhan,G., Willett, W., Speizer, F., Spiegelman, D., Stampfer, M., Comparison of dietary calcium with supplemental calcium and other nutrients as factors affecting the risk for kidney stones in women. *Ann Intern Med.*,126, 497–504(1997).

- 32- Curhan, G.C., Willett, W.C., Knight, E.L., Stampfer, M.J., Dietary factors and the risk of incident kidney stones in younger women (Nurses' Health Study II). *Arch Intern Med.*, 164, 885–891(2004)
- 33- Prien, E.L., Frondel, C. J., *Urology.* 57,949(1947).
- 34- Murphy, B.T., Pyrah, L.N., Br. J., *Urology.*, 34, 129(1962).
- 35- Nancollas, G.H., Koutsoukos, P.G.,*Prog. Cryst. Growth Charact.*, 3, 77(1980).
- 36 - van Kemenade, M.J.J.M., Bruyn, P.L. P.L., *Colloid Inter, J. J. Sci.*, 118, 564(1987).
- 37- Lundager, H.E., *Acta Chem., Scand.* 24, 1677(1970).
- 38- Spencer, M., *Chromatogr. J.*, 166, 435(1978).
- 39- Pak, C.Y.C., *Cryst, J. Growth.*, 53, 202(1981).
- 40- Bigi, A., Falini, G., Foresti, E., Gazzano, M., Ripamonti, A., Roveri, N., *J. Inorg. Biochem.*, 4(1993).
- 41- Amjad, Z., Koutsoukos, P.G. , Nancollas, G.H., *J. Colloid Inter. Sci.*, 101, 250(1984).
- 42- Salimi, M.H., Heughebaert, J.C., Nancollas, G.H., *Langmuir*, 1., 119(1985).
- 43- Grases, F., S'ohnel, O., Vilacampa, A.I., March, J.G.,*Clin. Chim. Acta.*, 244-245(1996).
- 44- Daudon, M., Bader, C.A., Jungers, P., *Scanning Microsc.*, 7, 1081(1993).
- 45- Leusmann, D.B., *Scand. J. Urology.*, 25, 141(1991).
- 46 Weiss, M., Liapis, H., Tomaszewski, J.E., Arend, L.J., Pyelonephritis and other infections, reflux nephropathy, hydronephrosis, and nephrolithiasis". In Jennette, JC, Olson, JL; Schwartz, MM. *Heptinstall's*

Pathology of the Kidney. 2 (6th ed.) "Chapter 22, Philadelphia: Lippincott Williams & Wilkins. pp. 991–1082(2007).

47- Moe, OW., "Kidney stones: pathophysiology and medical management". The Lancet., 367 (9507) 333–344(2006).

48-Johri, N., Cooper, B., Robertson, W., Choong, S., Rickards, D., Unwin, R. "An update and practical guide to renal stone management". Nephron Clinical Practice., 116 (3) 159–171(2010).

49- Halabe, A., Sperling, O., "Uric acid nephrolithiasis". Mineral and Electrolyte Metabolism., 20 (6): 424–431(1994).

50- Evan ,AP., Coe, F.L., Lingeman, J.E., Shao, Y., Matlaga, Kim, S.C., Renal crystal deposits and inhistopathology in patients with cystine stones. Kidney.,69,2227-2235(2006).

51- Grases, F., Costa-Bauza, A., Ramis, M., Montesinos, V., Conte, A.,Clin .Chim.Acta., 322 ,29–36(1998).

52-G. Schubert, Urology, Res., 34, 146–150(2006).

53- Tiselius ,HG., Berg, C., Fornander, A.M., Nilsson, M.A., Effects of citrate on the different phases of calcium oxalate crystallization. Scanning Micro sc., 7,381-389(1993).

54 -Rose ,GA., Stone analysis by wet chemistry. In: Rose GA, editor. Urinary Stones: Clinical and laboratory aspects Lancaster. MTP Press., p. 67-75(1982).

55- Larsson, L., Sorbo, B., Tiselius, H.G.O., On the analysis of urinary calculi. In: Brockis JG, Finlayson B, editors. Urinary calculus. International Urinary Stone Conference. Littleton (MA): PSG Publishing Company., p. 383-386(1981).

56- Otens, B., urinary stone analysis, methods, materials and value. Scand J Urology Nephrol Suppl .,71,1-109(1983).

- 57 - Hodgkinson, A.A., combined qualitative and quantitative procedure for the chemical analysis of urinary calculi. *J Clin Path.*,24,147-151(1971).
- 58- Larsson, L., Sorbo, B., Tiselius, H.G., Ohman, S.A., method for quantitative wet chemical analysis of urinary calculi. *Clin Chim Acta.*, 140, 9-20(1984).
- 59- Robertson, W.J., Peacock, M., Heyburn, P.J., Risk factors in calcium stone disease of the urinary tract. *Br J Urology.*, (1978)
- 60- hlstrand, A., Larsson, C., Tiselius, HG., Variations in urine composition during the day in patients with calcium oxalate stone disease. *J Urology.*, 131, 77-81(1984).
- 61- Huang, HS., Ma, M.C., Chen, C.F., Chen, J., Lipid peroxidation and its correlations with urinary levels of oxalate, citric acid, and osteopontin in patients with renal calcium oxalate stones. *J Urology.*, 62, 1123-1128(2003).
- 62- Abdel-Halim, R.E., Baglaf, A.O., Sibaei, A.I., Merzibani, M., Hadrami, M.S., Noorwali, I., Urolithiasis in the Western Region of Saudi Arabia: A clinical, biochemical and epidemiological study, Riyadh (KSA): King Abdul-Aziz City for Science and Technology. p. 74-278(1996).
- 63- Elliot, J.S., Calcium oxalate urinary calculi: Clinical and chemical aspects. *Medicine .*, 62, 36-43(1983).
- 64- Leonetti, F., Dussol, B., Berthezene, P., Thirion, X., Berland, Y., Dietary and urinary risk factors for stones in idiopathic calcium stone formers compared with healthy subjects. *Nephrol Dial Transplant.*,13, 617-622(1998).
- 65- Ibrahim, A., Urinary lithogenesis in Sudanese patients: A Study on 125 stone formers. *J Urology.*, 121,572-574(1979).

- 66- Ombra, M.N., Casulac, S., Biino, G., Maestrone, G., Cardia, F., Melis, P., Urinary glycosaminoglycans as risk factors for uric acid nephrolithiasis: case control Study in a Sardinian genetic isolate. *J Urology.*, 2, 416-420(2003)
- 67- Corbetta, S., Baccarelli, A., Aroldi, A., Vicentini, L., Fogazzi, G.B., Eller-Vainicher, C., Ponticelli, C., Beck-Peccoz, P., Spada, A., Risk factors associated to kidney stones in primary hyperparathyroidism. *J Endocrinol Invest.*, 28, 122-128(2005).
- 68- Kinder, J.M., Clark, C.D., Coe, B.J., Asplin, J.R., Parks, J.H., Coe, F.L., Urinary stone risk factors in the siblings of patients with calcium renal stones. *J Urology.*, 167, 1965-1967(2002).
- 69- Curhan, G.C., Willett, W.C., Speizer, F.E., Stampfer, M.J., Twenty four hour urine chemistries and the risk of kidney stones among women and men. *Kidney Int.*, 59, 2290-2298(2001).
- 70-Siener, R., Glatz, S., Nicolay, C., Hesse, A., Prospective study on the efficacy of a selective treatment and risk factors for relapse in recurrent calcium oxalate stone patients. *Eur Urology.*, 44, 467-474(2003).
- 71- Mittal, R.D., Kumar, R., Mittal, B., Prasad, R., Bhandari, M. Stone composition, metabolic profile and the presence of the gut inhabiting bacterium *Oxalobacter formigenes* as risk factors for renal stone formation. *Med Princ Pract.*, 12, 208-213(2003).
- 72- Yagisawa, T., Chandhoke, P.S., Fan, J., Metabolic risk factors in patients with first- time and recurrent stone formations as determined by comprehensive metabolic evaluation. *J Urology.*, 52, 750-755(1998).
- 73- Abomelha, M.S., Al-Khader, A.A., Arnold, J. Urolithiasis in Saudi Arabia. *J Urology.*, 35, 31-34(1990)

74. Husain, I., Badsha, SA., Al-Ali, I.H., Walton, M., Sahab, A., Jaffree, S., A Survey of urinary stone disease in Abu Dhabi Emirate, Medical Journal Supplement 1., 17-33(1979).
75. Hanash, KA., Bissada, N.K., Woodhouse, N.J.Y., Pattern of calcium metabolism in normo and hyper calciuric patients with calcium urolithiasis in Saudi Arabia. J Urol., 26(1985).
76. Ljunghall, S., Regional variations in the incidence of urinary stones. Letter to the editor. Br Med J., 18, 440(1978).
77. Thomas, J., Thomas, E., Charranson-Maistre, G., Barthelemy, C., Sleiman, B., Desgrez, P., An Epidemiologic Study of Calcium Oxalate Lithiasis. Urinary Calcium and Oxalate Excretion. Sem Hop Paris., 63, 1491-1498(1987).
78. Koide, T., Bowyer, R.C., Brockis, J.G., Comparison of urinary oxalate excretion in urolithiasis patients with and without hypercalciuria. Br J Urol 57, 505-509(1985).
79. Pizzato, AC., Barros, E.J.G., Dietary calcium intake among patients with urinary calculi. Nutr Res., 23, 1651- 1660(2003).
80. Pak, C.Y., Poindexter, J.R., Peterson, R.D., Koska, J., Sakhaee, K., Biochemical distinction between hyperuricosuric calcium urolithiasis and Gouty diathesis. J Urol., 60, 789-794(2002).
81. Coe, F.E., Kavalach, A.G.,Hypercalciuria and hyperuricosuria in patients with calcium nephrolithiasis. N Engl J Med., 291, 2344-2350(1974).
82. Watanabe, T., Proximal renal tubular dysfunction in primary distal renal tubular acidosis. Pediatr Nephrol., 20,86-88(2005).
- 83 . Yuen, Y.P., Lai, C.K., Tong, G.M., Wong, P.N., Wong, F.K., Mak, S.K., Lo, K.Y., Wong, A.K., Tong, S.F., Chan, Y.W., Lam, C.W., Novel

mutations of the AGXT gene causing primary hyperoxaluria type 1. *J Nephrol.*, 17,436-440(2004).

84. Font-Llitjos, M., Jimenez-Vidal, M., Bisceglia, L., DiPerna, M., de Sanctis, L., Rousaud, F., Zelante, L., Palacin, M., Nunes, V., New insights into cystinuria: 40 new mutations, genotype-phenotype correlation, and digenic inheritance causing partial phenotype. *J Med Genet.*, 42, 58-68(2005).

85. Moe, O.W., Bonny, O., Genetic hypercalciuria. *J Am Soc Nephrol* ., 16,729-745(2005).

86. Ryall, R.L., Harnett, R.M., Hibberd, C.M., Mazzachi, B.C., Mazzachi, R.D., Marshall, V.R., Urinary risk factors in calcium oxalate stone disease comparison of Men and Women. *Br J Urol.*, 60, 480-488(1987).

87. Tiselius, H.G., Almegard, L.E., The Diurnal urinary excretion of oxalate and the effect of Pyridoxine and Ascorbate on oxalate excretion. *Eur Urology.*, 3, 41-46(1977).

88- Ryall, R.L., Darroch, J.N., Marshall, V.R., The Evaluation of Risk Factors in Male Stone Formers attending a General Out-Patient Clinic. *Br J Urology.*, 56, 116-121(1984).

89 – Tiselius, H.G., Larsson, L., Validity of biochemical findings in the evaluation of patients with Urolithiasis. *Eur Urology.*, 6, 90-94(1980)

90- Tiselius, H.G., Relationship between the severity of renal stone disease and urine composition. *Eur Urology.*, 5, 323-327(1979).

91-Pierratos, A., Dharamsi, N., Carr, L.K., Ibanez, D., Jewett, M.A.S., Honey, R.J.D.A., Higher urinary potassium is associated with decreased stone growth after shock wave lithotripsy. *J Urology.*, 164, 1486-1489(2000).

- 92- Lavan, J.N., Neale, F.C., Posen, S., Urinary calculi. Clinical, Biochemical and Radiological studies in 619 patients. *Med J Aust.*, 2,1049-1061(1971).
- 93- Reis-Santos, J.M., Epidemiological Risk Factors in the South of Portugal. In: Jardin editors. Proceedings of the XXe Congress de la Societe Internationale Urologie. Paris: la Société Internationale Urologie., p. 244-246(1985).
- 94- Ljunghall, S., Danielson, B.G., Fellstrom, B., Holmgren, K., Johansson, G., Wilkstrom, B., Family history of renal stones in recurrent stone patients. *Br J Urology* 1., 57, 370-374(1985).
- 95-Curhan, G.C., Willett, W.C., Rimm, E.B., Stampfer, M.J., Family history and risk of kidney stones. *J Am Soc Nephrol.*, 8, 1568-1573(1997).
- 96- Bytyci, X., Mesaric, S., Epidemiological investigation of nephrolithiasis in the region of Sap kosovo in yugoslavia. In: Walker VR, Sutton RAL, Cameron ECB, Pak CYC, Robertson WG, editors. *Urolithiasis*. New York, London: Plenum Press., p. 699(1989).
97. Abdel-Halim, R.E., Baghlaif, A.O., Farag, A.B., Clinical chemical study of urinary stones in Saudi Arabia. I - Uric Acid stones. In: Schwille PO, Smith LH, Robertson WG, Vahlensieck W, editors. *Urolithiasis and related clinical research*. New York, London: Plenum Press., p. 715-718(1985).
98. Lee, Y.H., Huang, W.C., Lu, C.M., Tsai, J.Y., Huang ,J.K., Stone recurrence predictive score (SRPS) for patients with calcium oxalate stones. *J Urology.*, 170, 404-407(2003).
- 99 - Kok , D.J., Iestra, J.A., Doorenbos, C.J., Papapoulos, S.E., The effects of dietary excesses in animal protein and in sodium on the composition and the crystallization kinetics of calcium oxalate monohydrate in urines of healthy men. *J. Clin. Endocrinol. Metab.*, 71, 861-7(1990).

- 100 – Breslau, N.A., Brinskley ,L., Hill, K.D., Pak, C.Y.C., Relationship of animal protein-rich diet to kidney stone formation and calcium metabolism. *J. Clin. Endocrinol. Metab.*, 66,140–6(1988).
- 101- Reddy, S.T., Wang, C.Y., Brinkley, L., Pak, C.Y.C., Effect of low-carbohydrate high-protein diets on acid-base balance, stone-forming propensity, and calcium metabolism. *Am. J. Kidney Dis.*, 40, 265–274(2002).
- 102- Rule, A.D., Bergstralh, E.J., Melton, L.J. III., Li, X., Weaver, A.L., Lieske, J.C., Kidney stones and the risk for chronic kidney disease. *Clin J Am Soc Nephrol.*, 4,804-811(2009).
- 103 . Madore, F., Stampfer, M.J., Rimm, E.B., Curhan, G.C., Nephrolithiasis and risk of hypertension. *Am J Hypertens.*, 11, 46-53(1998).
- 104- Ramello, A., Vitale, C., Marangella, M., Epidemiology of nephrolithiasis. *J Nephrol.*, 13, 45-50(2000).
- 105-Caudarella, R., Rizzoli, E., Buffa, A., Bottura, A., Stefoni, S., Comparative study of the influence of 3 types of mineral water in patients with idiopathic calcium lithiasis. *J Urology.*, 159, 658-663(1998).
- 106- Coen, G., Sardella, D., Barbera, G., Ferrannini, M., Comegna, C., Ferazzoli, F., Urinary composition and lithogenic risk in normal subjects following oligomineral versus bicarbonate-alkaline high calcium mineral water intake. *Urol Int .*, 67, 49-53(2001).
- 107-Ilievski, P., Nakovski, R., Jankovski, V., Ilievska, S., Janevski, B., Soft Drinking Water and Urolithiasis. In: Walker VR, Sutton RAL, Cameron ECB, Pak CYC, Robertson WG, editors. *Urolithiasis*. London, New York: Plenum Press., p. 773(1989).
- 108- Vahlensieck, W., Influence of water quality on urolithiasis. In: Schwille PO, Smith LH, Robertson WG, Vahlensieck W, editors.

Urolithiasis and related clinical research. New York, London: Plenum Press., p. 97-103(1985).

109- Ilievski , P.M., Ilievska, S.S., Drinking water quality and urolithiasis. In: Schwille PO, Smith LH, Robertson WG, Vahlensieck W, editors. Urolithiasis and related clinical research. New York, London: Plenum Press., p. 105- 107(1985).

110- Bellizzi, V., De Nicola, L., Minutolo, R., Russo, D., Cianciaruso B, Andreucci M, et al. Effects of water hardness on urinary risk factors for kidney stones in patients with idiopathic nephrolithiasis. Nephron., 81, 66-70(1999).

111- Singh, P.P., Kiran, R., Pendse, A.K., Mathur, H.N., Bearing of drinking water quality on the prevalence of urolithiasis. In: Walker VR, Sutton RAL, Cameron ECB, Pak CYC, Robertson WG, editors. Urolithiasis. London, New York: Plenum Press., 771-772(1989).

112- Trinchieri ,A., Epidemiology of urolithiasis. Arch Ital Urology Androl ., 68, 203-249(1996).

113. Robertson, W.G., Nisa, M., Husain, I., Al-Faqih, S., Chakrabarty A, Qunibi W, et al. The Importance of Diet in the Etiology of Primary calcium and uric acid stone formation: the Arabian experience. In: Walker VR, Sutton RAL, Cameron ECB, Pak CYC, Robertson WG, editors. Urolithiasis. New York, London: Plenum Press, p. 735-739(1989).

114- Curhan, G.C., Willett, W.C., Rimm, E.B., Speizer, F.E., Stampfer, M.J., Body size and risk of kidney stones. J Amer Soc Nephrol ., 9, 1645-1652(1998).

115- Nishio, S., Yokoyama, M., Iwata, H., Takeuchi, M., Kamei, O., Sugamoto T, Obesity as one of the risk factors for Urolithiasis. Nippon Hinyokika Gakkai Zasshi ., 89, 573-580(1998).

- 116- Vagelli, G., Galabrese, G., Ferraris, V., Mazzotta, A., Pratesi, G., Gonella, M., Overweight and calcium stone disease. In: Walker VR, Sutton RAL, Cameron ECB, Pak CYC, Robertson WG, editors. Urolithiasis. London, New York: Plenum Press., 759-760(1989).
117. Shen, M., Shi, L., Li, L., Zhang, S., Zhang, C., Jiang, Y. A., case control study on urinary calculi and dietary factors. *Zhonghua Liu Xing Bing Xue Za Zhi.*, 23, 143-137(2002)..
118. Al Zahrani, H., Norman, R.W., Thompson, C., Weerasinghe, S., The dietary habits of idiopathic calcium stone formers and normal control subjects. *BJU Int* .,85, 616-620(2000).
119. Reyes ,R.L., Mirabal, M.M., Strusser, G.R., Clinicoepidemiologic behavior of urolithiasis in a rural Caribbean region. *Arch Esp Urolog,y.*,55,527-533(2002).
120. Curhan, G.C., Willett, W.C., Rimm, E.B., Stampfer, M.J. A., prospective study of dietary calcium and other nutrients and the risk of symptomatic kidney stones. *N Engl J Med* .,328: 833-838(1993).
- 121- Bailly, G.G., Norman, R.W., Thompson, C., Effects of dietary fat on the urinary risk factors of calcium stone disease. *Urology.*, 56, 40-44(2000).
- 122- Curhan, G.C., Willett, W.C., Speizer, F.E., Spiegelman, D., Stampfer, M.J., Comparison of dietary calcium with supplemental calcium and other nutrients as factors affecting the risk for kidney stones in women. *Ann Intern Med* ., 126, 497-504(1997).
- 123- Rofe, A.M., Bais, R., Conyers, R.A.J., The effect of dietary refined sugars and sugar alcohols on renal calcium oxalate deposition in ethylene glycol-treated rats. *Food Chem Toxicol* ., 24,: 397-403(1986).
- 124- Curhan, G.C., Dietary calcium, dietary protein, and kidney stone formation. *Miner Electrolyte Metab.*, 23, 261- 264(1997).

125- Trinchieri, A., Nespoli, R., Ostini, F., Rovera, F., Zanetti, G., Pisani, E. A., study of dietary calcium and other nutrients in idiopathic renal calcium stone formers with low bone mineral content. *J Urology* .,159,654-657(1998).

126- Curhan, G.C., Curhan, S.G., Dietary factors and kidney stone formation. *Compr Ther* ., 20, 485-489(1994).

127- Lewandowski, S., Rodgers, A.L.. Idiopathic calcium oxalate Urolithiasis: risk factors and conservative treatment. *Clin Chim Acta* .,345,17-34(2004).

128- von, U.,nruh, G.E., Voss, S.S.,auerbruch., T., Hesse, A., Dependence of oxalate absorption on the daily calcium intake. *J Am Soc Nephrol.*, 15, 1567-1573 (2004).

129- Rodgers A. Aspects of calcium oxalate crystallization: theory, in vitro studies, and in vivo implementation. *J Am Soc Nephrol* .,14: 351-354(1999).

130- Caudarella, R., Rizzoli,E., Buffa, A., Bottura, A., Stefoni, S., Comparative study of the influence of 3 types of mineral water in patients with idiopathic calcium lithiasis. *J Urol* .,159, 658-663(1998).

131- De Swart, P.M.J.R., Busemann Sokole, E., Wilmink, J.M., The interrelationship of calcium and magnesium absorption in idiopathic hypercalciuria and renal calcium stone disease. *J Urology.*, 159. 669-672(1998).

132- Curhan., G.C., Willett, W.C., Speizer, F.E., Stampfer, M.J., Intake of vitamins B6 and C and the risk of kidney stones in women. *J Am Soc Nephrol* .,10, 840-845(1999).

133- Curhan, G.C., Willett, W.C., Rimm, E.B., Stampfer, M.J. A., prospective study of the intake of vitamins C and B6, and the risk of kidney stones in men. *J Urol* .,155,1847- 1851(1996).

- 134- Drach ,G.W., Evaluation of urinary stone formers. *Semin Urology.*, 11, 12-19(1984).
- 135- Siener, R., Glatz, S., Nicolay, C., Hesse, A. ,The role of overweight and obesity in calcium oxalate stone formation. *Obes Res .*, 12, 106-113(2004).
- 136- Borghi, L., Schianchi, T., Meschi, T., Guerra, A., Allegri, F.,Maggiore, U., Comparison of two diets for the prevention of recurrent stones in idiopathic hypercalciuria. *N Engl J Med.*, 346, 77-84(2002).
- 137- Tiselius, H.G., Larsson, L., Urinary excretion of urate in patients with calcium oxalate stone disease. *Uroogyl Res .*,11: 279-283(1983;).
- 138- Tiselius, H.G., The effect of pH on the urinary inhibition of calcium oxalate crystal growth. *Br Urol .*,53: 470-474(1981).
- 139 - onigsberger, E.K onigsberger (Eds.) L.C. K., *Biom mineralization :MedicalAspects of Solubility*, Wiley,New York.,(2006).
- 140- Nijenhuis, T., Vallon, V., van der Kemp, A.W., Loffing, J.,Hoenderop, J.G., Bindels, R.J., Enhanced passive Ca²⁺ reabsorption and reduced Mg²⁺ channel abundance explains thiazide-induced hypocalciuria and hypomagnesemia. *J Clin Invest .*,115(6),1651-8(2005).
- 141- Pak,C.Y., Medical prevention of renal stone disease. *Nephron .*,81(suppl. 1),60-65(1999).
- 142- Goldfarb, D.S., Reconsideration of the 1998 NIH Consensus Statem66. Goldfarb DS. Reconsideration of the 1998 NIH Consensus Statement on prevention and treatment of kidney stones: are the recommendations out of date? *Rev Urology.*;4(2), 53-60 (2002).
- 143- Honow, R., Laube, M., Schneider Keble, T., and Hesse, A., Influence of grapefruit - orange - and apple - juice consumption on urinary variables and risk of crystallization. *Brit J Nutr. ,* 90,295 – 300(2003).

- 144 - Mack, M., Minimally Invasive and Robotic Surgery, Opportunities for Medical Research., vol. 285 (No. 5), p. 568-572,(2001).
- 145- Melzer, H.H. , Schur, M.O., Kunert, W., Buess, G., Voges, U., Meyer, J.U. , Intelligent Surgical Instrument System, Journal. Of Endoscopic Surgery. vol. 1, p. 165-170(1993).
- 146- Deml, B., Ortmaier, T., Seibold, U., The touch, and feel in minimally invasive surgery, IEEE Int. workshop on haptic audio visual environments and their applications ., p. 33-38(2005).
- 147- Ottermo, M., Stavdahl, O., Johansen, T., Palpation instrument for augmented minimally invasive surgery, Proc. IEEE/RSJ Int. conf. on intelligent robots, and systems ., p. 3960-3964(2004).
- 148- Schubert, G.,Urology Res. 34, 146–150(2006).
- 149- Carmona, P., Bellanato, J., Escolar, E., Biospectroscopy, 3, 331–346(1997).
- 150- Singh, V.K., Rai, A.K., Rai, P.K. Jindal, P.K. Laser Med. Sci. 24 749–759 (2009).
- 151- Abdel-Halim, V.R.E., Abdel-Halim, M.R. Saudi Med. J, 27 1462–1467.
- 152- Pollack, S.S., Carlson ,G.L.,A comparison of X-ray diffraction and infrared technics for indentifying kidney stones. Amer J Clin Pathol., 52: 656-560(1969).
- 153- Hodgkinson, A., combined qualitative and quantitative procedure for the chemical analysis of urinary calculi. J Clin Pathol .,; 24, 147-51(1971).

154- Meyers, R.A., Inductively Coupled Plasma/Optical Emission Spectrometry in Encyclopedia of Analytical Chemistry., pp. 9468–9485, John Wiley & Sons Ltd(2000).

155. Paluszkiewicz C., Jand ,S., Galka, M., Analysis of renal stones by FTIR spectroscopy. Microchimica Acta., 94 ,(1-6),45 – 48(1988).