

## References

- Abdel-Fattah AFM, Matsumoto K, Watanabe H (2000). Antinociceptive effects of *N. sativa* oil and its major constituent thymoquinone in mice. *Eur J Pharmacol* ; 14: 89-97.
- Abou BLI and Rashed MS (1995). Thin layer chromatography assay of thymoquinone in black seed oil (*Nigella sativa* L.) and identification of dithymoquinone and thymol. *J Liquid Chromatogr* ; 18: 105-15.
- Akhtar MS and Riffat S (1991). Field trial of *Saussurea lappa* roots against nematodes and *Nigella sativa* seeds against cestodes in children. *J Pakistan Med Assoc* ; 41(8): 185-187.
- Akram KM (1999). Chemical composition and medicinal properties of *Nigella sativa* Linn. *Inflammopharmacol.* 7 (1): 15-35.
- Al,Naggar TB, Gomez-Serranillos, MP, Carreto ME, Vilar AM (2003) Neuropharmacological activity of *N. sativa* L. Extract plant medical 67: 29-72.
- AL.Nageeb G, Ismail, M, AL Zuhairs A (2009). Fatty acid profile, tocopherol and total antioxidant activity of oil extract from *N. sativa* seeds, *J. Pharmacol* 10:244-50.
- Ali BH, Blunden G (2003). Pharmacological and toxicological properties of *Nigella sativa*. *Phytother. Res.* 17 (4): 299–305.

Aljassir MS (1992). Chemical composition and microflora of black cumin (*Nigella sativa*) seeds growing in Saudi Arabia. *Food Chemistry*; 45: 239-242.

Anantharmaiah GM,Brouillette CG,Engler JA,(1991). Role of Amphipathic helices in HDL Structure/function.Adv Exp Med Biol ;285:131-140.

Andaloussi, AB, louts,M; vorg ,T, Meddah,b.padma ,M, Abdellatit,S art. Haddat P (2001). The in VIOV Antidvabeted through. Activation Of The AM PK path way and increased muscle gliet 4 contont evince Based complementary and Alternative Medicine .

Anduloussi AB, Louts M, Vong T,Meddah B, Padma M, Abdellatif S, and Haddad P (2011). The in vivo anti diabetic activity of *Nigella sativa* is Medicated through activaton of the AMPK pathway and increased Muscle glut 4 content evidence based complementary and alternative Medicine.

Badary OA, Abdel-Naim AB, Abdel-Wahab MH, Hamada FM (2000). The influence of thymoquinone on doxorubicin-induced Hyperlipidemic nephropathy in rats. *Toxicol.* 143 (3): 219-26.

Bahram,B.G, Vahideh,EA, Maryam,R,and Aboltazi,g,Effect of deltry supplementation with nlegella sativa L. on seem.

Baldassarre D, Amato M,Pustina L. (2002). Increased carotid artery intima-media thickness in subjects with primary hypoalphalipoproteinemia. *Arterioscler Thromb Vase Biol* ;23:317-322.

Bamasa A, (1997). The effect of oral ingestion of *N. sativa* seeds on lipid blood parameters 5: 28-36.

Bamosa AO, Ali BA, al-Hawsawi ZA (2002). The effect of thymoquinone on blood lipids in rats. *Indian. J. Physiol. Pharmacol.* 46 (2): 195-201.

Barenholz Y.(2002). cholesterol and other membrane active sterols: from membrane evolution to “rafts” *prog lipid res* ;41:1-5.p.352  
Batiste MC, Schaefer EJ.(2002). Diagnosis and management of lipoprotein abnormalities .*Nutr Clin Care* 5:115-123.

Blankenborn, RICHARD Sheppard; Kumar, Vinay; Abbas, Abul K; Fausto, Nelson (2007). *Robbins Basic Pathology : with student consult online Access.* philadelphia: saunders.pp.345.ISBN 1-4160-2973-7 . 8<sup>th</sup> edition

Blankenhorn DH, Hodis HN (August 1993.) “ Atherosclerosis – reversal with therapy” *the western Journal of medicine* 159(2): 172 -9.

Burnett, JR, Barrett PH (2002). Apolipoprotein B metabolism :tracer kinetics, models, and metabolites, studies. *Crit Rev Clin Lab Sci* :39:89-137

Canonica L, Jommi G, Scolastico C and Bonati A (1963). The pharmacologically active principle in *Nigella sativa*. *Gazz Chim Ital* ; 93: 1404-1407.

Carmena R, Duriez P, Fruchart JC.(2004) Atherogenic Lipoprotein Particles in atherosclerosis circulation ;109(suppl III);III2-III7.

Castelli WP, Carrison RJ, Wilson PW. (1968). Incidence of coronary heart disease and lipoprotein cholesterol levels. the Framingham study. JAMA.;256:2835-2838.

Cinsberg HN (1994). lipoprotein metabolism and its relationship to atherosclerosis . Med clin Norh Am ;78:1-20.

Coleman MP, Key TJ, Wang DY. (1992). A propective study of obesity , lipids apoliporoteins and ischaemic heart disease in women . Atherosclerosis ;92:177-2192.

Cooper AD. (1997). Hepatic uptake of chylomicron remnats. J Lipid Res ;38:2173-2192.

Cooper EL, (2004). Complementary and alternative medicine when rigorous can be science. Exid .based complement. Alternative med (1):1-4.

Corder EH, Lannfelt L, Bogdanovic N. (1998) .The role of APOE polymorphisms in late -onset dementas . cell Mol life Sci ;54:928-934.

Couillard C, Bergeron N, Pascot A (2002). Evidence for impaired lipolysis in abdominally obese men: postprandial study of apolipoprotein B-48- and B-100- containing lipoproteins. Am J Clin Nutr .76:311-318.

Couillard C, Bergeron N, Pascot A (2002). Evidence for impaired lipolysis in abdominally obese men: postprandial study of apolipoprotein B-48- and B-100- containing lipoprotein . Am J Clin Nutr, 76: 311-318.

Davis HR, Veltri EP. Zetia (2007) :inhibition of Niemann-Pick CI like 1(NPCILI) to reduce intestinal cholesterol absorption and treat hyperlipidemia.J atheroscler Thromb;14:99-108.

Del Vecchio G, Del Vecchio V and Argenziano R (1964). Antibiotic action of quinines: 1. Antibiotic activity against *Staphylococcus pyrogens aureus* *in vitro* ; *Boll Soc Ital Biol* ; Sper 21: 1-3. drug, nigellone.*Egypt Pharm Bull* ; 42: 411-424 .

EL Dakha Khani, M, N.L, Mady and M.A, Halim, (2000). *Nigella sativa* L. oil protects against induce hepatotoxicity and improves serum lipid profile in rats. *Arzneimittelforschung*, 50(9): 832-6.

El Kadi M, Kandil O and Tabuni A M (1990). *Nigella sativa* and cell mediated immunity. *Arch of Aids Res* ; 232-235.

El-Dakhakhny M, Mady NI, Halim MA (2000). *Nigella sativa* L. oil protects against induced hepatotoxicity and improves serum lipid profile in rats. *ARZNEIMITTEL. FORSCH.* 50 (9): 832-6.

El-Din K, El-Tahir H, Bakeet DM (2006).The black seed (*Nigella sativa* Linnaeus)- a mine for multi cures: A plea for urgent clinical evaluation of its volatile oil. *JTU. Med. Sci.* 1: 1-19.

ElTahir KEH and Ageel AM (1994).Effect of the volatile oil of (*Nigella sativa*) on the arterial blood pressure and heart rate of the guinea-pig. *Saudi Pharm J* ; 2(4): 163-168.

ElTahir KEH, Ashour MM and Al-Harbi MM (1993). The respiratory effects of the volatile oil the black seed (*Nigella sativa*) in guinea-pigs; elucidation of the mechanism of action. *Gen Pharmacol* ; 24: 1115-1122.

Fabricant CG, Fabricant j (1999). Atherosclerosis induced by infection with Marek's disease herpesvirus chickens ". *Am Heart J* 138(5 pt 2): S465-8.

Friedewald WT, Levy RI, Fredrickson DS (1972). Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin chem*;18:499-502.

Fruchart JC, Nierman MC, Stroes ES (2004). New risk factors for atherosclerosis and patient risk assessment. *Circulation* ;109(suppl III):III15-III19.

Genest JJ Jr, Martin-Munley SS, McNamara JR (1992). Familial lipoprotein disorders in patients with premature coronary artery disease. *Circulation* ;85:2025-2033.

Ghannadi A (2005). An investigation of the analgesic and anti-inflammatory effects of *Nigella sativa* seed.

Gilani AH, Jabeen Q, Asad Ullah Khan M (2004). A review of medicinal.

Griffin BA (1994), Packard CJ. Metabolism of VLDL and LDL subclasses. *Curr Opin Lipidol* ;5:200-206.

Gurffat D, Durand D, Grualet B (1996). Relation of VLDL synthesis and secretion in the liver. *Reprod Nutr Dev* ;36:375-389.

Gursky O (2005). Apolipoprotein structure and dynamics .curr opin lipidol ;16:287-294.

Hanafy MSM and Hatem ME (1991). Studies on the antimicrobial activity of *Nigella sativa* seed (Black cumin). *J Ethnopharmacology* ; 34: 275-278.

Hevonoja T, Pentikainen MO, Hyvonen MT (2000). Structure of low density lipoprotein (LDL) particles : basis for understanding molecular change in modified LDL. *Biochim Biophys Acta* ;1488:189- 210.

Huff MW, Pollex RL, Hegele RA (2006). NPC1L1: evolution from pharmacological target to physiological sterol transporter. *Arterioscler Thromb Vasc Biol*;26:2433-2438.

JACKSON RL, McLEAN RR, Demel RL (1987). Mechanism of action of lipoprotein lipase and hepatic triglyceride lipase .*Am Heart J*;113:551-554.

Jaffar AR, Babb J, Movahed A (2004). Optimal management of hyperlipidemia in primary prevention of cardiovascular disease. *Int. J. Cardiol.* 97 (3): 355-366.

Jansen PCM (1981). Spices, condiments and medical plants in Ethiopia: Their taxonomy and agricultural significance. Publisher: Center for Agricultural Publishing and Documentation.; 76-85.

Javitt NB (1995). Cholesterol homeostasis: role of the LDL receptor. *FASEB J*;9:1378-1381.

Jialal I, Hirany SV, devaraj S (1995). comparison of an immunoprecipitation , ethod for direct measurement of LDL- cholesterol with beta – quantification (ultracentrifugation). Am J Clin pathol ;104:76-81.

Jorge R, Neutral,C, Ozaki R,ameida E (1998). Improvement in the endothelium dependent relaxation in hypercholesterolemia rabbitstreated with vitamin E, Atherosclerosis 140: 333-9.

Kostner KN, Kostner GM,(2002).nigma Lipoprotein(a): still an ? curr opin lipidol, 13: 391-396.

Kover J, Havel RJ, (2002). Sources and properties of triglyceride-rich lipoproteins containing opoB-48 and apo-100 in postprandial blood plasma of patients with primary combined hyberlipidemia. J Lipid Res 43: 1026 1034.

Kritchevsky D, (1998). Effect of triglyceride structure on lipid metabolism. Nutr Rev. 46: 177-181.

Kulkarni KR (2006). cholesterol profile measurement by vertical auto profile method clin Lab Med ;26:787-802.

Kurth HS (2002). Sequestration of aggregated low-density lipoproteins by macrophages. curr opin lipidol ;13:483-488.

Lane RM, Farlow MR (2005). lipid homeostasis and apolipoprotein E IN THE Development and progression of Alzheimers disease.J lipid Res ;:46:949-968.



Laposata M (1995). Fatty acids. Biochemistry to clinical significance. Am J clin pathol ;104:172-179.

Le PM, Benhaddou-Andaloussi A, Elimadi A, Settaf A, Cherrah Y, Haddad PS (2004). The petroleum ether extract of *Nigella sativa* exerts lipid-lowering action in the rats . JE thanopharmacolgy 44(2-3) 251-9.

Lee HM, lu K, Patel SB (2001). Genetic Basis of sitoserolemia .curr opin lipidol ;12:141 -149.

Levy E, spahis S, Sinnett D (2007). Intestinal cholesterol transport: an update and beyond .curr opin lipidol ;18:310-318.

Lund-katz S, liu L (2004), THUAHANI st. Hligh Density lipoprotein structure.front Biosci ;d1044-d1054.

Lusis AJ, Mar R, Pajukanta P (2004). Genetics of atherosclerosis. Annu Rev Genom hum Genet;5: 189-218.

Mahfouz M nd El-Dakhakhny M (1960). Chemical and pharmacological properties of the new antiasthmatic.

Mahley RW, Rall SC Jr (2000). Apolipoprotein E: far more than a lipid transport protein. Annu rev genomics Hum genet, 1: 507-537.

Mahmoud M, EL-Abhar Hand Saleh S (2002),the effect of *Nigella sativa* oil against the liver damage induce by schistosoma mansonii infection in mice. Journal of ethnopharmacology; 79: 1-11.

Marais AD, Firth JC, Blom DJ (2004). Homozygous familial hypercholesterolemia and its management. *Semin Vasc Med*;4:43-50.

Martin S, Padlla E, Ocete M, Galvez J, Jimenez J and Zarzuelo A (1993). Anti- inflammatory activity of the essential oil of *Bupleurum fruticosens*. *Planta Med* ; 59: 533-536.

Maton, Anthea; Roshan L. JeanHooooopkins, CHARLES William McLaughlin, susan.(1993).

McNamara JR, Shah PK, Nakajima K (2001). Remnant –like particle(RLP) cholesterol is an independent cardiovascular disease risk factor in women: results from the framingam heart study, *atherosclerosis*;154: 229-236.

Meral I, Yener Z, Kahraman T and Mert N (2001). Effects of *Nigella sativa* on glucose concentration, lipid peroxidation, antioxidant defence system and liver damage in experimentally- induced diabetic rabbits. *J Vet Med Ser A* ; 48: 593-9.

Michael L.P, Edward PF, Larry E.S (2010).Clinical chemistry principles, procedures and correlation, 10<sup>th</sup> ed. Lippincott Williams and wilkins, London: 314-320.

Mohamed HA, sayed IH, Mowad, M (2010) Protective effect of N. sativa seeds against dimetylaminoazobenzene induce liver carienogenesis 8;80-87.

Mott S, Yu L, Marcil M (2000). Decreased cellular cholesterol efflux is a common casue of familial hypoalpalipoproteinemia:role of the ABCAL gene Mutations.*Atherosclerosis* ;152:457-468.

Mustafa, A, E, (2002). Effect of *Nigella sativa* seeds on blood cholesterol level of rabbits, A thesis submitted in partial fulfillment for MSc degree in clinical chemistry sudan university of science and technology.

Najmi A, Haque SF, Khan RA, Nasiruddin M (2008). Therapeutic effect of *Nigella sativa* oil in different clinical and Biochemical parameters in metabolic syndrome. *Int. J. diabetes metabol*, 16:85-87.

Narain Vs, Gupta N, Sethi R , Puri A, Dwivedi SK ,Sarena R k, Puri VK (2008). Clinical correlation of multiple biomarkers for risk assessment in patients with acute coronary syndrome .*Indian Heart J. Nov.pec*;;60 (6)=536.42.

Nazrul Islam SK, Begum F, Ahsan T, Saiful Hage, MA (2004). Immunosuppressive and cytotoxic properties of *Nigella sativa*. *Phytotherapy Research* ; 18, 395-8.

Omer T, and Imtithal (2001). Biochemical immunological changes in rabbits blood as influenced by dietary black cumin (*Nigella sativa*) seeds. MSc thesis. University of Khartoum sudan.

Parasad K(1997). Dietary flax seeds in prevention of hypercholesterolemic atherosclerosis 132: 69-72 *Pharmacology*; Philadelphia : Saunders.PP.345.(SBN1-4166-2973-7.8<sup>th</sup> edition).

Plenz G, Robenek H (1998). Monocytes /macrophages in atherosclerosis. *Eur cytokine Netw*;9: 701-703. polyphenols. *J Med Food* ; 8:488-93.

Prasad K (2005). Hypcholesterolemic and antiatherosclerotic effect of flax lignan complex isolated from flaxseed. *Atherosclerosis*. 179 (2):269–275.

Pschierer V, Richter WO, Schwandt P (1995). Primary chylomicronemia in patients with severe familial hypertriglyceridemia responds to long-term treatment with (n-3) fatty acids. *J Nutr*;125:1490-1494.

Ramadan MF (2007). Nutritional value, functional properties and nutraceutical applications of black cumin (*Nigella sativa* L.): an overview. *Int. J. Food. Sci. Tech.* 42 (10): 1208–1218.

Reynisdottir S, Eriksson M, Angelin B (2003). Impaired activation of adipocyte lipolysis in familial combined hyperlipidemia. *J Clin Invest*;95: 2161-2169.

Ridgway ND, Byers DM, HW (1999). Integration of phospholipid and sterol metabolism in mammalian. *Prog Lipid Res*;38:337-360.

Rooney S and Ryan MF (2005). Effects of alpha-hederin and thymoquinone – a constituent of *Nigella sativa*, on human cancer cell lines. *Anticancer Res*; 25(38): 2197-2204.

Santamarina-Fojo S (1998). The familial chylomicronemia syndrome. *Endocrinol Metab Clin North Am* ; 27:551-567, viii.

Saad H, AZA, Zeh, H, Said, G (2005). Tradition a review: evidence based complementary alternative medicine (2) 474-9.

Schaefer EJ, Mc Namara JR (1997), Overview of the diagnosis and treatment of lipid disorders In: Rifai N, Warnick GR, editors Handbook of lipoprotein testing Washington, DC: AACC Press.

Segrest JP, Li L (2000), Amantharamaiah GM. Suture and function of apolipoprotein A-I and high – density lipoprotein. *Curr Opin Lipidol* ; 11:105-115.

Segui J , Gironella M, Sans Granell S, Gil F, Gimeno M, Coronel P, Pique JM, Panes J (September 2004). “Super oxide Dismutase ameliorates TNBS-induced colitis by reducing oxidative stress, adhesion molecule expression, and leukocyte recruitment into the inflamed” *J. Leukoc. Biol.* 76(3): 537-44.

Seo DM, Goldschmidt – Clermont PJ(2007). Unraveling the genetics of atherosclerosis: implications for diagnosis and treatment. *Exp Rev Mol Diagn*; 7:45-51.

Swamy SM, Tan BK.(2000). Cytotoxic and immunopotentiating effect of ethanolic extract of *Nigella sativa* L. seeds. *J Ethnopharmacol* (70): 1-7.

Tanne (1999). An egg a day is not harmful *British J. of Medicine* 7191-1094.

Topozada HH, Mazlou HA and El-Dakhakhny M (1965). The antibacterial properties of *Nigella sativa* seeds. Active principle of some clinical application. *J Egypt Med Ass* ; 48, Suppl. : 187-202. uses and pharmacological activities of *Nigella sativa*. *Pak. J. Biol.Sci.* 7 (4): 441-451.

Walden CC, Hegele RA, (1994). Apolipoprotein E in hyperlipidemia. L.

Warnick GR, McNamara JR, Boggess CN (2006) polyacrylamide gradient gel electrophoresis of lipoprotein subclasses. Clin lab Med ;26:802-846.

Worthea DR and Ghosh OA (1998). The in vitro anti tumor activity of some crude and purified components of black seed, *Nigella sativa*. Anticancer Res ; 18: 1527-32.

Yadav D, Pitchumoni CS (2003). Issues in hyperlipidemic pancreatitis. J Clin gastroenterol,36:54-62.

Zaoui A, Cherrah Y, Mahassini N, Alaoui K, Amarouch H, Hassar M (2002a). Acute and chronic toxicity of *Nigella sativa* fixed oil. Phytomedicine. 9(1): 69-74.