

5.4 References

1. Kollorz EK, Hahn DA, Linke R, Goecke TW, Hornegger J, Kuwert T. Quantification of thyroid volume using 3-D ultrasound imaging. *IEEE Trans Med Imaging* 2008; 27: 457–66.
2. Ghervan C. Thyroid and parathyroid ultrasound. *Med Ultrason* 2011; 13: 80–4.
3. Hegedüs L. Thyroid size determined by ultrasound: influence of physiological factors and non-thyroidal disease. *Dan Med Bull* 1990; 37: 249–63.
4. Knudsen N, Laurberg P, Perrild H, Bülow I, Ovesen L, Jørgensen T. Risk factors for goiter and thyroid nodules. *Thyroid* 2002; 12: 879–88.
5. Yildirim M, Dane S, Seven B. Morphological asymmetry in thyroid lobes, and sex and handedness differences in healthy young subjects. *Int J Neurosci* 2006; 116: 1173–9.
6. Gomez JM, Maravall FJ, Gomez N, Guma A, Soler J. Determinants of thyroid volume as measured by ultrasonography in healthy adults randomly selected. *ClinEndocrinol (Oxf)* 2000; 53: 629–34.
7. Wesche MF, Wiersinga WM, Smits NJ. Lean body mass as a determinant of thyroid size. *ClinEndocrinol (Oxf)* 1998; 48: 701–6.
8. Barrere X, Valeix P, Preziosi P, Bensimon M, Pelletier B, Galan P et al. Determinants of thyroid volume in healthy French adults participating in the SU.VI.MAX cohort. *ClinEndocrinol (Oxf)* 2000; 52: 273–8.
9. Rasmussen NG, Hornnes PJ, Hegedüs L. Ultrasonographically determined thyroid size in pregnancy and post partum: the goitrogenic effect of pregnancy. *Am J ObstetGynecol* 1989; 160: 1216–20.
10. Himanka E, Larsson L. Estimation of thyroid volume; an anatomic study of the correlation between the frontal silhouette and the volume of the gland. *ActaRadiol* 1955; 43: 125–31.
11. Van Isselt JW, de Klerk JM, van Rijk PP, van Gils APG, Polman LJ, Kamphuis C et al. Comparison of methods for thyroid volume estimation in patients with Graves' disease. *EurNucl Med Mol Imaging* 2003; 30: 525–31.
12. Nygaard B, Nygaard T, Court-Payen M, Jensen LI, Søre-Jensen P, Gerhard Nielsen K et al. Thyroid volume measured by ultrasonography and CT. *ActaRadiol* 2002; 43: 269–74.
13. Rago T, Bencivelli W, Scutari M, Di Cosmo C, Rizzo C, Berti P et al. The newly developed three-dimensional (3D) and twodimensional (2D) thyroid ultrasound are strongly correlated, but 2D overestimates thyroid volume in the presence of nodules. *J Endocrinol Invest* 2006; 29: 423–6.
14. Aksoy FG, Kesim Ö. Influence of cigarette smoking on thyroid gland volume: an ultrasonographic approach. *Turk J Med Sci* 2002; 32: 335–8.
15. Lucas KJ: Use of thyroid ultrasound volume In calculating radioactive iodine dose In hyperthyroidism. *Thyroid*, 2000; 10(2): 151–55
16. Ueda D: Normal volume of the thyroid gland in children. *J Clin Ultrasound*, 1990; 18: 455–62
17. Robins RJ, Schlumberger MJ: The evolving role of ¹³¹I for the treatment of differentiated thyroid carcinoma. *J Nucl Med*, 2005; 46: 28–37
18. Yousef M & Omer M, Evaluation of Normal Homogenous Thyroid Uptake by Using Tc99m, *Ozean Journal of Applied Sciences* 6(2), 2013

19. Ali W, Yousef M, Omer M, Gar- alnabi M, Ahmed M, Correlation of Thyroid Hormone Levels with Radioactive ^{99m}TcThyroid Uptakes, IOSR-JPBS, Volume 9, Issue 2 Ver. II (Mar-Apr. 2014), PP 165-169
20. Berman, M., HOFF, E., Barandes, M., Becker, D. V., SONENBERO, M., BENUA, R. and kotaras, D. A. : Iodine kinetics in man--a model. I. Clin. Endocrinol.28:1, 2014.
21. HINE, G. J. and LeoVINGER R. Application and radiation dose of radioisotopes in medicine. In Instrumentation in Nuclear Medicine, G. J. Hine, ed., Academic Press, New York, 1997, p. 626.
22. Brucer,M.: Thyroid radioiodine uptake measurement. A standard system for universal intercalibration. U.S. ABC Report ORINS-19, June, 1989.
23. Crespo G .G.and Vettreer ,H.: The calibration and standardization of thyroid radioiodine measurements, in tern. I. App1. Radiation isotopes 17:531, 1996.
24. Cameron, J. R. and BILL, D. B.: Thyroid uptake studies with ¹³¹I in very small doses. Radiology 79:452, 1992.
25. Berman, M., HOFF, E., Barandes, M., Becker, D. V.,SONENBERO, M., BENUA, R. and kotaras, D. A. : Iodine kinetics in man--a model. I. Clin. Endocrinol. 28:1, 1988.
26. Report of consultants of the International Atomic Energy Agency. The calibration and standardization of thyroid radioiodine uptake measurements. Phys. Med. Biol. 6:533, 1991.
27. A manual of radioactivity procedures. NCRP Report No. 28. National Bureau of Standards Handbook 80, U.S. Department of Commerce, Washington, D.C., 1961.
28. Alexander, W. D., Harden, R. MCG., and Shimmins J.,Nuc Med Tee 1966, Lancet, 2, 1041.
29. Andros, G., Harper, P. V., Lathrop, K. A., and Mc-Cardle, R. J. patho physiology of NM J. din. Endocr. Metab., 25, 1067. 1965.

30. Atkins, H. L., and Richards, P., General Tee of NM. J. nucl, Med., 9, 7. 1968,
Degrossi, O., Gotta, H., Olivari, A., Pecorini, V., and Chwojnik, A., Nud.-Med., 4, 383. 1995,
31. Sucker, H. J, and Cloud. Concise of human physiology, 2nd edition, chapter six page 212-
301, Country-company of publishing, 1989.