

6-2 المصادر الأجنبية.

Abbott, W. S. (1925). A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*, 18: 265-267.

Abd El-Ghany, N.M., Abdel-Razek, A.S., Ebadah, I.M.A. and Mahmoud, Y.A. (2016). Evaluation of some microbial agents, natural and chemical compounds for controlling tomato leaf miner, *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae). *Journal of plant protection research*. 56(4):372-379.

Abdel-Maksoud, N.; Zakhary, M. M.; Shehata, M. M.; El-Nowehi, A. M.; Grmaa, A. A. and Kelany, I. M. (1998). Effect of *Azadirachta indica* (Neem). against fruit fly *Bactrocera zonata* Saunders (Diptera: Tephritidae) under laboratory conditions *J. Applied Sci. Res.* 4 (2) 216- 223.

Abendroth. L; Emorei. R; Hartzler. B; McGrath. C; Mueller. D; Munkvold. G; Pope. R; Rice. M.E; Robertson. A; Sawyer. J; Schaefer. K; Tollefson. J; Tylka. G, (2009). A reference for identifying diseases, insect pests, and disorders of corn. Cooperative Extension Service, Iowa State University of Science and Technology, Ames, Iowa. 46-62.

Abul-Nasr, S. E. and A. K. El-Nahal, (1968). Some biological aspects of corn stem borer. *Sesamia cretica*, *Bull. Soc. Ent. Egypt.* 11: 429-444

Agaisse, H., and Lereclus, D. (1995). How does *Bacillus thuringiensis* produce so much insecticidal protein? Mini review. *Journal of Bacteriology*. 177, (21): 6027-6032.

- Ali, H.A.; Abdul – Russoul, M.S., and Swail, M.A.(1990).** Systematic list of Coccinellidae recorded for Iraq. Bul. Iraq nat. Hist. Mus. 8 (3) 45 – 51.
- Al-Jboori, I.M.S. Abdul-Rassol and S. J. Saleh. (2000).** New record of some biological enemies of citrus leaf miner *Phyllocnistis citrella* Stainton (Lepidoptera: Gracillariidae) in Iraq. accepted for publication in the Bull. Iraq. Nat. Hist. Mus.
- Allan, F. (2001).** Analysis of imidacloprid as a candidate for reduced-risk status. Issue of AE News. No. 186.
- Allsopp, P. G. and M, S. Sallam (2005).** prehardness for borer introduction into Australia. BSES, Aust, Johnvan Dyk 1996. 2: 93-115.
- Ambrose, M.L. (2003).** Characterization of the insecticidal properties of acetamiprid under field and laboratory condition. Thesis master. North Carolina state university. 71 pp.
- Amro, M. A. (2004).** Incidence of Certain Arthropod Pests and Predators Inhabiting Cowpea, With Special Reference to The Varietal Resistance of Selected Cultivars to *Bemisia tabaci* (GEN.) And *Tetranychus urticae* KOCH. Ass. Univ.Bull. Environ. Res. 7 (1):31-39.
- Andersh, W; Schwarz, M. (2003).** Clothianidin seed treatment the new technology of control of corn rootworm and secondary pests in u.s. corn protection Pflanzenchutz-Nachrichten Bayer 56(1): 147-172.
- Angus, T.A. (1954).** A bacterial toxin paralysing silkworm larvae. Nature (Lond), 173: 545-546. approaches. Kansas University, Manhattan, K S U. S. A. 423 pp.

- Arif. M.J. M.D. Gogi and G. Ahmed. (2004).** Role of Morpho-Physical Plant Factors Imparting Resistance in Cotton Against Thrips, *Thrips tabaci* Lind (Thripidae: Thysanoptera). Arab J. PL. Prot. 24:57-60.
- Asiye, D. and H, Kavut (2001).** Investigation on the attractive of some traps ageinest the *Osterinia nubilalis* (Hbn) moths in Aegan regian. Abstract of papers presented at the xxjst meeting in Venice, Italy.
- Askarianzadeh, A., K., Taher-Khani and A.R. Narrei. (2005).** Biological control of sugarcane pink stem borer, *Sesamia* spp. (Lepidoptera: Noctuidae) by the parasitoid 79 wasp *Platytesenomus hylas* (Hymenoptera: Scelionidae) in Iran. Silver Jubilee Congress, Guatemalla. January 30 - February 4, 2005. Biology commission. Abstracts of posters. (www.issct.org/BiologyAbstPoters2005.htm).
- Babu, G. R.; Krishnayya, P. V.; Rao, P. A. and Rao, V. S. (2008).** Field efficacy of *Bacillus thuringiensis* var. *kurstaki* in combination with plant oils on pest complex of cauliflower. Annals of Plant Protection Sciences, 16 (1): 53 57.
- Baker, T.C. (2011)** Insect Pheromones: Useful Lessons for Crustacean Pheromone Programs. Springer Science+Business Media PP 531 - 550.
- Barham, M. and Hajji, L. (2012).** Management of *Tuta absoluta* (Lepidoptera: Gelechiidae) with Insecticides on Tomatoes. Chapter 15. In: Insecticides – Pest engineering, Perveen, F. (ed.). pp: 333-354.
- Bekheit, H. K. M.; Moawad, G. M.; Elbedawy, R. A. and Mahgoub, M. M. (1997).** Control of potato tuber moth *P. operculella* in potato crop. Egyptian. J. Agric. Res. 75 (4): 923-938.

- Bozlagan, I., A. Ayvaz, F. Ozturk, L. Acik, M. Akbulut, and S. Yilmazi. (2010).** Detection of the *cry1* gene in *Bacillus thuringiensis* isolates from agricultural fields and their bioactivity against two stored product moth larvae. Turk. Jou. Agric., 34: 145-154.
- Bray, D. P.; Alves, G. B.; Dorval, M. E.; Brazil, R. P. and Hamilton, J. G. (2010).** Synthetic sex pheromone attracts the leishmaniasis vector *Lutzomyia longipalpis* to experimental chicken sheds treated with insecticide. Parasites & vectors, 3(1), 16
- Brian, M. A.G Porcar. F. and Yvan, R. (2009)** Effects of *Bacillus thuringiensis* δ -endotoxins on the pea aphid, *Acyrthosiphon pisum*. Appl. Environ. Microbiol. 75, 14: 4897-4900.
- Broughton, H. B.; Ley, S. V.; Slawin, A. M. Z.; Williams, D. J. and Morgan, E. D. (1986).** X-ray crystallographic structuredetermination of detigloyldihydro azadirachtin and reassignmentof the structure of the limonoid insect antifeedant azadirachtin.J. Chem. Soc. Chem. Commun., 2: 46-47.
- Brunner, J. F.; E. H. Beers (2005).** Role of neonicotinoide insecticides in Washington apple. Integrated pest managemnt control of Lepidoptera pest. J. of Insect. Science 5: 5-14.
- Burges, H.D. (2001).** *Bacillus thuringiensis* in pest control. retired from hosticulture research international. Jor. The royal of chemistry. 90-98 pp (U K)
- Butron, A., P. Soenagas, P. Revilla. 2001(a).** Effect of resistance to stem borer attacking fliut maize population. mission biological. of Galicia pantereda, Spain. cited by htt/agro. Sci 21: 125-130.

Butterworth, J. H. and Morgan, E. D. (1968). Isolation of a substance that suppresses feeding in locusts. *J. Chem. Soc.Chem. Commun.*, 1: 23-24.

Cabello T, Gallego JR, Fernández FJ, Soler A, Beltrán D, Parra A, Vila E (2009). The damsel bug *Nabis pseudoferus* (Hem.: Nabidae) as a new biological control agent of the South American Tomato Pinkworm, *Tuta absoluta* (Lep: Gelechiidae) in tomato crops of Spain. *OIBC/WPRS Bull* 49:219–223.

Cabello, T.; Gallego, J.R.; Fernandez-Maldonado, F.J.; Soler, A.; Beltran, D.; Parra, A. and Vila, E. (2013). The damsel bug *Nabis pseudoferus* (Hem.: Nabidae) as a new biological control agent of the South American tomato pinworm, *Tuta absoluta* (Lep.: Gelechiidae), in tomato crops of Spain. Meeting of the IOBC/WPRS Working Group-Integrated Control in Protected Crops, Mediterranean Climate. Mediterranean Agronomic Institute of Chania(MAICh), Crete, Greece.

Chailleux A, Biondi A, Lambion J, Zappala` L, Desneux N (2016). Indigenous natural enemies attacking *Tuta absoluta* (Lepidoptera: Gelechiidae) in Southern France. *Egypt J Biol Pest Control* 23:117–121.

Chancy, W.E.; E.T. Netwick, W.J. Bentley and N.C. Toscano. 2007. UC IPM Pest Management Guidelines. Cole Crops.Pub. No. 3442.PP4.

Charles, J.F.; Delecluse, A., and Nielsen –LeRoux, C. (2000) Entomo – pathogenic bacteria: from laboratory to field application. Kluwer Academic Publishers, Dordrecht, The Netherlands.

- Chen, Z.S. (1994).** Biological properties of *Bacillus thuringiensis*. International Course of *Bacillus thuringiensis*: Production" Wuhani P.R. China 10-25 /11/1994: 6-17
- Chiwada, P. and C. Owega. 2001.** Distribution of Durra stem borer. *Sesamia cretica* plant products distrectorate (C F I A) insect Sci, and It's Appli. 22: 470-478.
- CIMMYT (2000)** Wheat in the developing world.
http://www.Cimmyt.org/research/weat/map/developing_world/index.html.
- Cole, P. Horn, G.: Paul, A. (2006).** The impact of On *Micromus tasmainiae* (Walker) (Neuroptera: Hemerobidae) And the implication for pest control Lettce crops. Australian J. Of Entomol. 45: 244-248.
- Copland, J. J.; Kring, T. J. (2002).** Predaceous Coccinellidae in biological control. Ann. Rev. Entomol. 43: 295-321. Isikber, A. A. &, Obrycki M. J. W.2002. Effects of various aphid foods on *Cycloneda sanguinea*.Entomolgia Experimentalis at Applicata. 102, 93-97.
- Davidson, M.M.; Jacobs, J.M.E.; Reader, J.K. and Butler, R.C. (2002).** Development and evaluation of potatoes transgenic for a cry 1 Ac 9 gene conferring resistance to potato tuber moth. J. Amer. Soc. Horticul. Sin. 127(4): 590-596.
- De Angelis, J.D.; R.E. Berry and G.W. Krants. (1983).** A photosynthesis, leaf conductance and leaf chlorophyll content in spider mite (Acari: tetranychidae) injuries peppermint leaves. Environ. Entomol, 12:345-349.

- Demirbağ, Z., R. Nalçacıoğlu, H. Katı, İ. Demir, K. Sezen & Ö. Ertürk, (2008).** Entomopatojenler ve Biyolojik Mücadele. Esen Ofset Matbaacılık, Trabzon, 325 s.
- Dharmananda, S. (2004).** Matrine and Oxymatren: Subject of Chinese research. www.itmonline.org 4pp.
- Dimova, R. and A. Dekov. (1990).** Field crop of tropic and semitropic area. p:432.
- Dinarvana, Raajabpour, zandi sohani and Farhari (2020)** Effect of weedy culture on population densities, spatial distributions and sampling procedures of *Spodoptera exigua* and *Sesamia cretica* (Lep., Noctuidae) in corn fields. 2020 Feb;110(1):84-95. doi: 10.1017/S0007485319000312. Epub 2019 Jun 13.
- Du Pont.(2005).** Avaunt Insecticide "A Growing Partnership With Nature", H-64922. Du Pont, Wilmington, Delaware 19898. pp. 8.
- Edgington S, Moore D, Elbouhssini M, Sayyadi Z (2007)** Beauveria bassiana for control of sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae) and aspects of the insect's daily activity relevant to a mycoinsecticide. Biocontrol Science and Technology 17(1-2): 63-79.
- Ehlers, R.U. (2006).** Current and future use of nematodes in biocontrol: practice and commercial aspects with regard to regulatory Police issues. Biocont. Sci. Techn., 6, 303–316.
- Ekesi, S., N. K. Maniania, and S. A. Lux. (2002).** Mortality in three African tephritid fruit fly puparia and adults caused by the entomopathogenic fungi *Metarrhizium anisopliae* and *Beauveriabassiana*. Biocontrol Sci. Technol. 12: 719-729.

Elamin, E, M. (1988) (b). same as pests of the ecology of the stem borer *Sesamia cretica*. in sugar cane and sorghum in the sudan Beitrage - zur tropis. chen. land wirts chaft. and veterina remdizin (German D. R.) 26:33-37.

El-Aw, M.A; Draz, K.A.A; Hashem, A.G and El-Gendy, I.R. (2008). Mortality comparsion among spinosad –Actara, Malathionand Methomyl containing baitsentomopathogenic isolates of *Bacillus thuringiensis* on the fertility and reproductive system of the female mothe of cotton leaf worm *Spodoptera littoralis* (Boisp). Ain Shams Univ.Sci. Rep.

Elbert, A.; B. Becker; J. Hartwig and C. Erdelen. (1991). Imidacloprid a new systemic insecticide. Pflanzenschutz Nachrichten Bayer 44:113-116.

Elekcoglu Z. and Uygun N. (2006). Institute of plant protection Adana. Turkey. 2005. The parasitoid complex of citrus leafminer, *phylloncistis citrella stainton* (Lepidoptera: Gracillariidae) in the East Mediterranean region of Turkey and their role in biological control, Turk J. Zool. 30: 155-160.

Elimi, M. A.; M. A. Monzer and Y. A. A. EI–Deeb. (2006). Evaluation of two commericial sexpheromone lures for monitoring males of pink boll worm•*Pectinophora gossypiella* and spiny bollworm•*Earias insulana* in Egypt. J. Biol. P. cont 12 (1). 2002.

ElrlandsonS, Moore D, Elbouhssini M, Sayyadi Z (2007) Beauveria bassiana for control of sunn pest, *Eurygaster integriceps* (Hemiptera: Scutelleridae) and aspects of the insect' s daily activity relevant to a mycoinsecticide. Biocontrol Science and Technology 17(1-2): 63-79.

Environmental Protection Agency (EPA). (1998). EPA. Reregistration Eligibility Decision (RED) *Bacillus thuringiensis* US.EPA, prevention pesticides and toxic substance, EPA 738-R-98-004, 19P.

Ernesto, C. B.; Nestor, B. M.; Jorge V. G. Hugo C. A. B. and Antionio H. P. (2004). Instituto Nacional Autonomo de Investigation Agropecuaria Casilla Postal 100 Estacion Experimental Portoviejo – Ecuador. Florida Entomologist 87 (1).

EUROPEAN FOOD SAFETY AUTHORITY 2013- Conclusion on the peer review of the pesticide risk assessment of the active substance aminopyralid. EFSA Journal 11, 3031–3075.

Ezzeldin.H. A: Sallam. A. A: Helal. T. Y: Fouad. H. A. (2009). Effect of some materials in sesamia cretica infesting some maize and sorghum varieties. Archives of phytopathology and plant protection, 42 (3), P: 277 – 290.

Fagan, L.L., McLachlan. A., Till C.M. and Walker M.K. (2009). Synergy between chemical and biological control in the IPM of currant-lettuce aphid (*Nasonovia ribisnigri*) in Canterbury, New Zealand. Bull Entomol Res. Jul 2:1-7.

FAO. (2004). Chlief publishing sevice information. Ddivision, viale delle termedi-Italy.10pp.

FAO. (2006). FAOSTAT, statistical database.<http://www.FAO.org>.

FAO. (2012). Food and Agriculture Organisation. FAOSTAT. FAO static. Division, Rome.

FAO/IAEA/USDA. (2014). Product quality control for sterile mass-reared and released tephritid fruit flies. Version 6.0. Vienna, International Atomic Energy Agency. 164 pp.

- Faragalla and Ibrahim. (1990).** A Preliminary Response of Exotic Vs. Local Corn Hybrids to Natural Infestation of Two Stem Borers, (LEPIDOPTERA: NOCTUIDAE). Journal: Journal of King Abdulaziz, University: Science. ISSN 1319-1012. Volume: 2; Issue: 1; Start page: 79. 1990.
- Fargues, J.; Ouedraogo, A.; Goettel, M. and Lomer, C. (1997).** Effects of temperature, humidity and inoculation Method on susceptibility of *Schistocerca gregaria* to *Metarhizium flavoviride*. Biocont. Sci. Technol. 7: 345-356.
- Faust, P.G. (1972).** The δ -endotoxin of *Bacillus thuringiensis* DI. A rapid method for separating parasporal bodies from spores. Journal of Invertebrate Pathology, 20: 130-149
- Federic, B.A.; Luthy, P. and Ibarra. (1990).** Parasporal body of *Bacillus thuringiensis israelensis*. In Bacterial Control of Mosquitoes and Blackflies, Rutgers University Press. pp: 16-44.
- Fluence Ltd. Company China. (2006).** Oxymatrine 2.4 EC. Company profile .4pp.
- Franja, M. and mustea, D. 2001.** Reducing possibility of environment poll. through utilization one of biological methods. J, Eco, Entomol. 224(2):3131-3137.
- Getu, E., W.A. Overhoh and E. Kairu. (2001).** Distribution and species composition of stemborers and their natural enemies in maize and sorghum in Ethiopia. Insect Science 21: 353-359.

Ghribi, D.; L. Abdelkefi-Mesrati; H. Boukedi; M. Elieuch; S. Ellouze-Chaabouniand S. Tounsi. (2011). The impact of the *Bacillus subtilis* SPB1 biosurfactant on the midgut histology of *Spodoptera littoralis* (Lepidoptera: Noctuidae) and determination of its putative receptor.J. Invertebr. pathol.109(2):183-186.

Giustolin, T.A.; Vendramim, J.D.; Alves, S.B.; Vieira, S.A.; Pereira, R.M. (2001). Susceptibility of *Tuta absoluta* (Meyrick) (Lep. Gelechiidae) reared on two species of *Lycopersicon* to *Bacillus thuringiensis* var. *kurstaki*. J. Appl. Entomol. 125: 551–556.

Glare, T. R. and O'Callaghan, M. (2000). *Bacillus thuringiensis*: Biology, Ecology and Safety. John Wiley and Sons, Ltd. UK. pp. 350.

Gonzalez-Cabrera, J., Molla, O., Monto', H. and Urbaneja, A. (2011). Efficacy of *Bacillus thuringiensis*(Berliner) in controlling the tomatoborer, *Tutaabsoluta* (Meyrick) (Lepidoptera: Gelechiidae). BioControl .56:71–80.

Gorashi, N.E.; Tripathi, M.; Kalia, V.and Gujar, G.T. (2014) Identification and characterization of the Sudanese *Bacillus thuringiensis* Strain for their efficacy against *Helicoverpa armigera* and *Tribolium castaneum*. India journal of experimental biology,52:637-649.

Gordon, R.D. (1985). The Coccinellidae (Cdeoptera) of mrica north of mexico. J. Ny Entomol. Soc. 93: 1 – 912.

Gutierrez, C, P. and C, P. Torres. (1986). Wound induced chang in DIMBOA concentration in maize plant scausd to *Sesamia nanogrioides*. Annals of App. Biol. 113: 25-34.

- Hafez, M.; F. N Zaki; A, Moursy and Sabbour, M. (1994).** Biological effects of the Entomopathogenic fungus, *Beauveria bassiana* on the potato tuber moth *Phthorimaea operculella* (Seller). J. Islamic Academy of Sciences, 7(4): 1 – 4
- Halawa, Safaa. M. (2006),** Evaluation of some plant extracts to control the pink stem borer *Sesamia cretica* (Led.), Annals of Agric, Sc, Moshtohor, 44 (4): 1946 – 1946.
- Heimpel, A.M. (1967).** A critical review of *Bacillus thuringiensis* var. *thuringiensis* Berliner and other crystalliferous bacteria. Annu. Rev. Entomol., 12: 287-322.
- Hela, E. (1986).** Ulta structure of egg membrances of newly deposited egg of pink borer *Scesamia cretica*, Alexandrai J. of Agric. Sci. 31(3): 329-345.
- Henderson, C.F. and E.W. Telton. (2007).** Test with acaricides against the brown wheat mite. J. Econ. Entomol. 48: 157-161.
- Hodek, I. (1973).** Biology of Coccinellidae. Academia czechosl – ovakia, Prague. 260 pp.
- Höftele, H. and Whiteley, H.R. (1989).** Insecticidal crystal proteins of *Bacillus thuringiensis*. *Microb. Rev.*, 53: 242-255.
- Höftele, H. and Whiteley, H.R. (1989).** Insecticidal crystal proteins of *Bacillus thuringiensis*. *Microb. Rev.*, 53: 242-255.
- Holbrook, F.R. (1977).** Aldicarb and thiofanox effect on the The feeding activity of green peach aphids. J. of. Econ. Entomol. 7 (6); 742-744.

Hosary, Rasha A. EL. (2011). Evaaluuation of some essential oils against sesamia called. under field condtions. Plant Protection Dept., Fac. of Agric., Benha University, Egypt. Journal of American Science. 7, (3).

Huang, F. Subramanyam, B., Towes, T.D. (2004). Susceptibility of laboratory and field strains of four stored product insect species to spinosad. J. Econ. Entomol., 97(6): 2154-2159.

Hwan kim, Hyeong, Sung Hun Youn, Taek Su Shin, (2012). Biological control potentials of insect-parasitic nematode *Rhabditis blumi* (Nematoda: Rhabditida) for major cruciferous vegetable insect pests. Applied Entomology and Zoology. 47, 389 – 397, Cite this article.

Ismail, H. (2002). Field observation on the oviposition behaviaer of different corn borers and estimation of economic threshold _Acta phytopathological. Entomol. Hungariea 24:111-116.

Jankevica, L. (2004). Ecological association between entomopathogenic fungi in pest insects recorded in Latvia. Entomology, 41: 60-65.

Jepson, W. F. (1954). Aoritical review of the world literature on the Lepidopterous stalk borers of tropical graminaceous crops. (cited by ph. D. Thesis, Al exandria University.)167 pp.

Johovan, D. (1996). How corn is damaged by the ECB, north centrol regian Extension public. NCR. 327. Cited by <http://1astate.edu/corn borer/htm>.

Kaaya, GP. and S. Hassan. (2000). Entomogenous fungi as promising biopesticides for Stick control. *Experimental and Applied Acarology* 24: 913–926.Ekesi, S., N. K.

- Kagan, M. and E. E. Ortman. (1978).** Antixenosis a new term proposed to define painter's non-preference modality of resistance. Bull. Entomol. soc. Amer 24: 175-176.
- Karus, W.; Bokel, M.; Klenk, A. and Pohul, H. (1985).** The structure of azadirachtin and 22, 23-dihydro-23 b-methoxyazadirachtin. Tetrahedron Lett., 26: 6435-6438.
- Kauma, M T.; S. Kyamanywa; J. A. Ogwang; O. C. omwege, and H. R. willson. (2001).** Cereal stem borer species complex and establishment of *Cotesia flavipes* Cameron in eastern Uganda. Insect Science and its Application. 21(4):317-325.
- Kennedy, G.G. (2007).** Resistance in tomato and other Lycopersicon species to insects and mite pests. Improve solanceous crops.2:488-519.
- Koul, O. & S. Wahab. (2004).** Neem: Today and in The New Millenium.Kluwer AcademicPublishers New York, Boston, Dordrecht, London, Moscow, p.291.
- Kumar, H and. G. O. Asino. (1996).** Resistance of maize to *Chilopartellus* effect of plant phenology. J. Econo. Entomol. 86: 969-973.
- Kumar, L. V., A. Prabhuraj, A. K. Chakravarthy. (2006).** Management of sorghum shoot bug, *Peregrinus maidis* (Ashmead) (Homoptera: Delphacidae) through seed treatment chemicals. In: Goel SC, ed. Advances in Indian Entomology: Productivity and Health (a Silver Jubilee, Supplement No. 3, Volume II; Insect and Environment. Uttar Pradesh Zoological Society, Muzaffarnagar. pp. 137–140.

Kumpmann, S., J. Kienzle, C. Schulz, C. P. W. Zebitz, A. Trapp & M.

Kelderer, (2002). "Control of *Adoxophyes orana* F.v.R. with entomopathogens and NeemAzal-T/S: first approach for optimal combination strategies. 10th International Conference on Cultivation Technique and Phytopathological Problems in Organic Fruit Growing and Viticulture Proceedings of a Conference, Weinsberg. laboratory and field strains of four stored product insect species to . 42-35

Lacey, A.L.; E. Riga; and W. Snyder. (2004). The potential for using insect specific pathogens for control of insect pest of potato. Journal of potato progress. vol. IV. no.1.

Leyden, P. (2014). Efficacy of *Bacillus thuringiensis* spvay applications for control of lepidoptera pests. Master thesis. potchcfstoom campus of North west Lniversity. Africa south .49 pp.

Lezama-Gutierrez, R.; Hamm.J.J.; Molina-Ochoa, J.; Lopez-EdwardPescador- Rubio, A.; Gonzales-Ramirez, M.and Styer, E. L. (2001). Occurrence of entomopathologen of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) in the Mexican states of Michoacan, Colima, Jalisco and Tamaulipas. Florida Entomol.84(1):24-30.

Lozzia, G. C. and Barbara, M. (2003) Susceptibility of *Ostrinia nubilalis* to *Bacillus thuringiensis* var. *kurstaki*. Bulletin of Insectology 56 (2): 215-219.

Madian S. Kasem, Sameer J. Assaf, Ahmad J. Ibrahim, Mohamed

Ibrahim, Omran Yousef and Haleem Yousef. (2006). the fieldefficacy of thiamethoxam and imidacloprid as assd treatment compared to spray application to control some insect pests in faba bean, wheat and corn in Syria. ninth arab congress of Plant Protection, damascus, Syria. 19-23 november

Maiens fisch, P.H.; Huertimann .H; Rindisbacher. A; Gsell.L Dettwiler. H; Haettenschwiler. J.; Syeger.E.and Walti.M (2001). The discovery of thiamethoxam asecond generation neonicotiniod. pest Manage. sci. 57; 165-176.

Malvar. R.A; Butron. A; Alvarezi. A; Padilla. G; Cartea. M.E; Revilla. P; Ordas. A, (2007). Yield performance of the European Union Maize Landrace Core Collection under multiple corn borer infestations, 26(5): 775-78.

Mao Moore, K.C.and Erlandson, M.A. (1990). Isolation of Aspergillus parasitic spear and *Beauveria bassiana* Vuillemin melanopline, grasshoppers and demonstration of there pathogenicity in *Melanoplus Sanguinipes*. 120(10):989-991.

Martinez, S.S. (2002). O nim *Azadirachta indica*: natureza, usos múltiplos, produção. Londrina: Instituto Agronômico do Paraná. 142 pp.

Masaad Zolfagharian, Asaeedizadeh and Habib Abbasipour. (2016). Efficacy of Two Entomopathogenic Nematode Species as Potential Biocontrol Agents against the, Diamondback, Moth, *Plutella, xylostella*, (L.) DOI:10.18641/jbc/30/2/88697Corpus ID: 88861389.

- Mesbah, H, A, Mourad, A, K. El-Nimr H. M, Massoud, M. A. and A, A. Abd El-Aziz. (2002).** The role of some agro-practices and fertilizer type on both the incidence of stem borers infestation and corn yield in Egypt. Biolo. weten schappen (Belgium)p. 67 :575-589.
- Mesbah, H, A. and A, A. Gaber. (2002).** The effect of sequential apply of foliar nutrients biofertilizers and sawing dates on the incidence of corn stem borer. Egypt (Bel) 67:407-497.
- Metcalf, R. L., and W.H. Luckmann. (1975).** Introduction to Insect pest management-A witey. Intersci pub. New yourk. 587pp.
- Michaud JP, Mccoy CW and Futch SH (June 2002).** Lady beetle as biological control agents in citrus. EDIS. <http://edis.ifas.ufl.edu/HS138> (10 september 2002).
- Mirkarimi, A. (1988).** Biological studies on sorghum borer, Sesamia cretica Led. (Lepidoptera: Noctuidae). Iranian J. of Agri. Sc. (Iran) 18: 17-36. 20- Moyal, p.; M.M.
- Mohammed A.S.A. (2009).** A study of characters contributing to yield in some genotypes of maize J. of Tik. Univ. for Agri. (5) 2: 1 – 9.
- Mohammed B. Alhussein, Sara A.A. Kehail and Hashim A. Mohamed, (2014).** maize research program, arc, wad medani, sudan maize research Program, arc, new halfa, sudan the agricultural research corporation center of crop protection the 3rd pest management conference of the suoan, arc, wad medani, sudan, february 3rd – 4th

Monro, H.A. U. (1961). The dura stem borer sesamia cretica. Led. A, new problem in imported European room corn. canad. Ent (cited by ph.D. Thesis Elexandria University)190 – 180 :(10 – 9) 79 .

Monro, R.E. (1961). Protein Turnover and the Formation of Protein Inclusions during Sporulation of *Bacillus thuringiensis* Biochem. J. 81:225-232.

Mushtaha, R.M. (2013). Characterization and Bioassay of Different Commercial Products of *B. thuringiensis* Against Four Larval Stages and Adults of an insect *Tutaabsoluta* in Laboratory. Master thesis. The Islamic University –Gaza. Science college.154-166

MYERS, S. AND WEDBERG, J.1999. Development of an Economic injury level for European corn borer (Lepidoptera: Pyralidae) on corn grown for silage, Journal of Economic Entomology, Vol. 92, No. 3, 1999, 624- 630.

Navon A. (1993). Control of lepidopteran pests with *Bacillus thuringiensis*. in *Bacillus thuringiensis*, an environmental biopesticide: theory and practice.eds Entwistle P. F., Cory J. S., Bailey M. J., Higgs S. (John Wiley and Sons, Chichester, England), pp 125–146.

Niedmann, L.L. and Meza-Basso, L. (2006). Evaluació'n de cepas nativas de *Bacillus thuringiensis* como una alternativa de manejo integrado de la polilla del tomate (*Tuta absoluta* Meyrick; Lepidoptera: Gelechiidae) en Chile. Agric Te'c 66:235–246.

- Nithiskarani, Anita, and Vetrivelkalai, Selena Nelson. (2019).** Biocontrol potential of entomopathogenic Nematodes, Steinernema glaseri (Steiner, 1929) and Heterorhabditis indica (Poinar, Karunakar & David, 1992) against brinjal ash weevil (*Myllocerus subfasciatus*). Published 2019. Biology.
- Niyibigira, E. I.; Z. S. Abdallah; W. A. Overholt; V. Y. Lada, and A. Vanhuis. (2001).** Distribution and abundance, in maize and Sorghum, of Lepidopteran stem borers and associated indigenous parasitoids in Zanzibar. Insect Science and its Application. 21(4): 335-346.
- O'Brien, K. P., S. Franjevic & J. Jones. (2009).** Green Chemistry and Sustainable Agriculture the Role of Biopesticides. Barbara Smith Fund, Kendeda Sustainability Fund and the Johnson Family Foundation, pp.55.
- Oliveira a Jr., J. D. A.; Jain, S.; de, de SouzC. M. F.; Ayres, C. F. and Lucena, W. A. (2009).** Toxicity of a *Bacillus thuringiensis* israelensis – like strain against *Spodoptera frugiperda*. Bio. Control., 54: 467 – 473.
- Osman, H.H.; Badr El-Sabah, A.F.and Abeer, M.M. (2014).** The potency of Chloropyrifos and Camphor extract on *Spodoptera littoralis* (Boisd.). Acad.J.biolog.Sci.,5(2) :131-139.
- Ozdemir, N. 1988.** Corn borers *Ostrinia nubilalis* and *Sesamia cretica* Led. in the black sea region of Turkey. Symposium on corn borers and control measures proceedings. Pp. 25-26
- Padmaja, V. and G.Kaur. (2001).** Use of the fungus *Beauveria bassiana* (Bals). Vuill (Moniliales: Deuteromycetes) for controlling termites. *J. CurrentSci* .18 (6):645-647.

- Panhwar, F.(2005).** The Neem tree *Azadirachin indica*, the natural pesticide practice in Pakistan. Chem.lin-virtual labrotary chemistry. J. econ. Entomo.81(3):17-21.
- Pathak, M. D. and R, C. sexena. (1979).** Insect resistance pp:220-292. in plant breeding perspectives (Edt) J, and Hendriksen, A. J. Center for Agric. publishing and Documentation, Wageningen 460 pp.
- Pathak, P.H and Krishna, S. S (1991).** Postembryonic development and reproduction in *Corcyra cephalonica* (Lepidoptera) on exposure Eucalyptus and neem.J.Chem.Ecol.17(12):2553-2558. (abst).
- Persley, G.T(1996)**Biotechnology and integrated pest management . CAB. International, Washington, USA, 51 – 283.
- Prabal, L., R. Shivay, and I. pandey . 2000.** Metting world maize needs Technological opportunites and prioties for public sector. cited by ent . Google htta:/ Aesop. Rutgers. edu/ 12. htm.
- Qazzaz, F.O.; Al-Masri, M. I. and Barakat, R. M.(2015).** Effectiveness of *Beauveria bassiana* native isolates in the biological control of the mediterranean fruit fly (*Ceratitis capitata*). Advances in Entomology 3: 44-55.
- Ramanaidu, K. and G. CH. Cutler. (2012).** Different toxic and hermetic responses of *Bombus impatiens* to *Beauveria bassiana* *Bacillus subtilis* and spirotetramat. © Society of Chemical Industry.

Ricano, J., B. Guerri-Agullo, M. J. Serna-Sarrias, G. Rubio-Llorca, L.

Asensio, P. Barranco and V. Lopez-llorcal, (2013). Evaluation of the pathogenicity of multiple isolates of *Beauveria bassiana* (Hypocreales: Clavicipitaceae) on, *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae) for the assessment of asolid formulation under simulated field conditions. -Fla. Entomol. 96: 1311-1324.

Rochat, D.; Mohammadpoor, K.; Malosse, C.; Avand-Faghih, A.; Lettere, M. Beauhaire, J.; Morin, J.P.; Pezier, A.; Renou, M. and Abdollahi, G. A. (2004). Male aggregation pheromone of date palm fruit stalk borer *Oryctes elegans*.J. Chem. Ecol. 30:387-407.

Rosell,G., C. Quero, J. Coll & A. Guerrero. (2008). Biorational insecticides in pest management. The Journal of Pesticide Science, 33(2): 103–121.

Royer, L. and J. N. McNeil. (2004). Changes in calling behavior and mating success of the European corn borer, *Ostrinia nubilalis*, caused by relative humidity. Entomologia. Ex. Applicata vol.61 (2):131-138.

Saad, A.F.(1979). Arial spraying for control of *Sesamia cretica* in Iraq. (Cited. Rev. Appl. Ent.) 67(9):211-215.

Sabbour, M.M.(2002). The role of chemical additives in enhancing the efficacy of *Beauveria bassiana* and *Metarrhizium anisopliae* against the potato tuber moth *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). Pakistan J.of Biological Science. 5(11): 1155-1159.

- Saha S., S. Walia & B.S. Parmar .(2011).** Exploring the diversity of neem bioactives as eco-benign pesticides: A reappraisal. *Toxicological Environmental Chemistry*, 93: 1508-1546.
- Sajan ,S, S. and S, S. Sekhvon. 1987.** Antibiosis in maize stem borer *Chilopartellus* (Swinboe)in India- Tropical pest management 33:55-60.
- Samson, A.R.; Evans, D. and Latge, J. (1988).** Atlas of entomopathogenic-fungi. printed in the Netherland, New York . 187 pp.
- Samson, M. K.; R. C Brakke. (1981).** Effect of temperature on susceptibility of normal and aberrant ratio corn stocks to barley stripe mosaic and wheat streak mosaic virus. *Phytopathology*, 71: 823-824.
- Sarwar, M (2009).** Population synchronization of Aphids (coleoptera: coccinellidae) and exploitation of food, food attractants for predator. *J. Biological Diversity and conservation* 2(2):85-89.
- SAS Institute Inc. SAS state guide for personal computer. Version 6 ed. SAS Institute, Cary, NC. USA. (2010).** Abbott, W. S .1925. Method for computing the effectiveness of an insecticide. *J. Econ. Enstool.* 18: 265 – 267.
- Schmutterer, H. (2002).** The Neem Tree , Azadirachtaindica A .Juss Other Meliaceous plants Sources of Unique Natural products for Integrated pest management medicine and other purposes .Mumbia , India.

- Schnepf, E.; Crickmore, N.; Van Rie, J.; Lereclus, D.; Baum, J.; Feitelson, J.; Zeigler, D.R.; Fean, D.H. (1998).** *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiol. Mol. Biol. Rev.* 62: 775–806.
- Scholte, E.J.; Knols BGJ; Samson R.A.and, Takken, W.(2004).** Entomopathogenic fungi for mosquito control : A review. *J. Insect Sci.*, 24 pp.
- Sebesta, K.; Farkas, J.; Horska, K. and Vankova, J.(1981).** Thuringiensin, the beta-exotoxin of *Bacillus thuringiensis*. In *Microbial Control of Pests and Plant Diseases 1970-1980* (Ed. H.D. Burges), Academic Press, London, New York, pp: 249-281.
- Senthil Nathan, S. S., K. Kalaivani & K. Murugan. (2006).** Behavioural responses and changes in biology of rice leaffolder following treatment with a combination of bacterial toxins and botanical insecticides. *Chemosphere*.
- Senthil, K. P. and P. Bharthi. 2009.** Studies on relationship between gca and sca effects in maize (*Z.mays L.*), *Elect. J. Plant Breed.*, 1:24-27.
- Seraj.A.A.(2001).** Damage and Assessment of Losses Caused on Sugar cane by *Sesamia, cretica* (Lepidoptera: Noctuidae)inKhuzestanRegion. *Journal. Journal of Science and Technology of Agriculture and Natural Resources.* ISSN 1028-7655. Volume: 5; Issue: 2; Start page: 169.
- Shah, P.A . ,and M.S . Goettel . (1999) .** Directory of microbial control products and services, 2 nd edn. Division on microbial Control. Society, for invertebrate Pathology. Division on microbial Control, Gainesville, USA. P. 81.

Shalaby, F. F., M. M. Assar, A. F. Lutfallah, and S. S. Yacoub. (2013).

Field Application of Onion preparations for Controlling *Sesamia cretica* Led. Infesting Maize Plants. Workshop Cairo. Egypt. February 10-11.

Shutong, W.H. ; Tongle, Z. F. and Forrer, H. R. (2007). Screening for plant extracts to control potato late blight. Frontiers of Agriculture in china, 1:43-46.

Smith.C.M. (2005). Plant resistance to Arthropods, molecular and conventional.*thuringiensis*. *Microb. Rev.*, 53: 242-255.

Soares, M. A., Leite, G. L. D., Zanuncic, J. C., Rocha, S. L., SÁ, V. G. M., and Serrao, J. E. (2007). Flight capacity, parasitism and emergence of six 6 Trichogramma (Hymenoptera: Trichogrammatidae) species from reforested areas with eucalyptus in Brazil. *Phytoparasitica* 35: 314-318.

Sohati, P. H.;E. M. Musonda, and M. Mukanga. (2001). Distribution of cereal stem borers in Zambia and release of *Cotesia flavipes* Cameron, an exotic natural enemies of *Chilo partellus* (Swinhoe). *Insect Science and its Application*. 21 (4):311-316.

Sridhar, p. (2006). Plant Resistance to Insects citd by Googl's <http://11aesop.rutgers.edu/~11lecture12.htm> as retrieved 21=46-57.

Tamaki, G.; Weeks, R. E., (1981). The impact of predators on population of green peach aphids on field grown sugar beets. *Envirn. Entomol.*, 2: 345-349.

- Tams, W, T. and Bowden, J. (1953).** A revision Africa species of Sesamia guene and related genera . Bul. of Entomol. Research. 43:645-678. Doi :10 .1002/ejlt .200900138the chemical industry". European Journal of Lipid Science and *toosendan* (Meliaceae). Pestic. Biochem. Physiol. 53, 34–41.
- Teander, O. (1985).** Review of straw carbohydrate in progress in biotechnology 1. new approaches to resistance on cereal carbohydrates . H. 11. R. Dandmnnek, L (Edts) pp:217-230. Elesvier sci, public. Amsterdam.
- Temerak, A.A., S.A. and Sayed M.L. Lyasandrou. (2010).** The use of different insect control regimes using three green chemicals to combat Viracola livia on date palm fruit in Egypt. Furth international date palm conference. Abu Dhabi, United Arab Emirates,15-17 March 2010.
- Thiery and Frachon, E. (1997).** Identification, Isolation, culture and preservation of entomopathogenic bacteria. In: Manual of Techniques in Insect Pathology, Academic Press, pp: 54-77
- Thomas, Y. M. T. Betherod, and I. Pelzuelo . (2003).** American of the ECB *Ostrinia nubilalis* 1-sex pheromone Abbreviated Evolution 57:261-273.
- Thomason, J.A., B. E. Hartley, J. D. Gibson, R. Sui, and S. W. Searcy. (2009).** Characteristics of High-Biomass Sorghum as a Biofuel American Society of Agri. and Biological Engineers, St. Joseph, Michigan Paper No. 090015.

- Torres, J. B and Ruberson, J. R. (2004).** Toxicity of Thiamehoxam and Imidaclorpid to *Podisus nigrispint* (Dallas)(Heteroptera :Pentomidae) nymphs associated to Aphid and white fly control In cotton . Neotropical entomology . 33(1): 99-109.
- Trindade, R.C.P.; Marques, I.M.R.; Xavier, H.S. and Oliveira, J.V. (2000).** Extrato metanólico da amêndoа da semente de nim e a mortalidade de ovos e lagartas da traça-dotomateiro. Scientia Agricola, Piracicaba, Vol. 57, No. 3, P. 407-413.
- UCIPM Davis, R. M. (2003).** Maize dwarf mosaic virus. How to manage pests Pest Management Guidelines:Corn. UC ANR Publication 3443. <Http://A: 113101011. htnl>.
- Ujvary ,I. (2001).** Pest Control Agents from Natural Products. In handbook of pesticide toxicology, 23rd (Editör: R. I. Krieger) Academic Press. San Diego, 109-179.
- USDA-ERS. 2012. 5 July , (2012),** posting date. Adoption of genetically engi – neered crops in the U.S.: recent trends in GE adoption. USDA,Washington,DC.<http://www.ers.usda.gov/data-products/adoption-of-genetically-engineered-crops-in-the-us/recent-trends-in-ge-adoption.aspx>.140 – 176.
- Van Deventer, P.(2009).** Leafminer threatens tomato growing in EuropePages 10-12 in Agri-& Horti World, Fruit & Veg Tech.
- Waghmare J.T., A.M. Ware & S.A. Momin.(2007).** Neem oil as pesticide. *Journal of Dispersion Science and Technology*, 28: 323-328.

- Ward, M. P. (2001).** Effectiveness of a synthetic lure to reduce blow fly strike incidence: preliminary observations. *Vet parasitol.* 97:77 – 82. cited by www.rspca.org.au.
- Welty, C. (2001).** Insecticide News. *VegNet.* 8(3) February.
- Wessner, M, Champion B, Girault J-P, Kaouadji N, Saidi B and LafontR (1992).** Ecdysteroids from *Ajuga iva*. *Phytochemistry*, 31: 3785-3788.
- Whalon, M. E. & B. A. Wingerd. (2003).** Bt: mode of action and use. *Archives of Insect Biochemistry and Physiology*, 54: 200-211.
- White, P. J. and L. A. Johnson. (2003).** Corn: Chemistry and Technology.2nd Edn., *American Association of Cereal Chemists*, St. Paul, MN., USA., ISBN-13: 9781891127335, PP: 892.
- Wiltshire , E . P . (1948).** The Lepidoptera of the kingdom of Egypt. *Bull. Soc. fonadler. Entom.* xxxII: 203 – 298. www.itmonline.org. 4pp.
- Wyatt, R. D., P.B. Hamilton, and H. R. Burmaster. (1973).** The effects of T-2 toxin in broiller chickens. *Poultry Sci.* 52:1853-1859.
- Wyatt, T.D. (2003)** Pheromones and Animal Behavior Chemical Signals and Signatures .second edition . cambrige university press . pp 66.
- Yaqti, R., C. B. meister, M. W. Idraw, and E. Al-jouri.(2007).** Effect of NeemAzal T/S and Aqueous Extract for Mature Fruits (*Melia azedarach* L.) in Controlling the Corn Stem Borers and Their Roles in Increasing Corn Yield. *J. of Biolo. Scie. Syria.* 26(3).

- Yun, J.S., Kim, H.H., Kim, D.W., Lee, S.M., Kim, D.S., Lee, D.W. (2004).** Pathogenicities of entomopathogenic fungi, Beauveria bassiana and Metarhizium anisopliae against lepidopterous insect pests, *Agrotis segetum*, *Artogeia rapae*, *Mamestra brassicae*, *Plutella xylostella*, *Spodoptera exigua* and *Spodoptera litura*. *Asian J. Turfgr. Sci.*, 18, 221–229.
- Zue-Yun, Y. and Bo-Guang, Z. (2007).** Antifungal activities of matrine and oxymatrin and their synergetic effects with chlorthalonil. *J. fores. Res.* 17:323-325.