

# ***DEDICATION***

***This Research is dedicated to my  
respective mother who have been my  
constant source of inspiration. Without  
her love and support this project  
would not have been made  
possible...!!!***

# ***Acknowledgement***

My deep thanks to my supervisor Dr. Hussain Ahmed Hassan for his constant supervision, inexhaustible patience & unlimited help.

My thanks extend to Dr. Caroline and my colleagues whose help me to complete this research.

## **Abstract**

This study was executed to demonstrate dimension of renal pelvis (length and width in three levels ); the study was conducted at CT departments in Nilain private center and ALripat hospital from march 2013 to may 2013.

Random samples of fifty patients whom referred to CT department with CTU request, 25 male and 25 females with different ages were collected; axial and sagittal cuts in addition to reconstructed coronal section were done to calculate the renal pelvis clearly.

From the study the RT. Renal pelvis was greater than the LT. Renal pelvis dimensions because the liver press the right kidney.

Determine the renal pelvis dimension help in accuracy of diagnoses because there was direct impact between width of renal pelvis and hydronephrosis.

Multidetector computed tomography is the modality of choice to determine dimension of renal pelvis and to visualize some pathological condition.

## ملخص البحث

أجريت هذه الدراسة لمعرفة الأبعاد التشريحية لحوض الكلية (الطول والعرض في ثلاثه مستويات) وقد أجريت الدراسة في كل من مركز النيلين- ومستشفى الرباط الجامعي في

الفترة من مارس 2013 الي مايو 2013

أخذت عينة عشوائية من خمسين مريض تم تحويلهم إلى قسم الأشعة المقطعية بطلب فحص أشعة مقطعية ملونه للجهاز البولي (25 من الرجال و 25 من النساء) من مختلف

الأعمار، وقد خضع جميع المرضى- لإجراء الوضعين (المحوري والسهمي) بالإضافة للوضع التاجي المركب وذلك لقياس حجم حوض الكلية بوضوح.

أوضحت الدراسة أن حوض الكليه اليمين أكبر من حوض الكلية اليسار وأن تحديد أبعاد حوض الكليه يساعد في سهولة ودقه التشخيص لأن لعرض حوض الكلية علاقة بالتموه الكلوي.

أوضحت الدراسة أيضا أن الأشعة المقطعيه متعددة المجسات هي الخيار الأمثل لتحديد أبعاد حوض الكلية وتحديد الأمراض.

## List of contents

<u>Topics</u>	<u>page</u>
Dedications	I
Acknowledgment	II
Abstract (English)	III
Abstract (Arabic)	IV

List of contents	V
List of figures	VII
List of tables	IX
<b>CHAPTER ONE</b>	
1-1 Introduction	1
1-2 Problem of study	3
1-3 Objectives	3
1-4 The study area	4
1-5 Over view of the study	4
<b>CHAPTER TWO:</b>	
2-1 Anatomy of the kidneys	5
2-1-1 Blood supply	8
2-1-2 Nerve supply	9
2-2 Pathology	10
2-2-1 Renal stone	10
2-2-2 Renal mass	11
2-2-3 Hydronephrosis	12
2-2-4 Compensatory renal hypertrophy	13
2-2-5 Chronic Pyelonephritis	13
2-3 Physiology	16
2-3-1 Maintenance of homeostasis	16
2-3-2 filtration	18
2-3-3 Storage and excretion of waste	19

2-3-4 Production of hormones	19
2-4 Multidetector computed tomography	21
2- 5 Detector layout and slice definition	23
2- 6 Z-interpolation, pitch and mAs per slice conception	25
2- 7 Multidetector CT of the kidney	28
2- 8 Technique and Scanning Protocol	30
2- 8-1 protocol	31
2-9 previous study	32
<b>CHAPTER THREE</b>	
3- Material and Methods	33
3-1 Patients	33
3-2 Machine	33
3-3 CTU protocol	33
3-4 image interpretation	35
<b>CHAPTER FOUR</b>	
4- Results	36
<b>CHAPTER FIVE</b>	
5- 1 Discussion	41
5-2 Conclusion	42
5-3 Recommendations	42
<b>REFERENCES</b>	43
<b>APPENDIX</b>	44

## List of figures

<b>Figure #</b>	<b>Details</b>	<b>pages #</b>
(2-1)	Anatomy of the renal pelvic	<b>6</b>
(2-2)	coronal section show measurement of pelvic	<b>35</b>
(4-1)	The gender frequency	<b>36</b>
(4-2)	The age classes and frequency	<b>36</b>



## List of tables

<b>Table #</b>	<b>Details</b>	<b>pages #</b>
--------------------	----------------	--------------------

(4-1)	the Gender, Frequency and Percentage	<b>36</b>
(4-2)	the age classes and Frequency	<b>37</b>
(4-3)	the mean and Standard Deviation (STDV) of the RT kidney Measurements For the Total Sample	<b>37</b>
(4-4)	the mean and Standard Deviation (STDV) of the LT kidney Measurements For the Total Sample	<b>38</b>
(4-5)	The mean and Standard Deviation (STDV) of the RT kidney Measurements for the Normal and Sample with Hydronephrosis at <i>P-Value=0.05</i>	<b>38</b>
(4-6)	The mean and Standard Deviation (STDV) of the LT kidney Measurements for the Normal Sample and Sample with Hydronephrosis at <i>P-Value=0.05</i>	<b>39</b>
(4-7)	Cross Tabulation between RT Kidney measurement and diagnosis as RT, LT Bilateral hydronephrosis and other causes as stone	<b>39</b>
(4-8)	Cross Tabulation between LT Kidney measurement and diagnosis as RT, LT Bilateral hydronephrosis and other causes as stones	<b>40</b>

## ABBREVIATION

ADH	Antidiuretic Hormone
CT	Computed Tomography
CTU	Computed tomography urology
IVP	Intravenous Paleography
IVU	Intravenous Urography
MDCT	Multidetector Computed Tomography
MRI	Magnetic Resonance Imaging
PCS	Pelvicalyceal system
PTH	Prathyroid hormone
RAS	Renin Angiotensin System
SST	Single Slice CT
VRT	Volum Rendring Technique