

References

- Alao**, S.J. and Balanve, D. (1984). Growth and carcass composition of broiler fed sunflower oil and olive oil. Br. Poult. Sci., 25:209-219.
- Allam**, S. (2000). Poultry Breeding and Husbandry. The Egyptian Anglo Library, 165, M Fareed street, Cairo, Egypt.
- Allison**, J.B. (1955). Biological evaluation of protein. Physiol Riv., 35: pp 664.
- AOAC** (1988). Official Methods of Analysis (12th ed), Association of official Analytical chemists, Washington D.C, USA.
- Artman**, N.R. (1964). Interactions of fats and fatty acids as energy sources for the chick. Poult. Sci., 43:994-1004.
- Baracos**, V.E. and Olomu, J.M. (1989). Effects of flaxseed oil on prostaglandin production and performance of broiler chickens. Poultry Sci., 68 (1) : 9 (Abstr.).
- Bartov**, I., Bornstein S. and Lipstein, B. (1974). Effect of calorie to protein ratio on the degree of fatness in broilers fed on practical diets. Br. Poultry Sci., 15:107-117.
- Bartov**, I. (1987). Combined effect of age and ambient temperature on the comparative growth of broiler chicks fed tallow and soybean oil. Poultry Sci., 66:273-279.
- Bensadoum**, A. and Rathfield (1972). Form of adsorption of lipid in chickens. Proc. Sci., Exp. Miol. Med. 141:814-817.
- Brake**, J. (1989). The effect of dietary fat on broiler breeder performance. Poultry Sci., 68 (1): 173 (Abstr.) .
- Butolo**, J.E. (2002.). Qualidade de ingredientes na alimentação animal. Campinas colégio Brasileiro de Nutrição Animal, 430p, Av. Andrade Neves, Castelo, Brazil.

- Carew**, L.B.Jr. and Hill F.W. (1964). Effect of corn oil on metabolic efficiency of energy utilization by chicks. *J.Nutr.*, 83:293-299.
- Carew**, L. and Hill, F.W. (1967). Effect of diethylstilbestrol on energy and protein utilization by chicks fed a diet high in fat content. *J.Nut.*, 92:393-398.
- **Carew**, L.B.Jr.; Hopkins, D.T.; Nesheim, M.C. (1964). Influence of amount and type fat on metabolic efficiency of energy utilization by the chicks. *J.Nutr.*, 83:300-306.
- Carew**, L.B.; Nesheim M.C. and Hill, F.W. (1963). The relationship of dietary energy level and density to growth response of chicks to fats. *Poultry Sci.*, 42:710-718.
- Carroll**, K.K. (1965). Dietary fat and fatty acid composition of tissue lipids. *J. Am or. Oil. Chem. Soc.* 42:516-528.
- Carver**, D.S.; Rice E.E.; Gray, R.E. and Mone, P.E. (1955). The utilization of fats of different melting points added to broiler feed. *Poultry Sci.*, 34:544-546.
- Cary**, M.C. (1983). Lipid digestion and adsorption. *Ann. Rev. Physoil*, 45:651-672.
- Chakkodaby**, L., R.; Ranjite, P. and Ganguly, J. (1980). The effect of dietary unsaturated oil on brothsynthesis of cholesterol and biliary and faecal excretion of cholesterol and bile acid in rat. *J. Nutr.*, 110:2140-2158.
- Charles**, D.R.; Goom, C.M. and Bray, T.S. (1981). The effects of temperature on broilers: interactions between temperature and feeding regime. *Br. Poultry Sci.*, 22: 475-481.
- Callcomb**, D.M.; Richardson J.M. and Kir, S. Ed. (1974). Anaerobic bacteria and deconjugated bile salt in upper small intestine of infant with G1T Disorder. *Poult. Sci.*, 63:97-108.

- Christmas**, R.B. and Harms, R.H. (1988). Effect of energy on the performance of broiler chicks fed various levels of monensin. *Poultry Sci.*, 67:452-454.
- Chung**, H.C.; Guenter, W.; Rotter, R.C.; Crow, G.H. and Stanger, N.E. (1993). Effects of dietary fat source on Sudden death syndrome and cardiac sarcoplasmic reticular calcium transport in broiler chickens. *Poult. Sci.*, 72:310-316.
- Cocks**, L.V. and Rede, C. V. (1966). Laboratory Handbook for oil and fat analysts. Baily's industrial oils and fat product ,England.
- Coon**, C.N.; Becker, W.A. and Spencer, J.V. (1981). The effect of high energy diets containing supplemental fat on broiler weight gain, feed efficiency, and carcass composition. *Poultry Sci.*, 60:1250- 1253.
- Corew**, L.B.; Neshien, M.C. and Hill, F.W. (1963). The relationship of dietary energy level and density to growth response of chick to fat. *Poult. Sci.*, 42:710-718.
- Crespo**, N. and Nsteve-Garcia, E. (2002). Dietary polyunsaturated fatty acid decrease fat deposition in separable fat depot but not in the remainder carcass. *Poultry Science*; 81:512-518b.
- Dale**, N.M. and Fuller, H.L. (1979 b). Effects of diet composition on feed intake and growth of chicks under heat stress. I-Dietary fat levels. *Poultry Sci.*, 58:1529-1534.
- Dale**, N.M. and Fuller, H.L. (1979 a). Effects of low temperature, diet density and pelleting on the preference of broilers for high fat ration. *Poultry Sci.*, 58: 1337-1339.
- Deaton**, J.W.; McNaughton, J.L.; Rece, F.N. and Lott, B.D. (1981). Abdominal fat of broilers as influenced by dietary level of animal Fat. *Poultry Sci.*, 60:1250-1253.

- DeBasilio**, V.; J.J. Montilla and Vargas, R.E. (1989). Effect of palm oil , palm kernel oil , corn oil and lard on performance and fatty acid pattern of breast and thigh muscles of broilers. *Poultry Sci.*, 68(1): 39 (Abstr.).
- DeGroote**, G. (1968 a). Energetic evaluation of unstablized and net energy for maintenance and growth. *Feed stuffs*, 40(51):26-28.
- DeGroote**, G.; Reyntens, R. N. and Amich-Gali, J. (1971). Fat studies.2. The metabolic efficiency of energy utilization of glucose, soybean oil and different animal fats by growing chicks. *Poultry Sci.*, 50:808-819.
- Donaldson**, W.E.; Combs, G.F., Romser, G.L. and Spittle, W.C. (1957). Studies on energy levels in poultry rations. 2. Tolerance of growing chicks to dietary fat. *Poultry Sci.*, 36:807.
- Douglas**, J.H.; Sullivan, T.W. and Bond, P.L. (1988). Dietary protein and fat level for broiler 42 to 76 days of age. *Poultry Sci.*, 67 (1): 80 (Abstr.) .
- Edwards**, H.M.Jr. and May, K.N. (1965). Studies with menhaden oil in practical-type broiler rations. *Poultry Sci.*, 44:685-689.
- Farrel**, D.J. (1978). Efficiency of utilization by growing chicken of the energy of dietary fat and oil. *Br. Poultry Sci.*, 19:105-109.
- Fedde**, M.R.; Waible initials P.E. and Burger, R.E. (1960). Factors affecting the absorption of certain fat in chick. *J. Nut.*, 70:447-452.
- Ferreira**, W.M. (1999) Diegestão e metabolismo dos lipídios. Belo Horizonte: Escola de Veterinária da UFMG; P. 1-34.
- Freeman**, C.D. (1969). Properties of fatty acid in dispersion of emulsified lipid and bile salt and significance of these in fat absorption in pig and sheep. *Br. J. Nut.*, 23: 249-264.
- Fuller**, H.L. and Groots, M.E. (1983). Influence of the intestinal microflora on nutrition. In: *Physiology and Biochemistry of Domestic Fowl*.

- Fuller**, H.L. and Redon, M. (1977). Energetic efficiency of different dietary fats for growth of young chicks. *Poultry Sci.*, 59: 549- 557.
- Furman**, R.H. (1968). Medium Chain Triglycerides. Univ. Pennsylvania press, Philadelphia, U.S.A.
- Griffin**, H.D.; Whitehead, C.G. and Broodbent, L.A. (1982). The relationship between plasma triglycerides concentration and body fat content for male and female broiler .Br. *Poultry. Sci.*, 23 (1) :15.
- Gurr**, M.I. and James, A .T. (1980).Lipid Biochemistry. Chapman, London and New York.
- Hallander**, D. and Morgan, D. (1979).Increase in cholesterol intestinal adsorption with age in rat .Exp. gerntal., 14:201-204.
- Hardy**, L.B.; Auger, V. and Wilcax, F.H. (1962).Genetic differences in serum cholesterol in chickens .Amer.J. physoil., 202:997-1001.
- Hildon**, J.A; Jones, E.C. and Khead, A.T. (1934). The direct effect of dietary fat limitation on fatty acid composition of carcass fat of bird .Biochem. J., 28:786-795.
- Hopkins**, D.T. and Nesheim, M.C. (1961).Essential fatty acids for poultry. Proc. Cornell Nutrition Conference, Ithaca, New York.
- Hurwitz** , S .;Bar , A. ; Kalz , M. ; Sklan , D. and Budowskip initials E. (1973). Using C¹⁴ labelled fatty acids and non absorbed marker reported evidence for identifying the jejunum as the major site of fat absorption in chick and laying hens .J.Nut.,103:543-547.
- Huyghebaert**, G; De munter, initials C. DeGroote, G. (1988). The metabolizable energy (AMEN) of fats for broilers in relation to their chemical composition. *Animal feed Science Technology*, vol. 20:45.
- Ikegwounu**, F.I. and Aim, T.A. (1977). Age influence of cholesterol and ATP activity infest of prepuberal chick. *Poult. Sci.*, 56:1158-1180.

- Jackson**, N.; Kilpatrick, H.R. and Fulton, R.B. (1969). An experimental study of the utilization by the laying hen, of dietary energy partially supplied as animal fat. *Poultry Sci.*, 10:155-126.
- Jones**, J.E. and Barnett, B.D. (1973). The effect of dietary fat on turkey hen resistance to high environmental temperature. *Poultry Sci.*, 52:1506-1509.
- Kahnke**, I.M.J. (1966). Atomic Absorption spectrophotometer Applied to Determination of Zinc in formalinized Human Tissues. *News*, 1:5-7.
- Kalmbach**, M.P. and Patter, L.M. (1959). Study in evaluating energy control of feed for chick and comparative value corn oil and tallow. *Poult. Sci.*, (38):1271.
- Ketels**, E. and Groote, G.D. (1987). Effect of fat source and level of fat inclusion on utilization of fatty acids in broiler diet. *Archive Fur Geflege L Kunoh*, 51(4) : 127-132.
- Klarich**, E. initials and **Fumer**, B.E. (1989). Influence of fishing diet on phospholipid composition on fatty acid profile of individual phospholipid in lean muscle of beef cattle. *J. Anim. Sci.*, 67:2282.
- Kleiber**, M. (1975). *The Fire of Life : an introduction to animal energetics*. Krieger Publishing Company, Huntington, N.Y.
- Lara**, U.C. (2004) Efeito da fonte lipídica em dietas para frangos de corte sobre o desempenho, rendimento e composição da carcaça [dissertação]. Belo Horizonte: Escola de veterinária, UFMG.
- Lessen** S., Summers, J.D. (2001). *Nutrition of the Chicken*. 4th ed. Ontario University Books; 413 PP.
- Leveiile**, G.A. and Serimshow D.R., (1961). Influence of dietary protein on serum protein component and cholesterol in growing chick. *J. Nut.*, 74:500.

- Lipstein**, B. and Bornstein E., (1975). Extra-caloric properties of acidulated soybean-oil soap stock for broilers during hot weather. *Poultry Sci.*, 54: 396-404.
- Machael**, E.; Lisano, C. and Janes, F.; Kelnnamer E.L. (1977). Value for several blood parameters in esters wild turkey. *Poult. Sci.*, 56: 157-166.
- Malmros**, G.G. and Wigand, J.L. (1957). Influence of carbohydrate supplemental fat source of metabolizable diet. *Poult. Sci.*, 59:2129-2135.
- Makki, Raga**. H. (1997). The Effect of Dietary Fat on Broiler Chicken Performance, Serum Cholesterol, Lipid and Protein. M.Sc. Thesis, University of Khartoum, Sudan.
- Marion**, J.E. and Woodroof, J.G. (1963). The fatty acid composition of breast, thigh and skin tissue of chicken broiler as influenced by dietary fats. *Poultry Science*, 42:1202-1207.
- McDonald**, P.; Edwards, R.A. and Greenhalgh, J.F.D. (1988). *Animal Nutrition*. 4th ed., John Wiley and Sons, Inc., New York.
- Menten**, J.F.M.; Gaiotto J.F. and Racanicci , A.M.C. (2003) Valor nutricional quality de óleos e gordura para frangos de corte. In: Simpósio sobre Manejo e Nutrição de Aves e suinos; campinas: Colégio Brasileiro de Nutrição Animal, P.93-135.
- Mittelstaedt**, C.W.; Murray, E.E. and Teeler, R.G. (1989). Ration caloric density effects on broiler growth and efficiency. *Poultry Sci.*, 68 (1): 194 (Abstr.).
- **Mohamed, Asma**. H.. (1996). Broiler Chickens Performance as Affected by Animal Fat and Plant Oil Under Hot Arid Condition of Sudan. M. Sc., thesis, University of Khartoum, Sudan.
- Moiorino**, P.M.; Alhozab, A.A., Rimiltchell G.G. initials and Reid , B.L. (1986).Animal fat effect on nutrient utilization. *Poult. Sci.*, 65:2304-2313.

- Montgomery**, R.; Robert, L.; Dryer, T. ; Conway, W. ; Arthur, A. and Pector, C. (1974). Biochemistry CA case oriented approach. The C.V. Mosby Company Saint Louis, U.S.A.
- Moreng**, R.E. and Avens, J.S. (1985). Poultry Science and Production. Restone Publishing Company inc., U.S.A.
- Nakhata**, N.and Anderson, J.O. (1982). Describing the relation between dietary protein and energy levels and chick performance by mathematical equations. *Poultry Sci.*, 61:891-897.
- Nascif**, C.C.C. ; Gomes, P.C. ; Albino, L.F.T. and Rostagno, H.S.(2004) . Detrminação dos valores energeticos de alguns Óleos e gorduras para pintos de corte machose fêmeas aos 21 dias de idade. *Revista Brasileira de Zootecnia*, 33(2): 375-385.
- NRC** (1984). Nutrient Requirements of Poultry. National Research Council, Washington, U.S.A.
- Newman** , R.E. ; Bryden , W.L. ; Fleck , E.; Ashes , J.R. ; Buttemer , W.A. ; Storlien, L.H. and Downing, J.A. (2002). Dietary n-3 and n-6 fatty acids alter avian metabolism: molecular-species composition of breast-muscle phospholipids. *British Journal of Nutrition*, 88:11-18.
- Obunike**, J.C.; Geachin, M.C; Gardner, F.A. and Bailey, C.A. (1988).Effect of corn and sorghum based diet, with and without supplement fat, on processing yields in broiler chickens. *Poultry Sci.*, 67:130 (Abstr.).
- Olomu**, J.M. and Baracos, V.E. (1989). The influence of dietary fat level and source on the performance of broiler chickens. *Poultry Sci.*, 68(1): 108 (Abstr.) .
- Parker**, M.M.; Humoller, F.L.L. and Mahler, D.L. (1967). Determination of copper and Zinc in biological materials. *Clin. Chem.*, 13: 40.

- Patterson**, B.W.; Wong, W.W.; Sheng, H.P.; Mersman, H.J.; Insull, W.; Klein, D.D.; Fioratt, M.L. and Pond, W.G. (1992). Neonatal genetically lean and abuse pig respond differently to dietary cholesterol. *J. Nut.*, 122:1820.
- Paul**, R. and Ganguly, J. (1976). Effect of unsaturated lipid on bile flow and biliary excretion of cholesterol and bile salt in rats. *Chemphysio. Lip.*, 17: 315-323.
- Powell**, D.W. and Barrier, H. (1981). Function of epithelia. *Am. J. Physiol.*, 241: 275-288.
- Prolin**, D. and Hussein, T.H. (1982). The effect of bile acid on lipid and nitrogen retention to carcass composition and dietary energy. *Poult. Sci.*, 61: 1697.
- Quart**, M.D.; Damron, B.L.; Martin, F.G.; Christmas, R.B. and Sloan, D.R. (1989). Effects of feed fats on broiler performance. *Poultry Sci.*, 68 (1): 109 (Abstr.) .
- Rand**, N.T.; Scott, H.M. and Kummerow, F.A. (1958). Dietary fat in the nutrition of the growing chick. *Poult. Sci.*, 37: 1075-1085.
- Reid**, B.L. and Weber, C.W. (1975). Supplemental dietary fat and laying hen performance. *Poult. Sci.*, 54: 422-428.
- Reiser**, R. (1951). The synthesis and interconversion of poly unsaturated fatty acids by laying hens. *J. Nut.*, 44:159.
- Renner**, R. (1964). Factors affecting the utilization of Carbohydrate- free diets by the chick. I. level of protein. *J. Nutrition*, 84: 322-325.
- Renner**, R. and Hill, F.W. (1973). Metabolizable energy values of fats and fatty acids for chickens. *Proc. Cornell Nutr. Conf.*, 95-100. Cited in Br. *Poultry Sci.*, 14:279-289.

- Renner, R.F. and Hill, F. (1961). Utilization of fatty acid by chickens of various ages. *J. Nut.*, 74: 259-264.
- Robert, R.B.; Morrissey, W.E. and Donaldson, J. (1977). Diet composition and cholesterol in Japanese Quail. *Poult. Sci.*, 56:2018-2110.
- Salmon, R.E. (1974). Effect of dietary fat concentration and energy to protein ration on the performance, yield of carcass component and composition of skin and meat of turkeys as related to age. *Br. Poult. Sci.*, 15:543-560.
- Sanz, M.; Flores, A; Perez, D.E.; Ayala, P. and Lopez-Bote, C. J. (1999). Higher lipid accumulation in broilers fed on saturated fats than in those fed unsaturated fats. *British Poultry Science*; 40: 95-101.
- Sanz, M.; Lopez-Bote, C. J.; Menoyo, D. and Bautista, J.M.(2000 b).Abdominal fat deposition and fatty acid synthesis are lower and α -oxidation is higher in broiler chickens fed diets containing unsaturated rather than saturated fat. *Journal of Nutrition*; 130: 3034-3073.
- Scheibel, M.S.; Coon, C.W. and Kelly, W. (1979). The heat increment of feed used in poultry diet. *Nut. Rev. Int.*, 20:87.
- Schettler, G.; Nüssel, E.; Arbeitsmed. Sozialmed. Praventivmed. 10(1975) 25. Determination of cholesterol in serum or plasma on photometric system.
- Scott, M.L.; Nesheim, M.C. and Young, R.J. (1982). *Nutrition of Chicken*. 3rd ed. M.L. Scott and Associates, Ithaca, New York.
- Sell, J.L. and Owings, W.J. (1981). Supplemental fat and metabolizable energy-to-nutrient rations for growing turkeys. *Poultry Sci.*, 60: 2293-2305.
- Sell, J.L. and Thampson, J (1965). The effect of ration pelleting and level of fat on the efficiency of nutrient utilization. *Br. Poult. Sci.*, 6: 345-354.
- Sensor, J.R. (1964). Intestinal absorption of fat. *J. lipid Res.*, 5:495-521.
- Serfin, J.A. and Nesheim, M.C. (1970). Influence of dietary at labile factors in soybean meal upon bile acid pool turnover in chick. *J.Nut.*, 100: 786-795.

- Shermer**, W.D. (1990). Effects of oxidation on the quality of ingredients and feed of Poult. In: Maryland Nutrition conference, Maryland, EUA.
- Sibbald**, I.R. and Kromer, K.G. (1980). Effect of the basal diet on utilization of fat as source of true ME lipid fatty acids. *Poult. Sci.*, 59: 316-324.
- Sibbald**, I.R. and Price K. (1977). Effect of level of dietary inclusion of Ca on true metabolizable energy value of fat. *Poult. Sci.*, 56: 2070- 2078.
- Sim**, J.S. (1977). Effect of dietary factors on serum cholesterol and egg yolk cholesterol level of laying hens. *Poult. Sci.*, 56: 1616-1621.
- Sim**, J.S.; Kills, W.D. and Bragg, D.B. (1980). Effect of dietary egg yolk on serum cholesterol level of White Leghorn cockerel. *Poult. Sci.*, 59 (8) : 1812-1817.
- Skinner**, J.; Park, T. and Waldroup, W. (1989). Effects of constant or changing dietary energy level on performance of male broilers. *Poult. Sci.*, 68: (1) :134. (Abstr.) .
- Sklon**, D.; Budowki, P. and Harwitz, J. (1977). Site of bile absorption in rat. *J. Lipid*, 21: 467.
- Spittle**, D.; Kavongroven, L.K. and Rovisubbiah, M.T. (1976). Concentrate change of bile acid in sequential segment of pigeon intestine and their bile acid absorption. *Biochem.-Biophy. Acta*, 441:32.
- Sulieman, Y.R. and Mabrouk, Afaf. A. (1999). The Nutrient Composition of Sudanese Animal Feeds. Bulletin iii Central Animal Nutrition Research Laboratory, Animal Production Research Center, Kuku, Khartoum North, Sudan.
- Sunde**, M.L. (1956). The effect of fats and fatty acids in chick ration. *Poult. Sci.*, 35: 362-368.

- Stanley**, V. G.; Krisdakasivivat, G. and Alden, R. (1988). Effect of growth-stimulants (supplementary 10 dine and bacitracin MD) on abdominal fat deposits in broilers fed four levels of dietary fat. *Poultry Sci.*, 67 (1): 161. (Abstr.) .
- Thacker**, P.A., and Cocmpbell, G.L.; (1994). Composition and nutritive value of acidulated fatty acids, degummed conola oil and tallow as energy sources for starting broiler chicks. *Animal feed and technology*, 46: 251-260.
- Thamas**, O.P. and Combs, G.E. (1967). Relationship between serum protein level and body composition in chick. *J. Nut.*, 91:468.
- Tufft**, L.S. and Jensen, S. (1990). Effect of dietary taurine on performance and fat retention of broilers and turkey poult fed varying levels of fat. *Poult. Sci.*, 69(1): 195 (Abstr.) .
- Valencia**, M.E.; Maiorino, P.M. and Reid, B.L. (1980 a). Energy utilization in laying hens. 11 Energetic efficiency and added tallow at 18.3 and 35°C. *Poultry Sci.*, 59: 2071-2076.
- Vargas**, R.E.; deBailio, V. and Montilla, J.J. (1989). Effects of palm oil, palm kernel oil and lard on total plasma cholesterol and lipoprotein in broilers. *Poult. Sci.*, 68(1): 151(Abstr.).
- Vermeersh**, G. and Vanschoubrock, F. (1968). The quantification of the effect of increasing levels of various fats on body weight gain, efficiency of food conversion and food intake of growing chicks. *British Poultry Sci.*, 9: 13-30.
- Volker**, L.and Amich-Gali, J. (1968). Recherches surles valeurs comparées de L énergie metabolisable et autres graisses das les ratios destinées aux broilers, National Renders Association, Inc., Rome, Publication No. 16.

- Weber**, F. (1981). Absorption mechanisms for fat soluble vitamin and effect of other food constituent nutrition in health and diseases. International symposium, XII Int. Congress of Nutr., P. 119-135.
- Weiss**, H.S. and Fisher, H. (1957). Plasma lipid and organ changes associated with feeding of animal fat in laying chicken. J. Nut. 61:267.
- Wiseman**, J. (1984). Fats in Animal Nutrition. Butterworths, London .
- Yau**, J.C.; Sams, A.R.; Denton, J.H. and Baily, C.A. (1989). Enrichment of selected fatty acids in broiler tissues. Poultry Sci., 68(1):162 (Abstr.) .
- Young**, A. , and **Gorett** B., (1963). Effect of oleic and linoleic acid on the absorbance of saturated fatty acid in chicks. J. Nut., 81: 321- 329.
- Zafrira**, N.; Advorin E. ; Miluot C. and Mokadi, S. (1990). Extra caloric effect of soybean oil at different dietary energy levels. Poultry Sci., 69 (1): 99 (Abstr.).
- Zumbado**, E. (1990). Nutritive vale of Palm oil and Palm kernel oil for broilers. Poultry sci., 69 (1) : 151 (Abstr.).