

## References:

***Achiron R and Achiron A:*** Transvaginal fetal neurosonography: The first trimester of pregnancy. In: Chervenak FA, Kurjak A and Comstock CH (eds.), Progress in Obstetric and Gynecology Sonographic Series, Ultrasound and Fetal Brain, Parthinon Pub, 1999; pp: 90-108.

***Anderson N, Allan R, Darlow B, et al:*** Diagnosis of intraventricular hemorrhage in the newborn. AJR Am J Roentgenol ,1994.

***Anderson N, Allan R, Darlow B, et al:*** Diagnosis of intraventricular hemorrhage in the newborn. AJR Am J Roentgenol, 2004.

***Athni SS, Cherches IM and Loftus B:*** Clinical neuroanatomy. In: L.A. Rolak (ed.): Neurology Secrets (2nd edition) . 1998; pp: 13-49.

***Babcock CJ, Cong BW, Salamat MS, et al:*** Sonographic anatomy of the developing cerebellum: normal embryology can resemble pathology. 1996; AJN 166 (2): 427-33.

***Barkovich AJ:*** MR and CT evaluation of profound neonatal and infantile asphyxia: AJNR Am J Neuroradiol, 1999; 13:959.

***Benjamin Y:*** Hypoxic-Ischemic Brain Injury: Imaging Findings from Birth to Adulthood1. RadioGraphics, 2008 ; RadioGraphics, 28, 417-439. 28, 417-439.

***Boichot Ch, Paul M, Jean B:***Term Neonate Prognoses after Perinatal Asphyxia: Contributions of MR Imaging, MR Spectroscopy,

Relaxation Times, and Apparent Diffusion Coefficients. *Radiology* 2006; 239(3): 839-848).

**Boog G:** Cerebral palsy and perinatal asphyxia (I--diagnosis). *Gynecol Obstet Fertil.* 2010; 38(4):261-77.

**Cai Q, Xue XD, Fu JH:** Progresses in studies on magnetic resonance imaging of neonatal hypoxic-ischemic encephalopathy. 2010; 48(3):227-31.

**Carlson N. Allyn A and Bacon K:** *Physiology of Behavior*, 5th ed. 1997; pp: 154-175.

**Chao CP, Zaleski CG, Patton AC:** Neonatal hypoxic-ischemic encephalopathy: multimodality imaging findings. *Radiographics.* 2006; 26 Suppl 1:S159-72.

**Chau V, Poskitt KJ, Sargent MA, et al:** Comparison of computer tomography and magnetic resonance imaging scans on the third day of life in term newborns with neonatal encephalopathy. *Pediatrics.* 2009; 123(1):319-26.

**Chris D:** Newborn Breathing Problems :Birth Asphyxia. 2009; PP: 740-783.

**Christine P, Christopher G. Zaleski, and Alice C et al:** Neonatal Hypoxic- Ischemic Encephalopathy: Multimodality Imaging Findings. *Radiographics* 2006;26:S159-S172.

**Chugani HT:** Functional brain imaging in pediatrics. *Pediatr Clin North Am*, 1992; pp:39-77.

**Connolly D, Widjaja E and Griffiths P:** Involvement of the Anterior Lobe of the Cerebellar Vermis in Perinatal Profound Hypoxia :*American Journal of Neuroradiology* 28, 2007; pp:16-19.

**Copp AJ:** Neural tube defects: *Trends Neuroscience*, 1997; 16: pp: 381-3.

**Cowan F, Rutherford M, Groenendaal F, et al:** Origin and timing of brain lesions in term infants with neonatal encephalopathy. *Lancet* 2003;361(9359): 736–742.

**De Vries LS and Jongmans MJ:** Long-term outcome after neonatal hypoxic-ischaemic encephalopathy. *Arch Dis Child Fetal Neonatal Ed.* 2010; 95(3):F220-4.

**di Pietro MA, Brody BA and Teele RL:** Peritrigonal echogenic ‘blush’ on cranial sonography: pathologic correlates. *AJR Am J Roentgenol*, 1996;146:1067-1072.

**Donald N, Di Salvo, MD:** A New View of the Neonatal Brain: Clinical Utility of Supplemental Neurologic US Imaging Windows. *Radiographics*. 2001;21:943-955.

**Edwards AD and Nelson KB:** Neonatal encephalopathies. Time to reconsider the cause of encephalopathies. *BMJ* ,1998;317(7172):15378.

**Encarta R** : Neurophysiology: Nervous System : The Physiology of Brain. 1994; pp: 381-392.

**Fahy J and King T**: Intrauterine Asphyxia. Clinical Implication For Providers of Intrapartum Care. Midwife Women's Health. 2005;

**Gallucci M, Capoccia S and Catalucci A**: Radiographic Atlas of Skull and Brain Anatomy.2003.

**Gassner I, Unsinn KM, Geley T, et al**: Sonography of the central nervous system in prematures and newborns. Radiology 40(1), 2000; pp:35-42.

**Gordon S and Johnson Jr**: A Guide to Brain Anatomy, Function and Symptoms. 2009; pp: 435-445.

**Graham EM, Ruis KA, Hartman AL, et al**: A systematic review of the role of intrapartum hypoxia-ischemia in the causation of neonatal encephalopathy: Am J Obstet Gynecol 2008; 199:587.

**Jones DK, Simmons A, Williams SC, et al**: Non-invasive assessment of axonal fiber connectivity in the human brain via diffusion tensor MRI. Magn Reson Med 2009; 42: 37–41.

**Jones RA, Palasis S and Grattan-Smith JD**: MRI of the neonatal brain: optimization of spin-echo parameters. AJR Am J Roentgenol 182(2), 2004; pp:367–372.

**Jouveta P, Cowana M, Coxa P, et al** : Reproducibility and Accuracy of MR Imaging of the Brain after Severe Birth Asphyxia: American Journal of Neuroradiology 20, 1999; 1343-1348.

**Ito Bashir A., Zakaria M. Al-Hawsaw, and Anwar H. Khan:**  
Hypoxicischemic encephalopathy Incidence and risk factors in North Western Saudi Arabia, Saudi Med J 2003; Vol. 24 (2): 147-153

**Kanik E, Ozer EA, Bakiler AR, et al:** Assessment of myocardial dysfunction in neonates with hypoxic- ischemic encephalopathy: is it a significant predictor of mortality? J Matern Fetal Neonatal Med. 2009; 22(3):239-42.

**King, T and Parer, J:** The physiology of fetal heart rate patterns and perinatal asphyxia: J perinatal neonatal nurse, 2000; pp:14-19.

**Kirby CL and Rosenberg HK:** The infant brain: Ultrasonography. In: Goldberg BB, Pettersson H, eds. 2006.

**Kliegman** : Nelson Text Book of Pediatrics, 18th ed. 2007; pp: 555-561.

**Lee BCP, Liper E, Naas R, et al:** MRI of the central nervous system in neonates and young children. AJNR 7:605- 16. 1986.

**Li AM, Chau V, Poskitt KJ, et al** : White matter injury in term newborns with neonatal encephalopathy. *Pediatr Res.* 2009; 65(1):859.

**Liauw L, van der Grond J, van den A, et al** : Is there a way to predict outcome in (near) term neonates with hypoxic-ischemic encephalopathy based on MR imaging? *AJNR Am J Neuroradiol:* 2008; 29(9):1789-94.

**Lilian T, Siea L, Marjo S, et al** : Early MR Features of Hypoxic-ischemic Brain Injury in Neonates with Periventricular Densities on Sonograms: *American Journal of Neuroradiology* 21, 2000; pp:852-861.

**Lishya L, Jeroen G, Annette A. et al** : Hypoxic-Ischemic Encephalopathy: Diagnostic Value of Conventional MR Imaging Pulse Sequences in Term-born Neonates. *Radiology* 2008; 247(1): 204-212).

**Logitharajah P, Rutherford MA and Cowan FM:** Hypoxic-ischemic encephalopathy in preterm infants: antecedent factors, brain imaging, and outcome. *Pediatr Res.* 2009; 66(2):222-9.

**Luna JA and Goldstein RB:** Sonographic visualization of neonatal posterior fossa abnormalities through the posterolateral fontanelle: *AJR Am J Roentgenol,* 2000; 174:561-567.

**lyнна y.litteton:** maternal, neonatal and women's health nursing. joan engebretson., fetal assesment. 2002; pp:178-183.

**Manohar M and Shroff MD:** Standard neonatal cranial ultrasound scan views: Neonatal cerebral ultrasound, 2007; 78-103.

**Manohar M. João P and Raybaud c:** MR Imaging for Diagnostic Evaluation of Encephalopathy in the Newborn- RSNA, 2010;

**Martin JH:** Development of the central nervous system. In: Martin JH (ed.): Neuroanatomy, Text and Atlas (2nd edition), 1996; pp: 33-59.

**McMinn RM:** Central nervous system. In: McMinn RMH (ed.), Last's Anatomy Regional and Applied (8th edition), Churchill Livingstone, 1990, pp: 579-99.

**Ment LR, Bada HS, Barnes P, et al:** Practice parameter: neuroimaging of the neonate: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. Neurology,2002; 58:1726.

**Miller SP, Cozzio CC, Goldstein RB, et al:** Comparing the diagnosis of white matter injury in premature newborns with serial MR imaging and transfontanel ultrasonography findings: AJNR Am J Neuroradiol 2003; 24:1661.

**Nagy T:** Neuropathology CNS pathology: special features; Cells and cellular pathology in the CNS. 2003; pp: 55-58.

**O`Rahilly R and Müller F:** The Embryonic Human: The central nervous system, 1994; 123-6.

**Ong LC, Kanaheswari Y, Chandran V, et al:** The usefulness of early ultrasonography, electroencephalography and clinical parameters in predicting adverse outcomes in asphyxiated term infants. Singapore Med J. 2009; 50(7):705-9.

**Paul G, Erik B, and Voet D:** An Atlas of Neonatal Brain Sonography: 2006 pp: 212-220.

**Perlman JM:** brain injury in term infant: seminperimatol, 2004; 28-415.

**Pottumarthi V and Prasad:** Introduction to Magnetic Resonance Imaging and Spectroscopy Magnetic Resonance Imaging. Methods and Biologic Applications.2006;2-55.

**Redline RW:** Severe fetal placental vascular lesions in term infants with neurologic impairment: Am J Obstet Gynecol , 2005; 192:452.

**Robertson RL, Ben-Sira L, Barnes PD, et al:** MR line-scan diffusion-weighted imaging of term neonates with perinatal brain ischemia: AJNR Am J Neuroradiol;20(9): 1999; pp:1658–1670.



**Ross MG and Gala R:** Use of umbilical artery base excess. Algorithm for timing of hypoxic injury. Obstetric and gynecological . 2002;

187-1.

**Ruis KA, Lehmann CU, Northington FJ, et al :** Neonatal brain imaging and the identification of metabolic acidemia and hypoxic-ischemic encephalopathy. J Matern Fetal Neonatal Med. 2009; 22(10):823-8.

**Rutherford M, Edward D, Saeed N, et al :** MR Imaging Quantification of Cerebellar Growth Following Hypoxic-Ischemic Injury to the Neonatal Brain: American Journal of Neuroradiology 25, 2004; pp: 463-468.

**Rutherford M, Ramenghi L, Edwards A, et al:** Assessment of brain tissue injury after moderate hypothermia in neonates with hypoxic–ischaemic encephalopathy: a nested substudy of a randomised controlled trial. Lancet Neurology, 2010; 9 (1): 39-45.

**Sarnat HB and Sarnat MS:** Neonatal encephalopathy following fetal distress. A clinical and electroencephalographic study. Arch Neurol ,1976;33 pp:696-705.

**Shaywitz BA. McMillan JA and Deangelis CD:** Perinatal asphyxia: Introduction hypoxic ischemic encephalopathy, 1999; PP: 235- 237.

**Shroff MM, Soares-Fernandes JP, Whyte H, et al:** MR imaging for diagnostic evaluation of encephalopathy in the newborn. Radiographics. 2010; 30(3):763-80.

**Sutton D:** Intracranial lesions (1). In: Sutton D (ed), A textbook of diagnostic radiology (6th edition), Churchill Livingstone, Inc., NY, Vol. (2) , 1998; pp: 1581-1618.

**Thornberg E, Thiringer K, Odeback A, et al:** Birth asphyxia: incidence, clinical course and outcome in a Swedish population. Acta Paediatr, 1995; 84:927.

**Van Kooij BJ, van Serial M, Nieuvelstein RA, et al:** MRI and neurodevelopmental outcome in 9- to 10-year-old children with neonatal encephalopathy. J Pediatr. 2010; 157(2):221-227.

**Williams k and Galeneau F:** Intrapartum fetal heart rate patterns in the predication of neonatal academia, 2003; 188.

**Wolf R, Robert A, Clancy R, et al :** Quantitative Apparent Diffusion Coefficient Measurements in Term Neonates for Early Detection of Hypoxic-Ischemic Brain Injury: Initial Experience. Radiology 2001, 218: 825-833).

**Yasin Dağ1:** Clinical outcomes of neonatal hypoxic ischemic encephalopathy evaluated with diffusion-weighted magnetic resonance imaging, 2007

**Zanelli A, Dirk P Stanley and David K :** Hypoxic-Ischemic Encephalopathy 2008; PP:136-167.

**Zupan Simunek V:** Birth asphyxia in term newborns: Diagnosis, prognosis, neuroprotection. Arch Pediatr. 2010;17(5):578-82.

## *References*