SUDAN UNIVERSITY OF SCIENCE AND THEHNOLOGY COLLEGE OF GRADUATE STUDIES INSTITUTE OF LASER

USING Nd:YAG LASER IN KELOID TREATMENT, A COMPARATIVE STUDY

A Dissertation submitted as partial Fulfillment for the Requirement of The Degree of Postgraduate Diploma

In LASER Application in Medicine

Dermatology

By: Dr. Ihsan Abdalrhman Abdalla

Supervisors:

Dr. Mubarak Elmahal Ahmed

Table Of Contents

Acknowledgement		I
		Ti
Abstract		
Chapter	One	
		Page
	DUCTION AND BASIC CONCEPT 1-37	
INTRO		1
1.1	Introduction Introduction	1
1.2	What Is Keloid?	2
	Pathophysiology	2
	Wound Healing	3
	Keloid And Hypertrophic Scars	4
1.2.1.3	Keloid And Immune System	4
1.2.2	Epidemiology	5
1.2.3	Dermatopathology	5
1.2.4	Differential Diagnosis	5
1.3	Treatment Of Keloids	5
1.3.1	Conventional Treatment Of Keloid	7
1.4	Laser Basics	7
1.4.1	Laser History	0
1.4.2	Medical Laser Systems Components	15
1.4.3	Characteristics Of Laser Beams	15
1.4.4	Laser Parameters	17
1.5	Laser Tissue Interaction	18
1.5.1	Skin Optics	18
1.5.2	Types Of Laser Tissue Interaction	21
1.5.3	Laser -Induced Interstitial Thermotherapy	25
1.5.4	Selective Photothermolysis	25
1.6	Lasers Used In Dermatology	26
1.6.1	Lasers Used In Treatment Keloid	27
1.7	The Anticipated Laser Hazards	28
1.7.1	Inherent Hazards To Laser Beam	28
1.7.2	Hazards Associated With Operator Work Procedures	29
1.7.2	Hazards Associated With Laser Use	30
1.7.4	Causes Of Laser Accident	31
	Types Of Laser Accident	31
1.7.6	Laser Hazard Classification	31

		Page
1.8	Requirements For Laser Safety	33
1.8.1	General Laser Safety Requirements	33
1.8.2	Specific Laser Safety Requirements	34
1.8.3	Complications Of Laser Surgery	36
1.9.5	Aims and Objective	37
2 47		
Chapter Two SUBJECTS AND METHODS 38 – 44		
2.1	Subjects	38
2.1.1	The Laser Medical System	38
2.1.2	The Patient	40
2.1.3	The Patient Records	41
2.1.4	Laser Safety	41
2.1.5	Other Materials Used	42
2.25	Methods	43
2.2.1	The Laser Surgery	43
2.2.2	Postoperative Care	44
	er Three	
DISCI	SSION AND CONCLUSION 45 - 69	
3.1	Introduction	45
0.1	Patients Records	45
3.2.1	Ages	46
3.2.2	Sex	48
3.2.3	Tribes	50
3.2.4	Site	51
3.2.5	Symptoms	51
3.2.6	Color	53
3.2.7	Duration	54
3.2.8	Size	55
3.3	Treatment	56
3.4	Response Of Keloids To Treatment	5/
3.5	Adverse Sequence and Complications	61
3.6	Laser Hazards	61
3.7	Conclusion and Recommendation	67
	References	0/

,		rage
Fig 1.1	Basic Laser System	9
Fig 1.2	Light Energy Distribution	11

		Page
Fig 1.1	Basic Laser System	9
Fig 1.2	Light Energy Distribution	11
Fig 1.3	Three Level Atomic System	13
Fig 1.4	Four Level Atomic System	13
Fig 1.5	Population Inversion Of Three And Four Level Laser System	14
Fig 1.6	Light Reflection, Absorption Scattering	18
Fig 1.7	Location Of Thermal Effects Inside Biological Tissue	24
Fig 1.8	Laser Labels	34
Fig 1.9	Goggle	37
Fig 2.1	ND: YAG Laser Machine	40
Fig 2.2	Display Panel Of ND: YAG	40
Fig 2.3	Application System And Metal Cannule	44
Fig 2.4	Insearching Of Cannule	44
Table 1.1	Differences Between Kelod And Hypertrophic Scar	2
Table 1.2	Thermal Effect Of Laser	23
Table 3.1	Patient Records	45
Table 3.2	Age	47
Table 3.3	Sex	48
Table 3.4	Tribes	49
Table 3.5	Sight Of Lesion	50
Table 3.6	Associated Symptoms	52
Table 3.7	Color Of The Lesion	53
Table 3.8	Duration Of The Lesion	54
Table 3.9	The Height Of The Lesion	55
Table 3.10	Treatment Of Group One	56
Table 3.11	Treatment Of Group Two	56
Table 3.12	Treatment Of Group Three	57
Table 3.13	Symptoms Response After Treatment	58

F T Briter

Table 3.14	Color Response After Treatment	58
Table 3.15	Texture Response After Treatment	59
Table 3.16	Flexibility Response After Treatment	59
Table 3.17	Size Response After Treatment	59
Table 3.18	Response Comparison Between Three Groups	60
Shape 3.1	Age	47
Shape 3.2	Sex	48
Shape 3.3	Tribe	49
Shape 3.4	Sight Of The Lesion	51
Shape 3.5	Associated Symptoms	52
Shape 3.6	Color Of The Lesion	53
Shape 3.7	Duration Of The Lesion	54
Shape 3.8	Height Of The Lesion	55
Photos:		
Group I		63
Group II		64
Group III		66

Abstract:

Keloids are disfigures wounds repair which cause problem to the patient and challenge for the treating doctors.

In this comparative study which was done in the period from June to July 2007 we used the Nd:YAG 1064 nm for treating keloids and compare it with one of the conventional treatment which is intralesional steroid. 22patients involved in this study divided into three groups. Group 1 received intralesional steroid. Group 2 received steroid plus LASER sessions. Group 3 had LASER treatment only.

Although the results obtained were same for the three groups, and considering the side effects of steroid ,LASER treatment was promising and needs more work up for more patients and longer time .

مرض الجدرة يحدث نتيجة خلل في عملية التئام الجروح الجلدية و يؤدي الى تشوهات التي تؤدي الى مشاكل للمريض و تحدي للطبيب لصعوبة علاجها.

في هذه الدراسة المقارنة التي تمت في الفترة بين مايو الى يوليو من عام ٢٠٠٧، تم العلاج بجهاز الليزر ND:Yag ذو الطول الموجي ١٠٦٤ نانومتر و تمت مقارنته بالعلاج التقليدي للجدرة و هو الحقن داخل الورم بمادة الكورنيزون.

تم اختيار حوالي ٢٢ مريض لهذه الدراسة قسموا الى ثلاث مجموعات. مجموعة تم علاجها بالليزر، و مجموعة تم علاجها بالليزر زائدا الحقن الموضعي بالكورنيزون، و مجموعة تم علاجها بالليزر نتائج العلاج أظهرت تماثلا في المجموعات علاجها بالنظر الى المضاعفات الناتجة من العلاجات الأخرى نجد أن العلاج بالليزر بالنسبة للجدرة علاج واعد و يحتاج الى مزيد من الدراسة لمرضى أكثر و لفتزة علاجية أطول.