Sudan University of Science and Technology College of graduate Studies Institute of laser

Use of Carbon Dioxide Laser (fractional mode 10600 nm) in Treatment of Skin Tags in Sudanese Patients

A Dissertation Submitted For Partial Fulfillment of
Postgraduate Diploma in Laser Applications
In Medicine-Dermatology

By: Nosiba Sayed AwadElkariem MBBS, MD. Dermatology



Supervisors:

Dr.Babikir Osman Elbashir (Ph.D. laser physics)

Dr.ShazaMohammed Yosif (Dermatologist, laser postgraduate Diploma)

ABSTRACT

Skin tags are common benign lesions composed of loose fibrous tissue and occurring mainly on the neck and majorflexures as a small soft pedunculated protrusion. This study aimed to evaluate the efficiency of CO2 laser (fraction mode) for the treatment of skin tags.

Ten patientswere included in this study 2(males) and 8(females). They were complaining of skin tags, beforetreatment laser safety was done. All patient were treated under local anesthesia. The laser system used in this study was the 10600 nm carbon dioxide laser in continuous mode with power 4.5 and 7 Watts. Patients were very satisfied upon the treatment results, because of fewer side effects and less recovery time.

The results obtained were encouraging the use of co2 laser in skin tags treatment because of its greater safety margin and precise ablation, and could replacing the other modalities of treatment for the ablation of many superficial lesions.



ملخص الدراسة

الزوائد الجلدية هي من أكثر الأورام الجلدية الحميدة شيوعا ، وهي تتكون من نسيج ليفي، وتوجد في الرقبة والاماكن ذات الانتناءات كنتوء رخو . هدف هذه الدراسة هو تقييم فعالية اليزر ثاني اوكسيد الكربون في ازالة الزوائد الجدية .

تم اشراك عشره مرضى في هذه الدراسه اثنان من الذكور وثمانيه من الإناث. كان لديهم زوائد جلديه في مناطق مختلفه من أجسامهم ، قبل المعالجه تم عمل كل اجراءات السلامه.

كل المرضى تم حقنهم بمخدر موضعي ، نظام اليزر الذي تم استخدامه هو ١٠٦٠٠ ثاني اوكسيد الكربون بنمط مستمر وقوه تتراوح بين ٧ الى ٢٥٪ واط.

كانت النتائج مرضيه جدا نسبه لقله الاعراض الجانبيه وقصر فتره النَّقاهه.

إن نتائج هذا البحث محفزه جدا على استخدام اليزر ثاني اوكسيد الكربون في ازالة الزوائد الجلديه نسبه لسلامته العاليه ودقته المتناهيه، ويمكنه أن يحل محل طرق المعالجه الأخرى في إزالة الافات الجلديه السطحيه.



Title		Page number
Abstract		I
Arabic abstract		11
Acknowledgment		III
Index		IV
List of figures		VIII
List of tables	7 17*	IX
Chapter one	, .	1
1.1 Introduction		1
1.2 Main hypothesis		1
1.3 Dissertation structure		1
1.4 Histology of the Skin		-2
1.4.1 Epidermis		2
1.4.1.1 Stratum Basalis		3
1.4.1.2 Stratum Spinosum		3
1.4.1.3 Stratum Granulosum		5
1.4.1.4 Stratum Corneum		6
1.4.2 Dermis		7
1.4.3 The subcutis		7
1.5.Skin tags		8
1.5.1Clinical features.		8
.5.2Incidence and aetiology		9
.5.3 Histopathology		10
.5.4 tratment		11

1.6 laser basics	12
1.3.1 History	12
1.3.2Unique properties of lasers	13
1.3.3 Basic components of a laser	14
1.6.4 Spontaneous and stimulated emission	15
1.6.5 Population inversion	17
1.6.6 The resonator	17
1. 7 Laser tissue interactions	18
1.7.1 Optical Properties of Tissue	18
1.7.2 Reaction Mechanisms	18
1.7.3 Non thermal, Chemical Reactions	19
1.7.3.1Biostimulation	19
1.7.3.2 Photodynamic Therapy	20
1.7.4 Thermal Reactions	20
1.8 Laser safety	22
Chapter Two	27
2.Material and methods	27
2.1 Introduction	27
2.2 study area	27
2.3 study time:	27
2.4 study design:	27
2.5 study population:	27
2.5.1 Inclusion criteria:	27

2.5.2 Exclusion criteria:	27
2.6 The patient records	28
2.7 The laser medical system	29
2.8 Safety features of the carbon dioxide system:	30
2.8.2 Laser room precaution:	30
2.8.3 Optical hazards precaution:	30
2.9 Informed consent	30
2.10 The method:	30
Chapter three	32
3. Results & Discussion	32
3.1 Introduction	32
3.2 Results	32
3.2.1 Personal data of the patients	32
3.2.2 Complains	33
3.2.3 Sites and number of the lesions	34
3.2.4 Methods of treatment	34
3.2.5 Evaluation	35
3.3 Discussion	39
3.4 Conclusion	41
3.4 Recommendation	41
4. References	42