# Dedication

To the soul of my father

And

To all those who supported me

#### **ACKNOWLEDGEMENT**

First of all I thank God for blessing, and giving me the strength, patience, and resilience to overcome the difficulties and the obstacles that I faced during this study.

Iam sincerely grateful to Prof .Shadia Abdalati Omer for her supervision, continuous support, patience, help, and kindness.

Special thanks to Dr Shamseldein Ahmed who facilitated the papers publication.

I would like to express my sincere thankfulness to General Ali Hassan the director of the police Directorate for Dogs of the year (2014), Khartoum State who gave me the permission to carry out my work and Dr.Abdein Mohammed for his support, advices and help.

Great thanks to clinicians at police Directorate for Dogs clinic, Dr. Hanaa Abaas Mohammed Ahmed, Dr. Jehan and Dr.Zobaida Ali Awad Alkareem, for their help and kindness in sample collection.

Thanks extend to Lab. Technicians Alaa Eldeen Al Fadil, Hamory Kafei and Amani Abdallah for their assistance and co-operation during blood sampling.

# **List of Contents**

No	Page
Dedication	I
Acknowledgment	II
List of contents	III
List of Tables	V1
List of figures	V11
Abbreviations	V111
Abstract	1X
Arabic Abstract	X1
Introduction	1
Chapter One: Literature Review	5
1.1 Complete Blood count (CBC)	5
1.1.1 Red blood cells	6
1.1.1.1 Erythropoieses	8
1.1.2 Hemoglobin	10
1.1.3 Hematocrit	10
1.1.1 Red blood cells indices	11
1.1.4.1 Mean corpuscular volume determination (MCV)	12
1.1.4.2 Mean corpuscular hemoglobin(MCH)	12
1.1.4.3 Mean corpuscular hemoglobin concentration (MCHC)	12
1.1.4.4 Red cell distribution width (RDW)	13

1.1.5 Leukocytes	13
1.1.5.1 Leukopoiesis	14
1.1.5.2 Neutophil	15
1.1.5.3 Eosinophil	16
1.1.5.4 Basophil	16
1.1.5.5 Lymphocyte	17
1.1.5.6 Monocyte	18
1.1.5.7 Total leukocyte count	19
1.1.6 Platelets	19
1.1.6.1 Platelets indices	20
1.1.7 Red blood cells and biochemical parameters in relation to the	
sex and age	22
1.2 Effect of the season on CBC and biochemical parameters	24
1.3 Exercise	26
1.4 Erythrocyte osmotic fragility	
2.1 Study area	30
2.2 Study design	30
2.3 Metrological data	30
2.4 Animals	31
2.5 Blood collection	31
2.6 Hematological examination	32
2.7 Effect of the season	32

2.8 Effect of exercise on CBC	34
2.8.1Before exercise	34
2.8.2 Immediately after exercise	34
2.8.3 One hour post exercise	34
2.9 Erythrocyte osmotic fragility	34
2.10 Clinical parameters	37
2.11 Statistical analysis	37
Chapter Three: Results	38
3.1 Effect of the sex on CBC	38
3.2 Correlation between platelets count and platelet indices	38
3.3Correlation between platelet indices and parallel RBC	44
3.4 Effect of the season	44
3.4.1 Metrological data	44
3.4.2Effect of the season on red blood cells and indices	44
3.4.3 Effect of the season on white blood cells and Platelets	44
3.4.4 Effect of season on biochemical parameters	44
3.5 Effect of exercise	45
3.6 Erythrocytes osmotic fragility	51
3.7 Clinical parameters	51
Chapter Four: Discussion Conclusion	55 62
Recommendation	64
References	65

# **List of Tables**

No	Page
(1): Red Cells Parameters and Platelets count and Indices in Male	40
and Female of German shepherd Dogs in the Sudan.	
(2): White blood cells parameter in Male and Female of German	41
Shepherd Dogs in the Sudan.	
(3): Correlation between Platelets count and Platelets indices of	42
German Shepherd dogs in the Sudan.	
(4): correlation between Platelet indices and parallel red blood cell	43
Parameters in Male and Female of German shepherd dogs in the Suda	n.
(5): Seasonal variation in the erythrocytes indices of German Shepherd	46
dogs in the Sudan.	
(6): Seasonal variation of the, WBC count and white blood cells	47
Differential count, platelets count and indices of German Shepherd	
dogs in the Sudan.	
(7): Seasonal variation in the concentration of some blood metabolites	48
and serum minerals of German shepherd dogs in the Sudan.	
(8): Effect of exercise on red blood cell parameters of German shepherd	49
dogs in the Sudan.	
(9): Effect of exercise on white blood cell count, differential count and platelets count and indices of German shepherd dogs in the Sudan.	50
(10): Effect of time of the day on some clinical parameters of German Shepherd dogs in the Sudan.	54

# **List of Figures**

No	Page
(1): Over all Erythrocyte Osmotic Fragility of Germen shepherd	52
dogs in the Sudan.	
(2): Effect of sex on erythrocyte Osmotic fragility of Germen	
Shepherd dogs in the Sudan.	

# **ABBREVIATIONS**

Symbol	Abbreviation
ALP	Alkaline phosphatase
ANOVA	Analysis Of Variance
AST	Aspartate amino Transferase
CBC	Complete Blood Count
CD4	Cluster of Differentiation 4
CK	Creatinine kinase
EDTA	Ethylene Diamine Tetra Acetic acid
EOF	Erythrocyte Osmotic Fragility
GGT	Gamma Glutamyl Transferase
Hb	Hemoglobin
HCT	Hematocrit
HPA	Hypothalamic-Pituitary-Adrenalaxis
IMT	Immune Mediated Thrombocytopenia
LDH	Lactate Dehydrogenase
MCH	Mean Cell Hemoglobin
MCHC	Mean Cell Hemoglobin Concentration
MCV	Mean Cell Volume
MHC	Major Histo compatibility Complex
MPV	Mean Platelet Volume
NK	Natural Killer
PCT	Platelet crit
PDW	Platelet Distribution Width
PLT	Platelets
PMN	Poly Morph Nuclear
RBC	Red Blood Cell
RDW	Red cell Distribution Width
SAM	Sympathetic-Adrenal-Medullary axis
SPSS	Statistical Package for Social Science
WBC	White Blood Cell

### **ABSTRACT**

The current study evaluated some hematological and biochemical parameters of German shepherd dogs under Sudan condition as well as the effect of sex, season, and exercise.

Thirty three healthy dogs (14 males, 19 females) aged 2 to 4 years belonging to the Police Directorate for Dogs at Khartoum State were used in this study. Five ml of venous blood were taken from the cephalic vein and CBC analysis was immediately performed using the Sysmex KX2 hematology analyzer, Mindray BS-2000 for Biochemical parameters analysis, and SpectrophotometerAVI -574 for erythrocyte osmotic fragility determination.

Females showed highly significant (p $\leq$ 0.02) value for red cell distribution weidth (RDW) than males (14.68±1.95vs14.14±0.53) whereas, males showed high numerical values of red cell count (6.60±0.81vs6.35±0.74×10<sup>12</sup>), hemoglobin concentration, (Hb) (18.25±2.26vs17.09±3.71 gm/dl), white blood cells count (WBC) (10.49±2.85vs9.41±2.78×10<sup>9</sup>/L), Neutrophil (79.50±13.73vs74.23±13.79%) and Mix cells (52.77±12.01vs50.61±9.42%). Females showed highly significant (p $\leq$ 0.02) value of platelets count (PLT) (193.11±80.21vs170.50±43.78×10<sup>9</sup>/L)and Platelet crit (PCT) (0.14±0.09vs0.11±0.67%) than males.

Highly positive significant (P $\leq$ 0.01) correlations were observed between PDW and MCH, MCV, but highly negative correlation was found between PDW and RDW, RBC in the all dogs. Moreover highly positive significant (P $\leq$ 0.01) correlation was found between PLT and PCT in female and all the dogs. No significant correlations found between PLT and MPV, PDW in the

two sexes. However, no significant correlations were detected between RBC and PLT, PCT, and HCT in the all dogs.

During winter  $PCT(0.120\pm0.07fL),$ PDW(16.03±0.55%) .and Neutrophil(83.16 $\pm$ 15.30%) were significantly( $P \le 0.05$ ) increased, while the  $MPV(9.42\pm0.81)$ , Total protein(6.94±1.02g/dl), and globulin (3.47±8.05g/dl) were decreased. Significant increase (P≤0.05) in PLT (205.58±69.31×10<sup>9</sup>/L), triglyceride (159.37±20.65mg/dl) and urea (44.15±10.60mg/dl) were observed in summer. Albumin significantly (P < 0.05)decreased (3.03±0.51g/dl) and lymphocyte increased (94.45±3.19%) during autumn.

RBC, HCT, MCV were decreased numerically immediately and after one hour of exercise.

Lymphocytes decreased immediately and at one hour post exercise. PLT significantly ( $P \le 0.04$ ) increased ( $118.60 \pm 39.67 \times 10^9 / L$ ) immediately post exercise.

Hemolysis of erythrocyte started at Nacl concentration of 0.55% in all the dogs. Males erythrocytes were more resistant than females.

All the obtained results were within the normal range values, that German shepherd dogs are well adapted to the climate condition of the Sudan.

Effect of sex, season and exercise should be considered in clinical interpretation of dogs' hematological and biochemical variables.

### ملخص الدراسه

قيّمت الدراسة الحالية بعض المعايير الدموية والبيوكيميائية لكلاب الراعي الألماني تحت ظروف السودان وكذلك تأثير الجنس والموسم وممارسة الرياضة.

ثلاث وثلاثون كلباً صحياً (14 ذكر و 19 أنثى) تتراوح أعمار هم بين سنتين و 4 سنوات تابعة للاداره العامه للكلاب الشرطيه بولاية الخرطوم استخدمت في هذه الدراسة .

تم أخذ خمسة مل من الدم الوريدي من الوريد الرأسي وأجري تحليل إجمالي عدد الدم Mindray BS-2000, Sysmex KX2 على الفور باستخدام محلل الدم Spectrophotometer AVI -574 لتحليل المعلمات البيوكيميائية ، و 574- Spectrophotometer AVI كرات الدم الحمراء.

أظهرت الإناث قيمة معنوية عالية ( $p \le 0.02$ ) في حجم توزيع الخلايا الحمراء ( $p \le 0.02$ ) في حجم توزيع الخلايا الحمراء ( $p \le 0.02$ ) مقارنة بالذكور ( $p \le 0.02$ ) ، في حين أظهرت الذكور قيم عددية عالية مقارنة بالذكور ( $p \le 0.02$ ) ، تركيز الهيموجلوبين , ( $p \le 0.02$ ) من عدد الخلايا الحمراء ( $p \le 0.02$ ) خلايا الدم البيضاء ( $p \le 0.02$ ) خلايا الدم البيضاء ( $p \ge 0.02$ ) خلايا الدم البيضاء ( $p \ge 0.02$ )

 $(79.50\pm13.73\text{vs}74.23\pm13.79\%)$  Neutrophil  $(10.49\pm2.85\text{vs}9.41\pm2.78\times10^9\text{/L})$  ( $(p\le0.02)$  المناث قيمة معنوية عالية  $(52.77\pm12.01\text{vs}50.61\pm9.42\%)$  و  $(9\le0.02)$  المناث قيمة معنوية عالية  $(52.77\pm12.01\text{vs}50.61\pm9.42\%)$  و نسبة من عدد الصفائح الدموية ((PLT) المناث المنائح الدموية في العينه (PCT) ((PCT) المناث المناز المناث المناز المن

, PCT ( $0.120\pm0.07fL$ ) من كل من كبيرمعنوي كل من ( $0.120\pm0.07fL$ ) و PDW( $16.03\pm0.55\%$ ) بينما انخفض متوسط حجم PDW( $16.03\pm0.55\%$ )

الصفائح الدمويه (MPV) (MPV), البروتين,(6.94 $\pm$ 1.02g/dl), الدمون (MPV), الدمون (MPV) القلوبيولين (3.47 $\pm$ 8.05g/dl). في حين زادت  $\pm$ 10.60mg/dl) واليوريا (159.37 $\pm$ 20.65mg/dl) بشكل ملحوظ الثلاثية (P $\leq$ 0.05) في فصل الصيف. انخفض بشكل ملحوظ الالبيومين (P $\leq$ 0.05) في فصل الصيف. انخفض بشكل ملحوظ الالبيومين (P $\leq$ 0.05) خلال الخريف.

انخفضت HCT, RBC انخفاض غير معنوي على الفور وبعد ساعة من ممارسة الرياضة.

من ناحية أخرى, انخفضت الخلايا الليمفاوية على الفور وفي ساعة واحدة بعد التمرين و از دادت PLT معنويا ( $P \le 0.04$ ) معنويا ( $P \le 0.04$ ) معنويا ( $P \le 0.04$ )

بدأ انحلال الدم في كريات الدم الحمراء عند تركيز كلوريد الصوديوم بنسبة 51٪ في جميع الكلاب. كانت كريات الدم الحمراء لدى الذكور أكثر مقاومة من الإناث.

جميع النتائج التي تم الحصول عليها كانت ضمن قيم المدى الطبيعي, ذلك أن كلاب الراعى الألماني تتكيف بشكل جيد مع الظروف المناخية للسودان.

ينبغي النظر في تأثير الجنس، والموسم والتمرين في التفسير السريري لمتغيرات الكلاب الدموية والكيميائية الحيوية.