

References

- Abdel-Aale, S. M., and Attia, R. S. (1993): Characterization of black cumin (*Nigella sativa*) seeds. 2- Proteins. *Alex. Sci. Exch.*, **14**, 483– 496.
- Abdel-salam, I. M., Raafat, A. and Abdel-muniem, H. (1998): Radioprotective effect of *Nigella sativa* oil on the liver of Swiss albino mice. *Egypt. J. Biochem.*, **16**(1): 167–76.
- Abdel-sater, K. A. (2009): Garoprotective effects of *Nigella sativa* oil on the formation of stress gastritis in hypothyroidal rats, *Int. J. Physiol. Pathophysiol. Pharmacol.*, **1**: 143–9.
- Aboul ela, M. A., El-shaer, N. S. and Ghanem, N. B. (1996): Antimicrobial evaluation and chromato- graphic analysis of some essential and fixed oils, *Pharmazie*, **51**(12): 993–4.
- Aboutable, E. A., El-azzouny, A. A. and Hammerschmidt, F. J. (1986): Aroma volatiles of *Nigella sativa* L. seeds, in Brunke EJ (ed.) *Progress in Essential Oil Research, Proceedings of The International Symposium on Essential Oils*. de Gruyter, Berlin, 44–55.
- Adamu, H. M., Ekanem, E. O. and Bulama, S. (2010): identification of essential oil components from *Nigella sativa* seed by gas chromatography mass spectroscopy, *Paki. J. Nutri.*, **9**(10): 966–7.
- Agarwal, R., Kharya, M. D. and Shrivastava, R. (1979): Antimicrobial and anthelmintic activities of the essential oil of *Nigella sativa* Linn., *Indian J. Exp. Biol.*, **17**: 1264–5.
- Ahmad, J., Khan R. A. and Malik M. A. C (2009): Study *Nigella sativa* oil in the management of wheeze associated lower respiratory tract illness in children, *Afr. J. Pharm. Pharmacol.*, **3**(5): 248–51.
- Alhaj, N. A, Shamsudin, M. N, Zamri, H. F and Abdullah, R. L (2008): Extraction of essential oil from *Nigella sativa* using supercritical carbon dioxide: study of antibacterial activity, *Am. J. Pharmacol. Toxicol.*, **3**(4): 225–8.
- Ali, A., Abu sayeed, M., Shahinvr, M., Sayeamin, A., Khan, A.M. and Ida, I. M. (2012): Characteristics of oils and nutrient contents of *Nigella sativa* Linn and *Trigonella Foenum Graccum* seeds. *Chem. Society of Ethiopia.*, **26**(1):55-64.
- Akhtar, M. S. and Aslam, M. (1997): Anticestodal principles of *Nigella sativa* Linn. (Kolanji) seeds, *Pak. J. Pharmacol.*, **14**(2): 7–14.

Akhtar, M. S., and Riffat, S. (1991): Field trials of *Saussurea lappa* roots against nematodes and *Nigella sativa* seeds against cestodes in children, *J. Pak. Med. Assoc.*, **41**: 185–7.

Alsamarai, A. M., AlObaidi, A. H. A, (2012): Evaluation of Therapeutic Efficacy of *Nigella sativa* (Black Seed) for Treatment of Allergic Rhinitis. In: Kowalski ML, ed. *Allergic Rhinitis*. Rijeka, Croatia: Intech; 197-214.

Al-yahya, M. A. (1986): Phytochemical studies of the plants used in traditional medicine of Saudi Arabia, *Fitoterapia*, **57**(3): 179–82.

Anon. (1998): *PFA Specifications*, up to 1997–98, DGHS, Ministry of Health, Government of India, A.05.10.01-10.

Ansari, A. A., Osman, S. M and Subbarama, M. R. (1975): Component acids of minor seed oils. 1., *J. Oil Technol. Assoc. India.*, **7**(1): 26–7.

Aqal, B. (1992): The calcium antagonistic effect of the volatile oil of *Nigella sativa* seeds, *Dirasat-Univ. Jordan, Ser. B.*, **19B**(1), 119–33.

Ansari, Z., Satish, T. (2013): Traditional uses of *Nigella sativa*, in Malegaon region of Nashik — a review. *Int J Pure App Biosci.* **1**(2):19-23.

Aqel, M. and shaheen, R. (1996): Effects of the volatile oil on the uterine smooth muscle of the anti-inflammatory activity of *Nigella sativa L.*, rat and guinea pig, *J. Ethnopharmacol.*, **52**: 23–6.

Argon, Z. V. (2016): Determination of Physicochemical Properties of *Nigella sativa* Seed Oil from Balıkesir Region, Turkey. *Chem. And process Eng. Research.*, **41**: 2224-7467.

Assayed, M. E. (2010): Radioprotective effects of black seed (*Nigella sativa*) oil against hemopoietic damage and immunosuppression in gamma-irradiated rats. *Immunopharmacol. Immunotoxicol.*, **32**: 284-296.

Atta, M. and Imaizumi, K. (1998): Antioxidant activity of Nigella (*Nigella sativa* L.) seeds extracts, *Nihon Yukagakkaishi*, **47**(5): 475–80.

Atta, M. B. (2003): Some characteristics of nigella seed cultivated in Egypt and lipid profile, *Elsevier food chem.*, **83**:63-68.

- Awney, H. A., Amara, A., El-masry, M. H., and Baghdadi, H. H. (1997): Effect of 12 plant extracts on hepatic microsomal benzo[a]pyrene hydroxylation and hydrogen peroxide production in mice treated with lindane, *Environ. Nutr. Interact.*, **1**(3–4): 129–42.
- Bashandy, S. A. E. (1996): Effect of *Nigella sativa* oil on liver and kidney functions of adult and senile rats, *Egypt. J. Pharm. Sci.*, **37**(1–6): 313–27.
- Babayan, V. K., Koottungal, D., and Halaby, G. A. (1978): Proximate analysis, fatty acid and amino acid composition of *Nigella sativa* L. Seeds. *Journal of Food Science*, **43**: 1315–1319.
- Bin Sayeed, M. S., Asaduzzaman, M., Morshed, H., Hossain, M. M., Kadir, M. F., Rahman, M. R. (2013) The effect of *Nigella sativa* Linn. seed on memory, attention and cognition in healthy human volunteers. *Journal of Ethnopharmacology*. **148**(3):780-786.
- Bin Sayeed, M. S., Shams, T., Fahim Hossain, S(2014): *Nigella sativa* L. seeds modulate mood, anxiety and cognition in healthy adolescent males. *Journal of Ethnopharmacology*. **152**(1):156-162.
- Blaschek, W. E. D. (2016): *Wichtl — Teedrogen und Phytopharma — Ein Handbuch für die Praxis*. 6th ed. Stuttgart, Germany: Wissenschaftliche Verlagsgesellschaft.
- Boskabady, M. H., Farhadi, J. (2008): The possible prophylactic effect of *Nigella sativa* seed aqueous extract on respiratory symptoms and pulmonary function tests on chemical war victims: a randomized, double-blind, placebo-controlled trial. *J Altern Complement Med.* **14**(9):1137-1144.
- Bourrel, C., Vilrem, G. and Perinean, E. (1993): Aromatic compounds of *Nigella sativa* L. Grains; evaluation of antibacterial and antifungal properties, *Riv. Ital. EPPOS*, **4**(10), 21–7.
- Buriro, M. A. and Tayyeb, M. (2007): Effect of *Nigella sativa* on lipid profile in albino rats, *Gomal J. Med. Sci.*, **5**(1): 28–31.
- Cheikh-Rouhou, S., Besbes, S., Hentati, B., Bagheri, C., Deroanne, C. and Attia, H. (2007): *Nigella sativa* – chemical composition and physicochemical characteristics of lipid fraction, *J. Food Chem.*, **101**: 673–81.

- Daba, M. H., and Abdel-rahaman, M. S. (1998): Hepatoprotective activity of thymoquinone in isolated rat hepatocytes, *Toxicol. Lett.*, **95**(1): 23–9.
- Dubey, P. N., Balraj, S., Mishra, B. K. ant, K. K. and Solanki, R. K. (2016): *Nigella (Nigella sativa)*: a high value seed spice with immense medicinal potential, *Indian Journal of Agricultural Sciences* **86** (8): 967–79.
- El-alfy, T. S., El-faty, H. M. and Toama, M. A. (1975): Isolation and structure assignment of an antimicrobial principle from the volatile oil of *Nigella sativa* L. seeds, *Pharmazie*, **30**: 109–11.
- El-dakhakhny, M., Barakat, M., El-halim, M. A. and Aly, S. M. (2000): Effect of *N. sativa* oil on gastric secretion and ethanol induced ulcer in rats, *J. Ethnopharmacol.*, **72**(1/2): 299–304.
- El-daly, E. S. (1998): Protective effect of cysteine and vitamin E, *Crocus sativa* and *Nigella sativa* extracts on cisplatin-induced toxicity in rats, *J. Pharm. Belg.*, **53**(2): 87–95.
- El-Mahdy, M. A., Zhu, Q., Wang, Q. E., Wani, G. and Wani, A. A. (2005): Thymoquinone induces apoptosis through activation of caspase-8 and mitochondrial events in p53-null myeloblastic leukemia HL-60 cells. *Int. J. Cancer*, **117**: 409-417.
- El-sayed, A. A., Hussiney, H. A. and Yassa, A. I. (1997): Constituents of *Nigella sativa* oil and evaluation of its inhibitory effect on growth and aflatoxin production by *Aspergillus parasiticus*, *Dtsch. Lebensm.-Rundsch.*, **93**(5): 149–52.
- El-shayeb, N. M. A. and Mabrouk, S. S. (1984): Utilization of some edible and medicinal plants to inhibit aflatoxin formation, *Nutr. Rep. Int.*, **29**(2): 273–82.
- El-tahir, K. E. H. and Ageel, A. M. (1994): Effect of the volatile oil of *Nigella sativa* on the arterial blood pressure and heart rate of the guinea pig, *Saudi Pharm. J.*, **2**(4): 163–8.
- El-tahir, K. E. H., Ashour M. M. S. and Al-harbi, M.M. (1993): The respiratory effects of the volatile oil of the black seed in guinea pigs – elucidation of the mechanism of action, *Gen. Pharmacol.*, **24**: 1115–22.
- Engel, G. and Brinckmann, J. (2017): Nigella, *Herbal Gram American Botonical council*, **114**: 8-16.
- Farhangi, M. A, Dehghan, P., Tajmiri, S., Abbasi, M. M. (2016): The effects of *Nigella sativa* on thyroid function, serum vascular endothelial growth factor (VEGF)-1, nesfatin-1 and

anthropometric features in patients with Hashimoto's thyroiditis: a randomized controlled trial.
BMC Complement Altern Med; **16**(1):471.

Ferdous, A. J., Islam, S. N., Ahsan, M., Hasan, C. M. and Ahmed, Z. U. (1992): *In vitro* antibacterial activity of the volatile oil of *Nigella sativa* seed against multiple drug resistant isolates of *Shigella* spp and isolates of *Vibrio-cholerae* and *Escherichia-col*, *Phytother. Res.*, **6**: 137–40.

Gad, A. M., El-Dakhakhny, M., and Hassan, M. M. (1963): Studies on the chemical composition of Egyptian *Nigella sativa* L. oil. *Planta Medica*, **11**: 134–138.

Gali-Muhtasib, H., El-Najjar, N., Schneider-Stock, R. (2006): The medicinal potential of black seed (*Nigella sativa*) and its components, **2**:133-153.

Gheita, T. A., Kenawy, S. A. (2012): Effectiveness of *Nigella sativa* oil in the management of rheumatoid arthritis patients: a placebo controlled study. *Phytother Res.* **26**(8):1246-1248.

Ghorbanibirgani, A., Khalili, A., Rokhfrooz, D. (2014): Comparing *Nigella sativa* oil and fish oil in treatment of vitiligo, *Iran Red Crescent Med J*, **16**(6):4515.

Hadi, V., Kheirouri, S., Alizadeh, M., Khabbazi, A., Hosseini, H. (2016): Effects of *Nigella sativa* oil extract on inflammatory cytokine response and oxidative stress status in patients with rheumatoid arthritis; a randomized, double-blind, placebo-controlled clinical trial. *Avicenna J Phytomed.*, **6**(1):34.43.

Hailat, N., Bataineh, Z., Lafi, S., Raweily, E., Aqel M., A1-katib, M. and Hanash, S. (1995): Effect of *Nigella sativa* volatile oil on Jurkat T-cell leukemia polypeptides, *Int. J. Pharmacog.*, **33**: 16–20.

Hannan, A., Saleem, S., Chaudhary, S., Barkat, M. M. and Arshad, M. U. (2008): Anti-bacterial activity of *Nigella sativa* against clinical isolates of methicillin resistant *Staphylococcus aureus*, *J. Ayub. Med. Coll Abbottabad*, **20**(3):72–4.

Haq, A., Abdullatif, M., Lobo, P.I., khabar, K. S. A., Sheth, K. V. and Al-sedairy, S. T. (1995): *Nigella sativa*— effect on human lymphocytes and polymorphonuclear leukocyte phagocyticactivity, *lmmunopharmacol.*, **30**: 147–55.

Hashimoto's, (2016): thyroiditis: a randomized controlled trial. *BMC Complement Altern Med.* **16**(1):471.

Heiss, A. G., Stika, H. P., De Zorzi, N., Jursa, M. (2012/13): *Nigella* in the Mirror of Time: A Brief Attempt to Draw a Genus' Ethnohistorical Portrait. *Offa-Zeitschrift. Berichte und Mitteilungen zur Urgeschichte, Frühgeschichte und Mittelalterarchäologie.* **69/70**:147–169.

Houghton, P. J., Zarka, R., Delasheras, B. and Hoult, J. R. S. (1995): Fixed oil of *Nigella sativa* and derived thymoquinone inhibit eicosanoid generation in leukocytes and membrane lipid peroxidation, *Planta Med.*, **61**: 33–42.

Huseini, H. F, Kianbakht, S., Mirshamsi, M.H., Zarch, A. B. (2016): Effectiveness of topical *Nigella sativa* seed oil in the treatment of cyclic mastalgia: A randomized, triple-blind, active, and placebo-controlled clinical trial. *Planta Med.*, **82**(4):285-288.

Ibraheem, N. K., Ahmed, J. K. and Hassan, M. K. (2010): The effect of fixed oil and water extracts of *Nigella sativa* on sickle cells: an in vitro study, *Singapore Med. J.*, **51**(3): 230–4.

Ibrahim, Z. S., Ishizuka, M., Soliman, M., ElBohi, K., Sobhy, W., Muzandu, K., Elkattawy, A. M., Sakamoto, K. Q. and Fujita, S. (2008): Protection by *Nigella sativa* against carbon tetrachloride-induced down regulation of hepatic cytochrome P450 isozymes in rats. *Jpn. J. Vet. Res.*, **56**: 119-128.

Isik, H., Cevikbas, A., Gurer, U. S, (2010): Potential adjuvant effects of *Nigella sativa* seeds to improve specific immunotherapy in allergic rhinitis patients. *Med Princ Pract.* **19**(3):206-211.

Jha, T. B., and Roy, S. C. (1979): Rhizogenesis from *Nigella sativa* protoplasts, *Protoplasma.*, **101**: 139–42.

Khan, M.A. (1999): Chemical composition and medicinal properties of *Nigella sativa* L., *Inflammopharmacol.*, **7**(1): 15–35.

Khan, M. A. U., Ashfaq, M. K., Zuberi, H. S., Mahmood, M. S. and Gilani, A. H. (2003): The *in vivo* anti-fungal activity of the aqueous extract from *Nigella sativa* seeds, *Phytother. Res.*, **17**(2): 183–6.

Khan, M. A., Chen, H., Tanina, M., and Zhang, D. (2011): Anticancer activites of *Nigella sativa* (Black cumin). *Afr J Tradit complement Altern Med.* **8**(s): 226- 232

Kokoaka, L., Havlik, J., Valterova, I., Sov, H., Sajfrtova, H. andI. jankovsha, I. (2008): Comparison of Chemical Composition and Antibacterial Activity of *Nigella sativa* Seed Essential Oils Obtained by Different Extraction Methods, *Journal of Food Protection*, **71**, 12: 2475–2480

Kumar, B. H., and Thakur, S. S. (1989): Effect of certain edible oils on growth regulation in *Dysdercus similis*, *J. Anim. Morphol. Physiol.*, **36**(2): 209–17.

Labib, R. M., Osama, A. B., Hafez, H. F. and Abd el-hamid, M. A. M. (2009): *Nigella sativa* L. oil ameliorates methotrexate-induced intestinal toxicity, *Gastrointestinal Oncology Conference*, 1–3 October, Cairo, Egypt.

Ma, C., Nakabayashi, T., Miyashiro, H., Hattori, M., El-meckawy, S., Namba, T., and Shimotohno, K. (1994): Screening of traditional medicines for their inhibitory effects on human immuno- deficiency virus protease, *Wakan lyakugaku Zasshi*, **11**(4): 416–17.

Mahfouz, M., and El-dakhakhny, M. (1966): The isolation of a crystalline active principle from *Nigella sativa*, *Pharm. Sci. United Arab Rep.*, **1**: 9–19.

Medenica, R., Mukerjee, S., Huschart, T., Koffskey, J. and Corbitt, W. (1993): *Nigella sativa* plant extract increases number and activity of immune competent cells in humans, *Exp. Hematol.*, **21**: 1186.

Mabrouk, G. M., Moselhy, S. S., Zohny, S. F., Ali, E. M., Helal, T. E., Amin, A. A. and Khalifa, A. A. (2002): Inhibition of methylnitrosourea (MNU) induced oxidative stress and carcinogenesis by orally administered bee honey and *Nigella* grains in Sprague Dawely rats. *J. Exp. Clin. Cancer Res.*, **21**: 341-346.

Mahdavi, R., Namazi, N., Alizadeh, M., Farajnia, S. (2015): Effects of *Nigella sativa* oil with a low-calorie diet on cardiometabolic risk factors in obese women: a randomized controlled clinical trial. *Food Funct.*, **6**(6):2041-2048.

Mahdavi, R., Namazi, N., Alizadeh, M., Farajnia, S. (2016): *Nigella sativa* oil with a calorie-restricted diet can improve biomarkers of systemic inflammation in obese women: A randomized double-blind, placebo-controlled clinical trial. *J Clin Lipidol*, **10**(5):1203-1211.

Malhotra, S. K. (2012): Hand book of Herbs and speices. *Indian council of Agricultural Research*, India: 393-409.

Moretti, A., Antuono, D., Filippo, L., and Elementi, S. (2004): Essential oil of *Nigella sativa* L. and *Nigella damascena* L. seed, *J. Essent. Oil Res.*, **16**(13): 182–3.

Morikawa, T., Xu, K., Kashima, Y., Matsuda, H., Ninomiya, K. and Yoshikawa, M. (2004a): Novel dolabel- lane type diterpene alkaloid with lipid metabolism promoting activities from the seeds of *Nigella sativa*, *Org. Lett.*, **6**: 869–72.

Morikawa, T., Xu F., Ninomiya, k., Matsuda, H. and Yoshikawa, M. (2004b): Nigellamines A3, A4, A5 and C ne dolabellane type diterpene alkaloid with lipid metabolism promoting activities from the Egyptian medicinal food black cumin, *Chem. Pharm. Bull.*, **52**: 494–7.

Musa, D., Dilsiz, N., Gumushan, H., Ulakoglu, G. and Bitiren, M. (2004): Antitumor activity of an ethanol extract of *Nigella sativa* seeds, *Biologia Bratislava*, **59**(6): 735–40.

Mutabagani, A. and El-mahdy, S. A. M. (1997): Study of anti-inflammatory activity of *Nigella sativa* L. and thymoquinone in rats, *Saudi Pharm. J.*, **5**(2–3): 110–13.

Muthu kumara, S. S. and Huat kwong, B. T. (2001): Extraction, isolation and characterisation of antitumor principle, α -hederin, from the seeds of *Nigella sativa*, *Planta Med.*, **67**(1): 29–32.

Madkarni, K. M. (2001): *Indian Plants and Drugs with their Medicinal Properties and Uses*. Asiatic Publishing House, Delhi: 259–60.

Magwa, M., El-sawi, and Hana, G. M. (2006): Effect of *nigella sativa* and activated charcoal as anti- oxidant on verrucarin induced hepatotoxicity in male rats, *Adv. Phytomed.*, **2**: 133–53.

Mair, S. C., Salomi, M. J., Panikkar, B. and Panikkar, K. R. (1991): Modulatory effects of *Crocus sativus* and *Nigella sativa* extracts on cis-platin induced toxicity in mice, *J. Ethnopharmacol.*, **31**, 75–83.

Nielsen, S. S. (2010): Food analysis, 4th edition, west lafayette: USA, pp 241-255.

Nikakhlagh, S., Rahim, F., Aryani, F. H., Syahpoush, A., Brougerdnya, M. G., Saki, N., (2011): Herbal treatment of allergic rhinitis: the use of *Nigella sativa*. *Am J Otolaryngol.* **32**(5):402-407.

Namba, T., Tsunezuku, M., Dissanayake, D. M. R. B., Pilapitiya, U., Saito, K., Kakiuchi, N. and Hattori, M. (1985): Studies on dental caries prevention by traditional medicines (Part VII). Screening of Ayurvedic medicines for anti-plaque action, *Jpn. J. Pharmacognosy*, **39**(2): 146–53.

Namazi, N., Mahdavi, R., Alizadeh, M, Farajnia, S. (2015): Oxidative stress responses to *Nigella sativa* oil concurrent with a low-calorie diet in obese women: A randomized, double-blind controlled clinical trial. *Phytother Res*, **29**(11):1722-1728.

Naqvi, S. A. H., khan, M. S. Y. and Yohora, S. B. (1991): Anti-bacterial, anti-fungal and anthelmintic investigations on Indian medicinal plants, *Fitoterapia*, **62**: 221–8.

Nickavara, B., Mmojab, F., Javidniab, K. and Amolia, M. A. R. (2003): Chemical composition of the fixed and volatile oils of *Nigella sativa* L. from Iran, *Z. Naturforsch.*, **58c**: 629–31

Noor, N. A. and Mourad, M. (2010): Evaluation of antioxidant effect of *Nigella sativa* oil on mono- sodium glutamate induced oxidative stress in rat brain, *J. Am. Sci.*, **6**(12): 13–18.

O'mahony, R., Al khtheeri, H., Weerasekera, D., Fernando, N., Vaira, D. and Holton, J. (2005): Bacterial and anti-adhesive properties of culinary and medicinal plants against *Helicobacter pylori*, *World J. Gastroenterol.*, **11**: 7499–507.

Omar, A., Ghosheh, S., Abdulghani, A., Houdi, A. and Crookscor, P. A. (1999): High performance liquid chromatographic analysis of the pharmacologically active quinones and related compounds in the oil of the black seed, *J. Pharm. Biomed. Anal.*, **19**: 757–62.

Ottai, M. E. S, Teixeira, D. A., Silva, J. A, Mahfouze, S. A., (2012): Phenotypic and chemotypic variation of four nigella (*Nigella sativa*) varieties. Medicinal and Aromatic Plant Science and Biotechnology, **6**(1):40-45.

Oysu, C., Tosun, A., Yilmaz, H. B., Sahin-Yilmaz, A., Korkmaz, D., Karaaslan, A. (2014): Topical *Nigella sativa* for nasal symptoms in elderly. *Auris Nasus Larynx*. **41**(3):269-272.

Pearson, D. (1970): Chemical analysis of food, 6: pp510- 528.

Pruthi, J. S. (2001): *Minor Spices and Condiments*. ICAR, New Delhi.

- Rahman, A., Malik, S., Sadiq, H. S, Iqbal, C. M., Ni c-z, and Clardy, J. (1995): Nigellidine – a new indazole alkaloid from the seeds of *Nigella sativa*, *Tetrahedron Lets.* **36**(12): 1993–6.
- Rajkapoor, B., Anandan, and Jayakar, B. (2002): Anti-ulcer effect of *Nigella sativa* L. against gastric ulcers in rats, *Curr. Sci.*, **82**(2): 177–9.
- Ramadan, R. F and Morsel, J. T. (2002): Characterization of phospholipid composition of black cumin (*Nigella sativa* L.) seed oil, *Nahrung.*, **46**(4): 240–4.
- Rathee, P. S., Mishra, S. H. and Kaushal, R. (1982): Antimicrobial activity of essential oil, fixed oil and unsaponified matter of *Nigella sativa* L., *Indian J. Pharma.*, **44**: 8–10.
- Rchid, H., Nmila, R., Bessière, J. M., Sauvaire, Y. and Chokairi, M. (2004): Volatile components of *Nigella damascene* L. and *Nigella sativa* L. seeds, *J Essent. Oil Res.*, **16**(6): 585–7.
- Rooney, S. and Ryan, M. F. (2005): Modes of action of alpha-hederin and thymoquinone, active constituents of *Nigella sativa*, against HEp-2 cancer cells. *Anticancer Res.*, **25**: 4255-4259.
- Sahebkar, A., Beccuti, G., Simental-Mendia, L.E, Nobili, V. and Bo, S. (2016): *Nigella sativa* (black seed) effects on plasma lipid concentrations in humans: A systematic review and meta-analysis of randomized
- Salem, M. A. (2001): Effect of some heat treatment on nigella seeds characteristics. 1-Some physical and chemical properties of nigella seed oil. *J. Agric. Res. Tanta Univ.*, **27**: 471–486.
- Salem, A. M., Bamosa, A. O., Qutub, H (2017): Effect of *Nigella sativa* supplementation on lung function and inflammatory mediatorsin partly controlled asthma: a randomized controlled trial. *Ann Saudi Med.* **37**(1):64-71.
- Salim, E.I. and Fukushima, S. (2003): Chemopreventive potential of volatile oil from black cumin (*Nigella sativa* L.) seeds against rat colon carcinogenesis. *Nutr. Cancer*, **45**: 195-202.
- Salomi, N. J., Nair, S. C., Jayawarhanan, K. K., Varghese, C. D. and Panikkar, K. R. (1992): Antitumour principles from *Nigella sativa* seeds, *Cancer Lett.*, **63**: 41–6.

- Shah, A. H., Katibi, A., Ageel, A. M. and Tanira M. O. H. (1988): Phytochemical and antimicrobial screening of some plants used in Saudi folk medicine, *Pak. J. Pharm. Sci.*, **1**(1): 53–60.
- Sharma, N. K., Ahirwar, D., Jhade, D. and Ggupta, S. (2009): Medicinal and pharmacological potential of *Nigella sativa*: A review, *Ethnobot. Rev.*, 946–55.
- Shariq, I. M., Israil, A. M., Iqbal, A., Brijesh, P. (2015): Morpho-physiological characterization of seeds and seedlings of *Nigella sativa* Linn.: Study on Indian germplasm. *International Research Journal of Biological Sciences*. **4**(4):38-42.
- Shimada, K., Fujikawa, K., Yahara, K., Nakamura, T. (1992): Antioxidative properties of xanthan on the antioxidation of soybean oil in cyclodextrin emulsion. *J Agric Food Chem.*, **40**:945-8.
- Solati, Z., Baharin, B. S., and Bagheri, H. (2014): Antioxidant property, Thymoquinone content and chemical characteristics of different extracts from *Nigella sativa* L. seeds, *J.of the Amer. Oil Chem. Soc.*, 91:295-300
- Takruri, H. R. and Dameh M. A. H. (1993): Study of nutritional value of black cumin seeds (*N. sativa*), *J. Sci. Food Agric.*, **76**: 404–10.
- Thabrew, M. I., Mitry, R R., Morsy, M. A. and Hughes, R. D. (2005): Cytotoxic effects of a decoction of *Nigella sativa*, *Hemidesmus indicus* and *Smilax glabra* on human hepatoma HepG2 cells. *Life Sci.*, **77**: 1319-1330.
- Toppozada, H., Mazloum, H. and El-dakhgakhny, M. (1965): The antibacterial properties of *Nigella sativa* seeds. Active principle with some clinical applications, *J. Egypt. Med. Assoc.*, **48**, 187–202.
- Turkay, S., Burford, M. D., Sangun, M. K., Ekinci, E., Bartle, K. D. and Clifford, A. A. (1996): Deacidification of black cumin seed oil by selective supercritical carbon dioxide extraction, *J. Amer. Oil Chem. Soc.*, **73**: 1265–70.
- Üstün, G., Kent, L., Çekin, N., and Clvelekoglu, H. (1990): Investigation of the technological properties of *Nigella sativa* (Black Cumin) seed oil. *JAOCs*, **67**: 958–960.
- Vavilov, N. I., Dorofeev, V. F., (1992): *Origin and Geography of Cultivated Plants*. Cambridge, U K : Cambridge University Press .

- Weiss, E. A. (2002): *Spices Crops*. CABI, Wallingford, 356–60.
- Yildiz, F., Coban, S., Terzi, A., Ates, M., Aksoy, N., Cakir, H., Ocak, A. R. and Bitiren, M. (2008): *Nigella sativa* relieves the deleterious effects of ischemia reperfusion injury on liver, *World J. Gas-troenterol.*, **14**(33): 5204–9
- Zawahry, M. R. (1963): Isolation of a new hypotensive fraction from *N. Sativa* seeds, *Kongr. Pharm. Wiss. Vortr. Origenatitt*, **23**: 193–203.