

	Page No.
CONTENTS	
Acknowledgements	
CONTENTS	1
List of Tables	V1
List of Figures	V111
Abstract	X
Abstract in Arabic	X1
CHAPTER ONE: INTRODUCTION	1
1.1 Oil seeds production in Sudan	1
1.2 Importance of oil seeds	2
CHAPTER TWO: LITERATURE REVIEW	5
2.1 Black mahlab	5
2.1.1 Origin and habitats	5
2.1.2 Climatic requirement	7
2.1.3 Planting and Cultivation	7
2.1.4 Uses and medicinal values	8
2.2 White mahlab	9
2.2.1 Plant description and cultivation	9
2.2.2 General uses	11
2.2.3 Medicinal and other uses	13
2.3 Importance of fats and oils	13
2.4 Chemical composition of fats and oils	14
2.4.1 Triglycerides	14
The minor components	15
Mono- and diglycerides	15
Free Fatty acids	16

Phosphatides	16
Sterols	16
Tocopherols and Tocotrienols	17
Pigments	17
Fatty Alcohols	17
Fatty acids	17
Classification of Fatty Acids	18
Saturated Fatty Acids.	18
Unsaturated Fatty Acids	18
Polyunsaturated Fatty Acids	19
Factors affecting physical characteristics of fats and oils	20
Degree of Unsaturation of Fatty Acids	20
Length of Carbon Chains in Fatty Acids	20
Physiochemical Properties of oil	21
Color	21
Refractive Index (RI)	21
Specific Gravity	22
Peroxide Value (PV):	22
Saponification Value (S.V)	23
Acid Value (AV)	23
Unsaponifiable Matter	23
Iodine Value (IV)	24
Protein	24
Amino acids profile	25
CHAPTER THREE: MATERIALS AND METHODS	26
Materials	26
Approximate chemical analysis	26

Moisture and volatile matter	26
Ash content	27
Crude protein	27
Crude fiber	28
Oil content	29
Oil physicochemical analysis	30
Physical parameters	30
Color	30
Specific Gravity (Sp.gr)	30
Refractive Index (R.I)	30
Chemical Parameters	31
Acidity (free fatty acid and acid value)	31
Acid Value	31
Saponification Value (S.V)	32
Iodine value (IV)	32
Unsaponifiable Matter	32
Peroxide Value (P.V)	32
Fatty acids composition by Gas Liquid Chromatography	33
Fatty acids extraction	33
Preparation of Methyl Ester for GC	33
Amino acid composition by Amino Acid Analyzer	34
Preparation of hydrolysate sample	34
Sensory evaluation of the two types of mahlab oils	34
Hedonic test	35
CHAPTER FOUR: RESULTS AND DISCUSSION	36
	36

Seed Characteristics and Composition	36
Seed description	36
Hundred Seeds Weight	36
Approximate chemical analysis	38
Physiochemical analysis of oil	39
Color	40
Fatty Acids Composition	42
Amino acid profile	46
Sensory evaluation of mahlab oil	47
CHAPTER FIVE: CONCLUSIONS & RECOMMENDATIONS	47
Conclusions	48
Recommendations	49
REFERENCES	56
APPENDICES	

List of tables

Table (4.1): Approximate chemical analysis of white and black mahlab seeds

Table (4.2) :) Physiochemical analysis of oil of white and

black mahlab seeds	37
Table (4.3):) Physiochemical analysis of some conventional Sudanese oils	38
Table (4.4): Fatty acids composition of the two mahlab oils	
Table (4.5):) Fatty acid composition (% of total) of white mahlab oils	40
Table (4.6): Amino acid composition of Black and White mahlab seed	41 42
Table (4.7): amino acid composition of five different foods	
Table (4.8): Sensory evaluation analysis of two mahlab oils oil compared with conventional oils	44
	45
	46

List of Figures

Figure 1: Black mahlab flowers and plant	6
Figure 2: Black mahlab seeds	7
Figure 3 : Flowering white mahlab tree	10
Figure 4 : White mahlab fruits	11

Figure 5 : White mahlab seeds	12
Figure 6 : Diagrams of simple and mixed triglyceride	15 16
Figure 7 : Diagrams of mono- and diglycerides	18
Figure 8 : Saturated and unsaturated linkages	19
Figure 9 : conjugated and non-conjugated bonds	

Abstract

Seeds from two varieties of mahlab, black and white mahlab were obtained from the local market Khartoum North, Khartoum State. They were then subjected to different tests for chemical analysis, and the extracted oil was subjected to various tests for physico-chemical analysis according to the standard methods.

Hundred seed weights of the two mahlab varieties were found to be (1.88g) , (4.13g) in black and white mahlab, moisture content (6.03%), (6.22%), ash content (2.81%), (2.12%) , protein (12%) , (28%) , (fiber 2.5%) , (1.87%) , oil content was found to be 13.15%, 30.95% in black and white mahlab seeds respectively. Saponification values in the two varieties were varied from 184.23 in white mahlab to 180.3 in black mahlab. Refractive index was found to be (1.4700), (1.4975), Specific Gravity (0.8167), (0.8511), free fatty acids (3.65%), (3.93%), acid value (7.30), (7.86) mg KOH/g, peroxide value (4.43), (2.54) meq O₂ / Kg oil and Unsaponifiable matter (0.66%), (0.92%) in black and white mahlab oil respectively. The major fatty acids were found to be Palmitic (4.5%), Stearic (16.0%), Oleic (48.3%) and Linoleic (31.6%) in black mahlab oil, while in white mahlab oil was found to be as follows palmitic (5.7%), oleic (46%), and linoleic acid (48%).

Protein content was found to be 21 and 28% in black and white seeds respectively. The total amounts of amino acids

were found to be 12.55 and 18.38 g / 16 g N in black and white seeds respectively. The percentage of essential amino acids in black mahlab seeds was 39.5% and in white mahlab seeds was 40.5%.

From the sensory evaluation analysis of the two mahlab oils, the results revealed that there was a significant difference ($P \leq 0.05$) between the average scores for color, odor and taste of the two mahlab oils and that white mahlab oil seemed to be more acceptable compared to the black mahlab oil by panelists.

الخلاصة

دراسة صفات بذور نوعين من المحلب، المحلب الأسود والمحلب الأبيض لتحديد نسب كل من الزيت والأحماض الدهنية والأحماض الأمينية.

جميع الاختبارات التي أجريت على العينتين تمت بالطرق القياسية، لقد تم تقدير تقريبي لوزن المئة حبة فكان 1.88 g لبذور المحلب الأسود و 4.13 g لبذور المحلب الأبيض. تم تقدير تقريبي للمحتوى الكيميائي للبذور والذي شمل: تقدير المحتوى الرطوبي الذي وجد 6.03%، 6.22%، الرماد 2.81%، 2.12%، الألياف 2.5%، 1.87%، البروتين 21%، 28%، الزيت 13.15%، 30.95% لكل من بذور المحلب الأسود والأبيض على التوالي.

وقد اظهرت نتائج التحليل الفيزيوكيميائي للزيتين النتائج التالية: معامل الإنكسار 1.47001، 1.4975 الكثافة النوعية 0.8176، 0.8511 لكل من المحلب الأسود والأبيض على التوالي. الأحماض الدهنية الحرة 3.65%، 3.93%، قيمة الحامض 7.30، 7.86، قيمة البيروكسيد 4.43، 2.54، رقم التصبن 180.3، 184.23، والمواد غير القابلة للتصبن 0.66، 0.92.

الأحماض الدهنية التي وجدت بالزيتين كانت كما يلي: بالمتيك 4.5% استياريك 16% أوليك 48.3% لينوليك 31.6% في زيت المحلب الأسود بينما وجدت في زيت المحلب الأبيض على هذا النحو بالمتيك 5.7% أوليك 46% لينوليك 48%.

تحليل الأحماض الأمينية اظهر أن بروتين البذرتين يختلفان في كمية الأحماض الأمينية الأساسية والتي تراوحت اعلى نسبة لها بين 43.5% في المحلب الأسود و 34.8% في الأبيض.

نتائج التحليل الإحصائي للتقييم الحسى للزيتين اظهرت فروق معنوية ($P \leq 0.05$) أن زيت المحلب الأبيض مقبول أكثر من الأسود.