

## References

- [1] Harzallah, O. & Drean, J.-Y. 2011. Macro and micro characterization of biopolymers: Case of cotton fibre. *Biotechnology of biopolymers*, 193.
- [2] Xu, B., Yao, X., Bel, P., Hequet, E. F. & Wyatt, B. 2009. High volume measurements of cotton maturity by a customized microscopic system. *Textile Research Journal*, 79, 937-946.
- [3] Chee, P. W. & Campbell, B. T. 2009. Bridging classical and molecular genetics of cotton fiber quality and development. *Genetics and Genomics of Cotton*. Springer, pp 283-311
- [4] Expert Engineers,(2012)'Cotton Fiber Physical and Chemical Properties of Cotton', *Textile Engineering and Fashion Design Blog* ,23 June. Available at <http://textilefashionstudy.com/cotton-fiber-physical-and-chemical-properties-of-cotton/> (Accessed 30 October 2013)
- [5] Gordon, S., Horne, S. & Vander Sluijs, M. 2009. Ginning and Fibre Quality Series: Moisture in Cotton-the Fundamentals. *Australian Cottongrower, The*, 30, 32.
- [6] Bange, M. 2007. Effects of climate change on cotton growth and development. *The Australian Cotton Grower*, 28, 41-45.
- [7] Ton, P. 2011. Cotton and Climate Change: Impacts and Options to mitigate and adapt.
- [8] Anthony, W., Van Dorn, D., Edwards, E., PEARSON, T., Malloum, I., Kechagia, U., Ozdemir, I. & Nyoni, D. 2001. Impact of ginning on fiber quality: The best ginning practices. *Report Of An Expert Panel On Ginning Methods, International Cotton Advisory Committee*.

- [9] Montalvo JR, J. G. 2005. Relationships between micronaire, fineness, and maturity. Part I. Fundamentals. *The Journal of Cotton Science*, 9, 81-88.
- [10] Montalvo JR, J. & Hoven, T. 2005. Relationship between micronaire, fineness and maturity. Part II. Experimental. *The J Cott Sci*, 9, 89-96.
- [11] Karademir, E., Karademir, C., Ekininci, R. & Gencer, O. 2010. Relationship between yield, fiber length and other fiber-related traits in advanced cotton strains. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 38, 111-116.
- [12] Chen, B.&Zhao,L.1991.Multiple correlation analysis of yield, fibre quality and plant characteristics in upland cotton. *PKV Res. J*, 23, 21-23.
- [13] Khan,M.,Sadaqat, H.&Tariq,M. 1991. Path coefficient analysis in cotton (*Gossypium hirsutum*). *Journal of Agricultural Research*, 29.
- [14] Zhar, F. M., Naveed, M. & Ali, A. 2004. Correlation analysis of seed cotton yield with fiber characteristics in *Gossypium hirsutum* L. *International Journal of Agriculture and Biology*, 4, 656-658.
- [15] Survey of fiber testing equipment. 2006. International Cotton Advisory Committee, Washington.
- [16] Ikiz, Y., Rust, J., Jasper, W. & Trussell, H. 2001. Fiber length measurement by image processing. *Textile Research Journal*, 71, 905-910.
- [17] Nair, A., Nachane, R. & Patwardhan, B. 2009. Comparative study of different test methods used for the measurement of physical properties of cotton. *Indian journal of fibre & textile research*, 34, 352.

- [18] Paudel, D. R. 2012. *Evaluating the potential of new testing methods for cotton (Gossypium hirsutum L.) breeding*. Texas Tech University.
- [19] USDA /Uster Technology. 2006. Quality characteristics used for cotton classification, Uster® HVI 1000.
- [20] Kretzschmar, S. D. & Furter, R. 2008. USTER® AFIS PRO 2.
- [21] Bushara, M. O. Year. Decomposing Total Factor Productivity Change of Cotton Cultivars (Barakat-90 and Barac (67) B) in the Gezira Scheme (1991–2007) Sudan. *In:* 2010 AAAE Third Conference/AEASA 48th Conference, September 19-23, 2010, Cape Town, South Africa, 2010. African Association of Agricultural Economists (AAAE) & Agricultural Economics Association of South Africa (AEASA).
- [22] Ali, B. E. B. 2004. *Sudanese Cotton Specifications (Barakat-90 and Acala Varieties)*. MSc Thesis, Sudan University of Science and Technology.