

DEDICATION

To my father, mother and my husband

To my sisters and brothers

To my friends and colleagues

With my respect

EZDHAR

ACKNOWLEDGEMENTS

First of all, praise to Allah for giving me health and patience to complete this work. Also, I would like to express my deep gratitude and sincere thanks to Dr Atif Elsadig Idris for his guidance, supervision, and great effort he has paid during this study.

Thank are extended to Prof. Yassin Dagash and My thanks were expressed to all Doctor's in Department of Agronomy Dr. Samia Osman, Dr. Ahmmed Ali, Dr. Sami Ali and all the staff in the section

Special words of thanks to University of Khartoum. And section of crop production, Sudan University of Science and Technologyd My thanks are extended to my friends and colleagues.

I would like to express my deep gratitude and sincere thanks to my father, mother and my husband. Contributed in the work.

TABLE OF CONTENTS

	Page No.
Dedication	i
Acknowledgement	ii
Table of Contents	iii
List of Tables	v
List of Figures	vi
Abstract	vii
Arabic Abstract	viii
CHAPTER ONE	
1. Introduction	1
CHAPTER TWO	
2. Literature Review	5
2.1 Rhodes Grass	5
2.1.1 General Bakground	5
2.1.2 Varieties	6
2.1.3 Plant Description	7
2.1.4 Establishment and adaptation	7
2.1.5 Productivity	8
2.1.6 Environmental requirements	8
2.2 Clitoria	9
2.2.1 Characteristics and uses of clitoria	9
2.2.2 plant description	10
2.2.3 Climatic requirement	11

2.3 Intercropping	13
2.3.1 Definition and benefits	13
2.3.2 Methods of intercropping	14
2.3.3 Importance of inter cropping	14
2.3.4 Intercropping grasses with legumes	15
2.4.5 Effect of legumes on plant growth yield of grasses	16

CHAPTER THREE

3. Materials and Methods	18
3.1 Experimental site description	18
3.2 The plant material	18
3.3 Land preparation, design and cultural practices	18
3.4 Treatments application and cutting	19
3.5 Data collection	19
3.5.1 Plant height (cm)	19
3.5.2 Number of leaves/plant	19
3.5.3 Leaf area (cm ²)	20
3.5.4 Leaf to stem ratio	20
3.5.5 Forage yield (t/ha)	20
3.5.6 Forage dry yield (t/ha)	20
3.5.7 Crude protein (%)	20
3.5.8 Statistical analysis	21

CHAPTER FOUR

4. Results	22
4.1 Plant height (cm)	22
4.2 Number of leaves/plant	22
4.3 Leaf area (cm ²)	22

4.4 Leaf to stem ratio	22
4.5 Forage fresh weight	32
4.6 Forage dry weight	32
4.7 Determination of crude protein (%)	32

CHAPTER FIVE

5. Discussion	39
---------------	----

CHAPTER SIX

6. Summary and Conclusions	41
6.1 Summary	41
Conclusions	42
REFERENCES	43
APPENDICES	50

LIST OF TABLES

Table	Page No.
3.1: The form of analysis of variance with a randomized complete block design for Rhodes grass some growth and quality characters as intercropping with Clitoria	21
3.2: The form of analysis of variance with a randomized complete block design for yield of Rhodes grass and Clitoria (pure stand and mixed)	21
4.1: The mean squares from individual analysis of variance for different characters of Rhodes grass as intercropping with Clitoria	23
4.2: The mean of plant height (cm) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	24
4.3: The mean of No. of leaves for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	26
4.4: The mean of leaf area (cm ²) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	28
4.5: The mean of leaf to stem ratio for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	30
4.6: The mean of fresh weight (t/ha) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	33
4.7: The mean of forage dry weight (t/ha) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	35
4.8: The mean of crude protein % for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	37

LIST OF FIGURES

Figure	Page No.
4.1: The mean of plant height (cm) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	24
4.2: The mean of No. of leaves for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	26
4.3: The mean of leaf area (cm ²) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	28
4.4: The mean of leaf to stem ratio for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	30
4.5: The mean of fresh weight (t/ha) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	33
4.6: The mean of forage dry weight (t/ha) for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	35
4.7: The mean of crude protein % for Rhodes grass plants evaluation in different ratios of intercropping between it and Clitoria	37

ABSTRACT

The experiment was conducted at the Experimental farm, college of Agricultural studies, Sudan University of Science and Technology, Shambat. Five ratios of intercropping between Rhodes grass and Clitoria were used in this study.

The experiment was laid out in a randomized complete block design (RCBD) with three replications. Seven different characters were measured for consequence cuts. These characters were plant height (cm), leaf area (cm^2), number of leaves/ plant, leaf to stem ratio, fresh forage yield t/ha, dry forage yield t/ha and crude protein for the third cut. The analysis of variance revealed non-significant difference between the four studied ratios for the three cuts for all growth, quality and yield (fresh and dry) except the dry forage yield of the second cut, it was significant ($P \leq 0.05$). For fresh and dry forage yield in all the three cuts for five treatments, the range of the forage yield was 46.83 to 62.66 t/ha for fresh yield and 6.11 to 7.3 t/ha for dry yield. The range of crude protein was 12.25 to 17.50 for the third cut.

الخلاصة

تم اجراء هذه التجربة بالمزرعة التجريبية بكلية الدراسات الزراعية، جامعة السودان للعلوم والتكنولوجيا (شمبات). تم إستخدام خمس نسب من الزراعة المختلطة بين حشيشة الرودس و علف الكلايتوريا تم إجراء التجربة بإستخدام تصميم القطاعات الكاملة العشوائية بثلاثة مكررات تم قياس سبعة صفات مختلفة لثلاث قطعات متتابعة والصفات هي طول النبات/سم، مساحة الورقة/سم²، عدد الأوراق في النبات،نسبة الأوراق إلى الساق، الإنتاجية للعلف الاخضر بالطن/هكتار، الإنتاجية الجافة للعلف بالطن/هكتار ونسبة البروتين الخام للقطعة الثالثة. أظهر تحليل التباين عدم وجود فروقات معنوية للنسب الخمسة من الزراعة المختلطة لكل صفات النمو والنوعيه والانتاجيه للعلف الاخضر والجاف للثلاثة قطعات في ماعدا القطعه الثانية للانتاجيه العلف الجاف فقد ظهرت بها فروقات معنويه ($P \leq 0.05$) كان المدى للمجموع الكلى للانتاجية العلف من ٤٦.٨٣ - ٦٢.٦٦ بالطن هكتار للعلف الاخضر ومن ٦.١١-٧.٣ بالطن هكتار للعلف الجاف. وكان المدى للبروتين الخام في القطعة الثالثة من ١٢.٢٥-١٧.٥٠ .