

Table o contents

الاستهلال.....	III
Dedication.....	IV
Acknowledgment.....	V
Abstract.....	VI
ملخص الدراسة.....	VII
CHAPTER ONE	
Introduction.....	1
CHAPTER TWO	
Literature review	
2.1 Feed additives.....	3
2.2 Antibiotics.....	4
2.3 Phytobiotic.....	5
2.4 Garlic (Allium Sativum).....	8
2.4.1 Scientific classification.....	8
2.4.2 Botanical description.....	8
2.4.3 Constituents of bulb (clove).....	9
2.4.4 Uses.....	10
2.4.4.1 Food use	10
2.4.4.2 Medicinal use.....	10
2.4.5 Pharmacological actions.....	10
2.4.5.1 Anti-atherosclerotic and cholesterol and lipid-lowering effects.....	10
2.4.5.2 Antimicrobial effects.....	11
2.4.5.3 Antioxidant effect.....	13
2.4.5.4 Immunomodulatory activity.....	13
2.5 Ginger (Zingiber officinale).....	14
2.5.1 Classification of ginger.....	14
2.5.2 Botanical description.....	14
2.5.3 Constituents.....	14
2.5.4 Uses.....	15
2.5.4.1 Food use.....	15
2.5.4.2 Herbal.....	16
2.5.5 Pharmacological actions.....	16
2.5.5.1 Anti-atherosclerotic and antioxidant activity.....	16
2.5.5.2 Antioxidant activity.....	17
2.5.5.3 Anti-inflammatory activity.....	18
2.5.5.4 Antimicrobial activity.....	19
2.6 Effect of garlic and ginger their mixture on broiler performance.....	19

CHAPTER THREE MATERIAL AND METHODS

3.1 Experimental chicks.....	23
3.2 housing.....	23
3.3 Experimental rations.....	24
3.4 Data collected.....	24
3.4.1 performance data.....	24
3.4.2 Slaughtering procedure.....	24
3.4.3 The taste panel.....	25
3.5 Experimental design and statistical data analysis.....	25

CHAPTER FOUR Results

4.1 response o broiler chicks to diet containing mixture of garlic and ginger essential oils.....	28
4.1.1 Performance.....	28
4.1.2 Carcass dressing percentage.....	29
4.1.3 Panel test (subjective meat attributes).....	29
4.1.4 Economical appraisal.....	29

CHAPTER FIVE

5 Discussion.....	33
5.1 Conclusion.....	36
5.2 Recommendation.....	36
References.....	38
Appendix1.....	55
Appendix2.....	56
Appendix3.....	57

Tables list

Table1: The ingredients percent composition of experimental diets.....	26
Table2: Calculated chemical analysis of experimental diets:	27
Table3 The Effects of various levels of the dietary mixture of garlic and ginger essential oils on the performance and carcass dressing percentage of broiler chicks for 6 weeks.....	30
Table 4: the effect of different dietary amount of mixture the garlic and ginger essential oils on percentage of subjective values of broiler chicks for 6 weeks.....	31
Table 5: The total cost, revenue and net profit of broiler chicks fed on different levels of garlic and ginger essential oils.....	32

الآية

بسم الله الرحمن الرحيم

قَالُوا تَعَلَّمَ إِنَّا فِي الْأَوْرَاقِ نَحْنُ نَحْنُ يَطْبِرُ بِرْجَنَاحَ يُهْ
ذَمَالُكُمْ مَا فَرَّطْنَا فِي الْكِتَابِ مِنْ شَيْءٍ ثُمَّ إِلَى رَبِّهِمْ
يُحْشَرُونَ (38) ((

صدق الله العظيم

سورة الأنعام الآية (38)

Dedication

To the fountain of patience and optimism and hope my dear mother

To my big heart - dear father

To who support me to I be better in my life my sisters and brother

To the people who paved our way of science and knowledge

All our teachers distinguished

To whom I enjoyed with them the best and beautiful moments my friends

I guide this research

Acknowledgment

Firstly and lastly thanks to **ALLAH** who gave me persistence, and patience to complete this work. No words can adequately express my deep gratitude to my supervisor **Prof.Dr.Kamal Abdelbagi** for generously providing and for patience, constant support, advices and insight was invaluable to me. He is always available not only for consultation but also to solve any difficulties. Then I wish to express grateful thanks to administration of **Sudan University of Science and Technology**, college of **Agricultural Studies** for allowing me to conduct my research and providing any assistance requested. Gratitude also extended to **all people and friends** that assisted me in this research. For their excitement and willingness to provide feedback made the completion of this research and enjoyable experience.

ABSTRACT

This experiment was conducted to evaluate the response of broiler chicks fed on graded levels of mixture garlic and ginger essential oils as natural growth promoter alternative to antibiotic. Experimental parameters covered growth performance, carcass dressing percentage, subjective meat quality and economical appraisal. The experimental design used the complete randomized design (CRD). A total number of (84), 7days-old, 130gm initial weight unsexed *AborAcer* strain of broiler chicks were randomly divided into four experimental groups with three replicates, each of seven chicks. The first group (A) fed on basal diet without feed additives (control group), the other groups B,C and D were fed basal diets supplemented with different mixture of essential oils at levels of 200gm/ton(100gm ginger+100gm garlic),400gm/ton(200gm ginger+200gm garlic)and 600gm/ton(300gm ginger +300gm garlic). The basal diet was formulated to meet the nutrients requirements of broiler chicks according to (NRC, 1994). Experimental diets were fed for 6weeks.The results indicated that there were no significant ($p>0.05$) differences among all treatment groups in the values of body weight gain, feed intake, feed conversion ratio, carcass dressing percentages and subjective meat quality attributes. No mortality was recorded throughout the experimental period. The economical evaluation showed that levels of dietary mixture garlic and ginger essential oils were economical feasible, but the values of profitability ratio(1.24) of group D (600gm/ton at mixture garlic and ginger essential oils)was the highest of the tested groups.

ملخص الدراسة

أجريت هذه الدراسة لتقييم مدى استجابة كتاكيت الدجاج اللحم المغذاة على مستويات متدرجة من خليط زيت الثوم والجزبيل كمحفز طبيعي للنمو بدلاً للمضادات الحيوية . شملت قياس التجربة الأداء الإنتاجي، نسبة تصافي الذبيحة، الصفات الانطباعية النوعية للحم والتقييم الاقتصادي. صممت هذه التجربة بإستخدام النظام العشوائي الكامل . تم استخدام 84 كتكوت عمر 7 أيام بوزن ابتدائي 130 جم غير مجنسة من سلالة الابرايكر ، تم تقسيمها عشوائياً إلى أربعة مجموعات تجريبية ، كل مجموعة ضمت ثلاثة مكررات ، بكل مكرر سبعة كتاكيت ، تمت تغذية المجموعة الأولى (A) على عليقه أساسية بدون أي إضافة (عليقه قياسية) ، أما المجموعات الأخرى D ، C ، B فقد تمت تغذيتها على العليقه الأساسية مضافة إليها خليط زيت الثوم والزنجبيل بنسبة 200 ، 400 ، 600 جم اطن على التوالي . تم تكوين العليقة القياسية لتقابل الاحتياجات الغذائية للدجاج اللحم الصادر من (NRC 1994) تمت التغذية على العلاقة التجريبية لمدة 6 أسابيع اثبتت النتائج المتحصل عليها بأنه لا توجد فروقات معنوية ($P > 0.05$) بين كل المجموعات التجريبية في قيم وزن الجسم المكتسب ، العليقة المستهلكة ، معدل التحويل الغذائي ، نسبة التصافي الذبيحة وصفات اللحم الانطباعية النوعية للدجاج اللحم. لم تسجل اي حالات للتفوق خلال فترة التجربة . اظهر التقييم الاقتصادي بإن كل المستويات الغذائية لخليط زيت الثوم والجزبيل كانت مجده اقتصاديا ولكن قيمة الربحية النسبية (1.24) في المجموعة D (600 جم اطن خليط زيت الثوم والجزبيل) كانت الاعلى بين المجموعات المختبرة .