

## References

- Albert, A., Rubbo, S., Goldacre, R., Darcy and Stove, J., (1945). the Influence of Chemical Constitution on Antibacterial Activity. Part II: A General Survey of the Acridine Series *J. Exp. Pathol.*, 26, 160.
- Arcamone, F.; et al., (1987), Synthesis and antitumour activity of new daunorubicin and adriamycin analogues. *Experientia*, 34, 1255.
- Bashir,S.J. and Maibach,H.I.,(2000),Quantative Structure analysis relationships in the prediction of skin sensitization potential.Biochem modulation skin React.,pp.61-64.
- Bate-Smith, E. C.; Westall, R. G., (1950), Some Physical Properties of Aqueous Solutiow of Tetramethylammonium Bromide and Tetramethylammonium
- Bell, P. H. Roblin, and R. O., (1942), Studies in Chemotherapy. VII. A Theory of the Relation of Structure to Activity of Sulfanilamide Type Compounds, *J. Am. Chem. Soc.*, 64, 2905.
- Broto, P., Moreau, G., Vandycke, C., (1984), Molecular structures perception, auto correlation descriptor and SAR studies—system of atomic contributions for the calculation of the normal octanol water partition coefficients. *Eur. J. Med. Chem.* 19, 71–78.
- Cortes C. and Vapnik V., (1995), Support-Vector Networks. *Mach. Learn.*, 20:273–297.
- Crum-Brown, A. Fraser, T.R. (1868) On the connection between chemical constitution and physiological action. Part 1. On the physiological action of the ammonium bases, derived from Strychia, Brucia, Thebaia, Codeia, Morphia and Nicotia, *Trans. Roy. Soc. Edinburgh* 25, 151- 203.
- Davies MN, Hattotuwagama CK, Moss DS, Drew MGB and Flower DR. (2006) Statistical deconvolution of enthalpic energetic contributions to MHC-peptide binding affinity. *BMC Struct Biol.*, 6:5, 1–5, 13.

Dearden, J. C.; Mays, P. K. J., (1985), Can molecular connectivity serve as a steric parameter in quantitative structure activity relationships. *Pharm. Pharmacol.*, 61, 203.

Debolt, S. E., and Kollman, P. A., (1995), Investigation of Structure, Dynamics, and Solvation in 1-Octanol and Its Water-Saturated Solution: Molecular Dynamics and Free-Energy Perturbation Studies. *J. Am. Chem. Soc.*, 117, 5316.

Dewar, M.J.S. and Walter Thiel.,(1977) , Ground States of Molecules, 38. The MNDO Method. Approximations and Parameters. *J. Am. Chem. Soc.*, 99:4899-4907.

Dewar, M.J.S.,McKee M.L.,(1977), Ground states of molecules. 41. MNDO results for molecules containing boron. *J. American chem.,Soc.*, 99, 5231.

Dewar, M.J.S., Zoebisch, E. G., Healy, E.F., Stewart, J. J. P. , (1985), "AM1: A New General Purpose Burckhardt, G. N., Ford, W. G. K., and Singelton, E., (1936) The hydrolysis of arylsulphuric acids. Part I, *J. Chem. Soc.*, 17, 78.

Donald J. A., (2003) Burger's Medicinal Chemistry and Drug Discovery. Sixth Edition, Volume 1, p 3. John Wiley& Sons, Inc.

Du H., Wang J., Hu Z. and Yao X., (2008) Quantitative Structure-Retention relationship study of the constituents of saffron aroma in SPME-GC-MS based on the projection pursuit regression method. *Talanta*,77:360–365.

Du HY., Watzl J., Wang J., Zhang XY., Yaol XJ. and Hu ZD., (2008), Prediction of retention indices of drugs based on immobilized artificial membrane chromatography using Projection Pursuit Regression and Local Lazy Regression. *J. Sep. Sci.*, 31:2325–2333.

Ferguson, J., (1939), The Use of Chemical Potentials as Indices of Toxicity Proc. R. Soc. London Ser. B, 127,387.

Friedman JH. and Stuetzle W., (1981), Projection Pursuit Regression. *J. Am. Stat. Assoc.*, 76:817–823.

Gesson J.P., Jacquesy J.C., Mondon M., (1980), The Synthesis of ( $\pm$ )-11-Deoxydaunomycinone via Regioselective Tandem Diels-Alder Reactions *Tetrahedron Lett.* 21, 3351.

Gharagheizi F, Tirandazi Band Barzin R. (2009), Estimation of aniline point temperature of pure hydrocarbons: A quantitative structure-property relationship approach. *Ind. Eng. Chem. Res.*;48:1678–1682.

Gombar,V.K. and Enslien K., (1996), Assessement of n-octanol/water partition coefficient., *J.chem.inf.comput.sci.*,vol 36 (6)pp.1127-1134.

Guha R., Dutta D., Jurs PC. and Chen T., (2006), Local lazy regression: Making use of the neighborhood to improve QSAR predictions. *J. Chem. Inf. Model.*, 46:1836–1847.

Gupta, S.P., (1991), QSAR Studies on Local Anesthetics, *Chem. Rev.*, 91, 1109-1119.

Hammett, L.P. (1935) Some Relations Between Reaction Rates and Equilibrium Constants. *Chem. Rev.*, 17, 125-136.

Hammett, L. P.,( 1970) Physical Organic Chemistry. Reaction Rates, Equilibria and Mechanism, 2nd ed., McGraw-Hill, New York.

Hammett, L. P., (1966) Physical Organic Chemistry in Retrospect, *J. Chem. Ed.*, 43, 464.

Hansch, C., Kurup, A., Garg, R., and Gao, H., (2001), Chem-bioinformatics and QSAR: a review of QSAR lacking positive hydrophobic terms, *Chem. Rev.*, 101, 619-72.

Hansch, C.; Fujita, T., (1964)  $\rho$ - $\sigma$ - $\pi$  Analysis. A Method for the Correlation of Biological Activity and Chemical Structure. *J. Am. Chem. Soc.*, 86, 1616-1626.

Hansch C, Maloney PP, Fujita T, Muir RM., (1962) Correlation of biological activity of phenoxyacetic acids with hammett substituent constants and partition coefficients. *Nature*, 194: 178-180.

Hansch, C. (1969) Quantitative approach to biochemical structure–activity relationships. *Acc. Chem. Res.*, 2, 232–239.

Hansch, C., and Fujita, T., (1964),  $\rho-\sigma-\pi$  analysis a method for the correlation of biological Activity and Chemical Structure. *J. Am. Chem. Soc.*, 86, 1616.

Hansch, C; Klein, T E. (1986), Molecular graphics and QSAR in the study of enzyme-ligand interactions. On the definition of bioreceptors, *Acc. Chem. Res.* 19, 392.

Hastie, T., Tibshirani, R. and Friedman, J., (2009), The elements of statistical learning, 2nd edition. Springer.

Hildebrand, J. H., (1979), Is there a “hydrophobic effect”? *Proc. Natl. Acad. Sci. USA*, 76, 194.

Iodide Biochem. Biophys. Acta., 4, 427-440.

Israelachvili, J. N. and Wennerstrom, H., (1992), Entropic forces between amphiphilic surfaces in liquids, *J. Phys. Chem.*, 96, 520.

Iwamura, H., Nishimura, K., and Fujita, T., (1985), Quantitative structure-activity relationships of insecticides and plant growth regulators: comparative studies toward understanding the molecular mechanism of action, *Environ Health Perspect.* 61: 307–320.

Jung, M. E., Node, M., Pfluger, R., Lyster, M. A. and Lowe, J. A.,(1982), "Regiospecific Synthesis of Bicyclic 6-Alkoxy-2-Pyrone and Their Use in the Production of Tetracyclic Inter-mediates for 11-Desoxyanthracycline Synthesis," *J. Org. Chem.*, 47, 1150-1152.

Karelson, M.; Lobanov, V. S., Katritzky, A. R., (1996) Quantum-Chemical. Descriptors in QSAR/QSPR Studies. *Chem. Rev.*, 96, 1027.

Katritzky AR., Lobanov VS. and Karelson M., (2007) , Comprehensive Descriptors for Structural and Statistical Analysis (CODESSA) Ref. Man. Version 2.7.10.

Karitzky,A.R., Petrukhin.R. and Tatham,D., (2001), interpretation of quantitative structure property and activity relationship, *J. chem. inf. comput. sci.*, vol 41(3),pp. 679-685.

Krapcho A.P., Maresh M. J., Helgason A.L., Rosner K.E., Hacker M.P., Spinelli S., Menta E., Oliva A., (1993), The synthesis of 6,9-bis[(aminoalkyl) amino] substituted benzo[g] quinoxaline-, benzo[g]quinazoline- and benzo-[g]phthalazine-5,10-diones via regiospecific displacements, *J.Heterocyclic chem.* 30, 1597.

Kita Y., Kirahara M., Fujii Y., Okunaka R., Akai S., Maeda H., Tamura, Y., J. Chem. Soc. Chem. Commun., (1990), Synthetic anthracyclines: regiospecific total synthesis of a D-ring pyridine analogue of 11-deoxydaunomycin, 136.

Klopman, G.; Namboordiri, K.; Schochet, M., (1985), Simple Method of Computing the Partition Coefficient. *J. Comput. Chem.*, 6, 28-38.

Klopman G., Li Jy, Wang Sm, et al., (1994) Computer Automated Log P Calculations Based On An Extended Group-Contribution Approach *J. Chem. inf comp Sci* 34: (4) 752-781.

Kubinyi, H., QSAR: Hansch Analysis and Related Approaches, 43.

Kubinyi, H.,(1977), Quantitative structure-activity relations. 7. The bilinear model, a new model for nonlinear dependence of biological activity on hydrophobic character, *J. Med. Chem.*, 20 (5), 625–629.

Leo, A. J., (1991), Hydrophobic parameter: measurement and calculation. *Methods Enzymol.*, 202, 544–591.

Leo A, Hansch C, and Elkins D (1971). "Partition coefficients and their uses". *Chem Rev* 71 (6): 525–616.

Liew Chin Yee and Yap Chun Wei, (2012), Statistical Modelling of Molecular Descriptors in QSAR/QSPR. First Edition,Wiley-VCH Verlag GmbH & Co. KGaA., 31.

Lipnick R L., (1986) Charles Ernest Overton: narcosis studies and a contribution to general pharmacology. Trends. *Pharmacol Sci.*;7:161–164.

Lippold, B. C. and Adel, M. S., (1972), Arch. Pharm., 305, 417.

Livingstone, D. J., (2000), Infrared Spectra Information and Their Correlation with QSAR Descriptors. *J. Chem. Inf. Comput. Sci.*, 40, 195.

Liu H., Papa E., Walker JD. And Gramatica P., (2007), In silico screening of estrogen-like chemicals based on different nonlinear classification models. *J. Mol. Graph. Model.*, **26**:135–144.

Liu HX., Yao XJ., Zhang RS., Liu MC., Hu ZD. And Fan BT., (2005), Prediction of the tissue/blood partition coefficients of organic compounds based on the molecular structure using least-squares support vector machines. *J. Comput. Aided Mol. Des.*, **19**: 499–508.

Li X., Luan F., Si H., Hu Z. and Liu M., (2007), Prediction of retention times for a large set of pesticides or toxicants based on support vector machine and the heuristic method. *Toxicol. Lett.*, **175**:136–144

Lorentz, J., and Sorana-Daniela, B.,(2007), Results from the Use of Molecular Descriptors Family on Structure Property/Activity Relationships, *Int. J. Mol. Sci.*, **8**: 189-203.

Mandal AS., and Roy, K., 2009, Predictive QSAR modeling of HIV reverse transcriptase inhibitor TIBO derivatives. *Eur. J. Med. Chem.*, **44**:1509–1524.

Mannhold, R. and Dross, K., (1996), Quant. Struct.-Act. Relat., **15**: 403.

Maria das G. C., David L.N., Antônia T. A., Custódio D. S., Alcilene A. P., Ana C. B. O., (2002), Reverse Phase Thin Layer Chromatography of Amino alkanethiosulfuric Acids, Mercaptoalkanamines and Aminoalkyl Disulfides, *Int. J. Mol. Sci.*, **3**: 755-763.

Miller, A., (2002). Subset selection in regression, 2nd edition. Chapman & Hall/CRC.

Moore, W. J., (1972), Physical Chemistry, 4th. ed., (Prentice-Hall, Englewood Cliffs, N.J.,

Muxfeldt H., (1962). Syntheses in the Tetracycline SeriesAngew. *Chem. Int. Ed. Engl.* **1**: 372.

Nathan B., (2012). *Bioisosteres in Medicinal Chemistry*. Wiley-VCH, 237.

Nusselder, J. J. H. and Engberts, J. B. F. N., (1991), Langmuir, **7**, 2089.

Paul, K. A., and Michael, E., (2000), Synthetic routes to Cleistopholine and methylated analogues, ARKIVOC, 43-50.

Priolo N., Arribere CM., Caffini N., Barberis S., Vazquez RN. and Luco JM., (2001), Isolation and purification of cysteine peptidases from the latex of Araujia hortorum fruits - Study of their esterase activities using partial least-squares (PLS) modeling. *J. Mol. Catal. B-Enzym.*, **15**:177–189.

Quantum Mechanical Molecular Model", *J. Am. Chem. Soc.*, **107**, 3902.

Rekker R.F., (1977), The Hydrophobic Fragmental Constant, Pharmacochemistry Library, Vol. 1, Elsevier, Amsterdam.

Ren YY., Liu HX., Li SY., Yao XJ. And Liu MC., (2007), Prediction of binding affinities to beta(1) isoform of human thyroid hormone receptor by genetic algorithm and projection pursuit regression. *Bio org. Med. Chem. Lett.*, **17**: 2474–2482.

Rose, G. D., Geselowitz, A. R., Lesser, G. J., Lee, R. H. and Zehfus, M. H., (1985), Hydrophobicity of amino acid residues in globular proteins Science, 229, 834.

Rozeboom M.D., Tegmo-Larsson I.M., Houk T.N., (1981), Frontier Molecular Orbital Theory of Substituent Effects on Regioselectivity of Nucleophilic Additions and Cycloadditions to Benzoquinones and Naphthoquinones, *J. Org. Chem.* **46**, 2338

Stewart. J.J.P. (1996), Application of localized molecular orbitals to the solution of semiempirical self consistent field equations. *Int. J . Quant. Chem.*, **58**: 133–146.

Schneider, H. J., (1991). Mechanisms of Molecular Recognition: Investigation of Organic. Host-Guest Complexes, *Angew. Chem. Int. Ed. Engl.*, **30**, 1417.

Selassie, C. D., (2003), History of Quantitative Structure-Activity Relationships, Burger's Medicinal Chemistry and Drug Discovery Sixth Edition, Volume 1: Drug Discovery, p 3-9. John Wiley & Sons, Inc.

Sharon H. and Hans C. A., (1993), Ab initio and semiempirical methods for molecular dynamics simulations based on general Hartree-Fock theory, *J. Chem. Phys.* **99** (l): 523-532.

Showalter H., Johnson L., Hoftiezer M., Turner R., Werbel M., Leopold R., Shillis L., Jackson C. and Elslager F. (1987). Anthrapyrazole Anticancer Agents. Synthesis and Structure-Activity Relationships against Murine Leukemias. *J. Med. Chem.* **30**:121-131 .

Specht DF., (1991), A general regression neural network. *IEEE Trans. Neur. Netw.*, 2:568–576.

Stewart, James J. P. (1989), "Optimization of parameters for semiempirical methods I. Method". *J. Comput. Chem.* **10** (2): 209.

Szaleniec M., Tadeusiewicz R. and Witko M., (2008), How to select an optimal neural model of chemical reactivity . *Neurocomputing.*, **72**:241–256.

Taft, R.W., (1952) Linear Free Energy Relationships from Rates of Esterification and Hydrolysis of Aliphatic and Ortho-substituted Benzoate Esters. *J. Am. Chem. Soc.*, 74, 2729.

Taft RW. (1952), Polar and steric substituent constants for aliphatic and o-Benzoate groups from rates of esterification and hydrolysis of esters1. *J Am Chem Soc*, **74**: 3120-3128.

Taft R. W., and Newman, M. S., (1956), Ed., *Steric Effects in Organic Chemistry*, John Wiley and Sons, New York, , p. 556.

Tute, M. S. and Ramsden, C. A. (1990), Ed., *Comprehensive Medicinal Chemistry. The Rational Design, Mechanistic Study and Therapeutic Application of Chemical Compounds*, Vol. 4, Quantitative Drug Design, Pergamon, Elmsford, NY, 18.

Vanelle P., Rathelot P., Rémusat V., (2002), Synthesis of 3-Alkenyl-1-azaanthraquinones via Diels-Alder and Electron Transfer Reactions, *Molecules* 7, 917

- Vanyur R., Heberger K., Kovacs I. and Jakus J., (2002), Prediction of tumoricidal activity and accumulation of photosensitizers in photodynamic therapy using multiple linear regression and artificial neural networks. *J. Photochem. Photobiol.*, **75**:471–478.
- Wang T., Si HZ., Chen PP., Zhang KJ. And Yao XJ., (2008), QSAR models for the dermal penetration of polycyclic aromatic hydrocarbons based on Gene Expression Programming. *QSAR Comb. Sci.*, **27**:913–921.
- Wang WJ., Xu ZB., Lu WZ. And Zhang XY., (2003), Determination of the spread parameter in the Gaussian kernel for classification and regression Neurocomputing, **55**, PII S0925-2312(02)00632-X.
- Wong C.M., Popien D., Schenk R., Te Raa J., (1971), Synthetic Studies of Hydronaphthacenic Antibiotics. I. The Synthesis of 4-Demethoxy-7-O-methyl Daunomycinone. *Can. J. Chem.* **49**: 2712.
- Word H., (1966), Research Papers in Statistics. Wiley; New York, NY, USA.
- Xia BB., Liu KP., Gong ZG., Zheng B., Zhang XY. and Fan BT., (2009), Rapid toxicity prediction of organic chemicals to Chlorella vulgaris using quantitative structure-activity relationships methods. *Ecotoxicol. Environ. Saf.*, **72**:787–794
- Xia BB., Ma WP, Zheng B., Zhang XY. and Fan BT., (2008), Quantitative structure-activity relationship studies of a series of non-benzodiazepine structural ligands binding to benzodiazepine receptor. *Eur. J. Med. Chem.*, **43**:1489–1498.
- Yuan YN., Zhang RS., Hu RJ. And Ruan XF., (2008), Prediction of volatile components retention time in blackstrap molasses by least-squares support vector machine. *QSAR Comb. Sci.*, **27**:535–542.
- Žofka, J. (1970), Application of the Hartree-Fock method in nuclear theory, Czechoslovak Journal of Physics B, Vol. 20, 926-938.