

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ



*Sudan University of Science and Technology*  
*College of Graduate Studies*



## **Measurement of Renal Size in Sudanese Adult Using Computed Tomography**

قياس حجم الكلي في البالغين السودانيين باستخدام الاشعة

المقطعية

A thesis submitted for partial Fulfillment for the Requirement of M.Sc  
Degree in Diagnostic Radiologic technology

**By:**

**Amira Hamza ALbashir Osman**

**Supervisor:**

**Prof. Dr. Hussein Ahmed Hassan**

**2016**

## الآية

قال تعالى:

( قُلْ لَوْ كَانَ الْبَحْرُ مِدَادًا لِكَلِمَاتِ رَبِّي لَنَفِدَ الْبَحْرُ قَبْلَ أَنْ تَنفَدَ  
كَلِمَاتُ رَبِّي وَلَوْ جِئْنَا بِمِثْلِهِ مَدَدًا )

سورة الكهف الآية ( 109 )

## ***Dedication***

*To soul of my father*

*To My mother*

*To my sister*

*To my Husband and my Kids maysam*

*To my teachers*

*To all those who help me in preparation of the study*

## *Acknowledgement*

*My deep thanks to my supervisor Prof. Dr. Hussein Ahmed Hassan for his contact supervision, inexhaustible patience & unlimited help.*

*My thanks extend to staff of Ibn- Alhitham Diagnostic Center and Dar-Alelaj specialized Hospital. and also to my colleague Azza Alshareef Ahmed whose help me to complete this research.*

*My thanks to everyone who helped, supported and provided any type of help through this study.*

*Finally, special thanks to my family and friends who were of helpful during the whole study period.*

## ABSTRACT

This is descriptive study was conducted in three months during the period from September 2016 to December 2016 in Ibn- ALhitham Diagnostic Center and Dar-ALelaj specialized Hospital.

The aims of this study were to evaluate normal renal measurements and texture in Sudanese adults using Computed Tomography and correlated to age and gender.

This study carried out in a sample of 50 patients (25 males and 25 females) who underwent to Computed Tomography Urography examination for different reason.

The main results of this study were that mean and standard deviation of all sample of the tow kidneys, the lengths  $10.44 \pm 0.85$ ,  $10.57 \pm 0.91$ , widths  $4.55 \pm 0.6$ ,  $4.71 \pm 0.66$  And the cortex thickness was  $0.73 \pm 0.21$ ,  $0.74 \pm 0.22$  cm for right and left kidneys respectively, also the study showed that the texture of Right, Left kidneys  $39.20 \pm 5.08$ ,  $39.20 \pm 6.27$  Hounsfield unit. also mean and standard deviation of male's kidney lengths  $10.47 \pm 0.87$  ,widths  $4.58 \pm 0.66$  And the cortex thickness  $0.74 \pm 0.22$  cm and texture  $39.30 \pm 6.04$  Hounsfield unit, The female's kidney length  $10.54 \pm 0.89$  width  $4.68 \pm 0.65$ , the cortex thickness  $0.72 \pm 0.21$  cm and texture  $39.10 \pm 5.04$  Hounsfield unit.

The concluded of this study were that the kidneys measurements decreased with age, and the texture of kidneys increased with age .also the study showed there were no different between males and females kidneys measurements and texture.

The study recommended the future studies should be done with other modalities (magnetic resonance Imaging, positron emission tomography / computed tomography).

## ملخص البحث

هذه الدراسة الوصفية أجريت في ثلاثة أشهر من سبتمبر 2016 الي ديسمبر 2016 بمركز ابن الهيثم التشخيصي ومستشفى دار العلاج الخاص .

الهدف من هذه الدراسة هو تقييم قياسات الكلي الطبيعية ونسيج الكلي في البالغين السودانيين باستخدام التصوير المقطعي وربطها بالعمر والنوع.

أجريت هذه الدراسة علي عينه من 50 مريض (25 ذكور و25 اناث) أخضعوا لفحص الاشعة المقطعية الملونة للكلي لأسباب مختلفة.

كانت اهم نتائج هذه الدراسة ان المتوسط والانحراف المعياري لكل العينات بالنسبة للكليتين الطول  $10.57 \pm 0.91$ ,  $10.44 \pm 0.85$  العرض  $4.71 \pm 0.66$ ,  $4.55 \pm 0.64$  وسمك القشرة  $0.74 \pm 0.22$ ,  $0.73 \pm 0.21$  سم للكلية اليمين والشمال علي التوالي وايضا اشارة الدراسة ان نسيج الكلي اليمن والشمال  $39.20 \pm 6.27$ ,  $39.20 \pm 5.08$  معامل التوهين الخطي وايضا المتوسط والانحراف المعياري بالنسبه لطول كلي الذكور  $10.47 \pm 0.87$  العرض  $4.58 \pm 0.66$  وان سمك القشرة  $0.74 \pm 0.22$  سم ونسيج الكليه  $39.30 \pm 6.04$  معامل التوهين الخطي وبالنسبه لطول كلي الاناث  $10.54 \pm 0.89$  العرض  $4.68 \pm 0.65$  وسمك القشرة  $0.72 \pm 0.21$  سم ونسيج الكلية  $39.10 \pm 5.04$  معامل التوهين الخطي.

خلصت الدراسة ان قياسات الكلي تقل مع العمر وان نسيج الكلي يزيد مع العمر. وايضا اظهرت الدراسة انه لا يوجد فرق بين الذكور والاناث في قياسات ونسيج الكلي.

توصي الدراسة الدراسات القادمة ان تجري باستخدام الرنين المغنطيسي والانبعاث الاشعاعي بالأشعة المقطعية.

## List Contents

No	Topic	Page
	الاية	I
	Dedication.	II
	Acknowledgement.	III
	Abstract.	IV
	ملخص البحث	V
	List of Contents.	VI
	List of Tables.	VI
	List of Figures.	VI
	List of abbreviations.	XII
<b>Chapter one</b>		
1_1	Introduction.	1
1_2	Problem of the Study.	2
1_3	Objectives.	3
1_4	Significant of the study.	3
1_5	Overview of the study.	3
<b>Chapter Tow</b>		
2_1	Anatomy.	4
2_2	Physiology.	11

2_3	Pathology.	13
2_4	CT scanners.	15
2_5	Previous studies.	18
<b>Chapter three</b>		
3_1	Materials.	19
3_1_1	Study sample.	19
3_1_2	Study area.	19
3_1_3	Place of study.	19
3_1_4	Duration of study.	19
3_1_5	Inclusion criteria.	19
3_1_6	Exclusion criteria.	19
3_1_3	Equipments.	19
3_2	Methods.	21
3_2_1	Technique used.	21
3_2_2	Image interpretation.	21
2_2_3	Method of kidneys Measurement.	21
<b>Chapter four</b>		
4_1	Results.	23



## Chapter five

5_1	Discussion.	34
5_2	Conclusion.	36
5_3	Recommendations.	37
	References.	38
	Appendices.	–

## List of Tables

NO	Title	Page
4.1	Show Study group gender distribution.	23
4.2	Study group Age distribution of Males.	24
4.3	Show Study group Age distribution of females.	25
4.4	Descriptive statistics of the Normal Sudanese Body Characteristics (Total Sample).	26
4.5	Descriptive Statistics Mean Standard deviation of Right Kidney Measurements for the total sample.	26
4.6	Descriptive Statistics Mean Standard deviation of LT Kidney Measurements for the total sample.	26
4.7	Descriptive Statistics Mean, Standard deviation of the variables for (25) Males.	27
4.8	Descriptive Statistics Mean, Standard deviation of the variables for (25) females.	27
4.9	Correlation between age and right kidney texture.	28
4.10	Correlation between age and right kidney length.	29
4.11	Correlation between age and right kidney width.	30
4.12	Correlation between age and left kidney texture.	31
4.13	Correlation between age and left kidney lengths.	32
4.14	Correlation between age and left kidney width.	33

## List of Figures

NO	Title	Page
2.1	Coronal view shows Location and External Anatomy of kidney.	5
2.2	A and B axial view show Location and External Anatomy of kidneys.	6
2.3	A and B show coronal image of internal anatomy.	7
2.4	Show anterior view Relations of the kidneys.	8
2.5	Blood Supply of Kidneys.	10
3.1	Toshiba sensation 4 slices Ibn-ALhitham Diagnostic center.	20
3.2	PHILIPS sensation 4 slices Dar-ALelaj specialized Hospital.	20
4.1	Show Study group gender distribution.	23
4.2	Show age group distribution of males.	24
4.3	Show age group distribution of females.	25
4.4	Scatter plot diagram shows a linear relation relationship between the age and the texture of RT kidney.	28
4.5	Scatter plot diagram shows a linear relation between the age and the length of RT kidney.	29
4.6	Scatter plot diagram shows a linear relation between the age and the width of RT kidney.	30
4.7	Scatter plot diagram shows a linear relation between the age and the texture of LT kidney.	31
4.8	Scatter plot diagram shows a linear relation between the age and the length of LT kidney.	32
4.9	Scatter plot diagram shows a linear relation between the age	33

	and the width of LT kidney.	
1	Coronal CT image for female (27years) show site measurements of RT Kidney length, width and cortical thickness diameters.	–
2	Coronal CT image for Male (33years) show site measurements of RT, LT Kidney length, width and cortical thickness diameters.	–

## **List of abbreviations**

BMI	Body mass index.
Ca <sup>+</sup>	Calcium ion.
CT	Computer Tomography.
CTU	Computer tomography urography.
HU	Hounsfield Unit.
IV	Intra venous.
IVC	Inferior Vena cava.
KV	Kilo volt.
L3	Third lumbar vertebrae.
LT	Left.
MA	Mill Amber.
ml/sec	Mill/second.
MRI	Magnetic Resonance Imaging.
Na <sup>+</sup>	Sodium ion.
NaCl	Sodium chloride.
PET	Positron Emission Tomography.
PH	Plasma hydrogen ion.
R2	Liner Correlation Coefficient.
RT	Right.
T12	Thoracic spine number 12.
TBW	Total body water.
UPJ	Uretero Pelvic Junction.

# **Chapter One**

## **Introduction**

# Chapter one

## Introduction

### 1-1 Introduction to CT:

Computed tomography (CT) of the renal system is a diagnostic imaging test used to help detect normal anatomy and diseases of the kidneys. CT scanning is fast, painless, noninvasive and accurate. Spiral CT provides many advantages in evaluation of the kidneys. Spiral scanning essentially eliminates data misregistration, and allows for visualization of the entire kidney during peak contrast enhancement. The generated dataset is free from motion artifact, which improves multiplanar reconstructions, these factors are importance for the evaluation and characterization of small renal tumors .Increased sensitivity in the detection of subtle asymmetries of the cortical nephrogram is also obtained, which is useful in diagnosing disorders such as renal artery stenosis, renal vein thrombosis, acute pyelonephritis, and renal obstruction .The advantages of spiral scanning in the evaluation of both renal inflammatory diseases and renal masses (hopkinsmedicine.org 2016).

The kidneys are retroperitoneal bean-shaped organs that lie in the paravertebral gutters against the posterior abdominal wall. They lie at an oblique orientation, with the upper poles more medial and posterior than the lower poles. They are located on each side of the spine between T12 and L4 and are embedded in perirenal fat (Lorrie, 2007).

The right kidney is usually slightly lower due to displacement by the liver. Each kidney is composed of an outer cortex and an inner medulla. Each kidney has 7 to 14 minor calyces that merge into 2 or 3 major calyces. The major calyces join to form the renal pelvis, which is

the largest dilated portion of the collecting system and is continuous with the ureters. The fat-filled cavity surrounding the renal pelvis is called the renal sinus. Surrounding the kidneys and perirenal fat is another protective layer called the renal fascia. The medial indentation in the kidney is called the hilum; it allows the renal artery and vein and ureters to enter and exit the kidney the primary function of the urinary system is to filter blood, produce and excrete urine, and help maintain normal body physiology (Lorrie, 2007).

Typically, the kidneys are fully developed in the early twenties (about 23 to 25 years of age). There should be very little or no changes to the size of the kidneys, after this age. Kidney disease is an important factor that may affect kidney size. In fact, the size of a kidney and changes in its size can provide indications of renal problems. Polycystic disease and hydronephrosis (a condition where fluids accumulate inside the kidney) can cause distension (enlargement) of the kidneys. Other chronic renal conditions can affect kidney size and cause the kidneys to decrease in size. This is known as atrophy and usually occurs over a period of time (usually years). Chronic Glomerulonephritis and Hypertensive Nephrosclerosis are examples of conditions, which may cause atrophy (Hilaire, 2006-2016).

### **1-2 Problem of the study:**

The variant in normal renal measurements which may be due to ethnic variation and socioeconomic status and regions of zone, also some pathology may reduce or increase in kidneys size and changes in renal measurements .The size of kidney in Sudanese population compared to international index. Also The organ measurement usually affected by body characteristic, this characteristic might lead to wrong diagnosis therefore we need to compare this measurement to the body characteristic and hence we can have our own index.



### **1-3 Objectives:**

#### **1-3-1 general objective:**

To measure normal renal size in Sudanese adult using computed tomography.

#### **1-3-2 specific objectives:**

- To measure the kidneys size (length, width, thickness).
- To evaluate the texture of the normal kidneys.
- To compare kidneys measurements and texture with gender.
- To correlate kidneys (length, width) and texture with age.

### **1.4 Significant of the study:**

This study provides good information about Sudanese kidneys measurement and it used as guide line to proper Sudanese index.

### **1.5 Overview of study:**

Chapter One\_ Introduction and objectives of the study.

Chapter two\_ Literature review and background studies.

Chapter three\_ Materials and Methods.

Chapter four\_ The Results.

Chapter five\_ Discussion, Conclusion, Recommendations, References and Appendix.

# **Chapter Two**

## **Literature Review and Back ground studies**

## **Chapter two**

### **Literature Review and Background studies**


#### **2-1Anatomy:**

The kidneys are retroperitoneal bean-shaped organs that lie in the paravertebral gutters against the posterior abdominal wall. They lie at an oblique orientation, with the upper poles more medial and posterior than the lower poles. They are located on each side of the spine between T12 and L4 and are embedded in perirenal fat; the right kidney is usually slightly lower due to displacement by the liver. Each kidney is composed of an outer cortex and an inner medulla. The renal cortex comprises the outer one third of the renal tissue and has extensions between the renal pyramids of the medulla. The cortex contains the functional subunit of the kidney, the nephron, which consists of the glomerulus and convoluted tubules and is responsible for filtration of urine. (Lorrie, 2007).


The renal medulla consists of segments called renal pyramids that radiate from the renal sinus to the outer surface of the kidney. The striated-appearing pyramids contain the loops of Henle and collecting tubules and function as the beginning of the collecting system. Arising from the apices of the pyramids are the cup-shaped minor calyces. Each kidney has 7 to 14 minor calyces that merge into 2 or 3 major calyces. The major calyces join to form the renal pelvis, which is the largest dilated portion of the collecting system and is continuous with the ureters. The fat-filled cavity surrounding the renal pelvis is called the renal sinus. Surrounding the kidneys and perirenal fat is another protective layer called the renal fascia (Gerota's fascia). The renal fascia functions to anchor the kidneys to surrounding structures in an attempt to prevent

bumps and jolts to the body from injuring the kidneys. In addition, the renal fascia acts as a barrier, limiting the spread of infection that may arise from the kidneys. The medial indentation in the kidney is called the hilum; it allows the renal artery and vein and ureters to enter and exit the kidney (Lorrie, 2007).


The kidneys can be divided into five segments according to their vascular supply: apical, anterosuperior (upper anterior), anteroinferior (middle inferior), inferior, and posterior the segmental classification helps with surgical planning for partial nephrectomies (Lorrie, 2007).



**Figure 2.1 Coronal view shows Location and External Anatomy of kidneys.** (classes.midlandstech.edu).




**Figure: A**




**Figure: B**

**Figure: 2.2: A and B axial view show Location and External Anatomy of kidneys. (classes.midlandstech.edu)**



**Figure: A**



**Figure: B**

**Figure 2-3: A and B show coronal image of internal anatomy of kidney. (classes.midlandstech.edu)**

### **Relations of the right kidney:**

Superiorly and Anteriorly: the right suprarenal gland and the liver.


Anteriorly: The second part of the duodenum and the right colic flexure.

Posterior :the diaphragm, cost diaphragmatic recess of the pleura, the 12th rib and muscles of the posterior abdominal wall (Butler et al. 2007).

### **Relations of the left kidney:**

Anteriorly: The left suprarenal gland, the spleen, the stomach, the Pancreas, the left colic flexure, and loops of jejunum.

Posteriorly: as for the right kidney (Butler et al.2007).



**Figure: 2.4 Show anterior view Relations of the kidneys.**

([www.studyblue.com](http://www.studyblue.com)).

## **Blood supply of the kidneys:**


Renal arteries, veins and lymphatic drainage: The right and left renal arteries arise from the abdominal aorta, at approximately the level of the superior margin of L2, immediately caudal to the origin of the superior mesenteric artery, there is usually a single artery supplying each kidney, although there are many anatomical variants, with up to four renal arteries supplying each kidney. The renal artery divides in the renal hilum into three branches. Two branches run anteriorly, supplying the anterior upper pole and entire lower pole, and one runs posterior supplying the posterior upper pole and mid pole. Five or six veins arise within the kidney and join to form the renal vein, which runs anterior to the artery within the renal pelvis . The right renal vein has a short course, running directly into the IVC. The left renal vein runs anterior to the abdominal aorta and then drains into the IVC. Occasionally, the left renal vein runs posterior to the aorta, known as a retro-aortic renal vein. The left renal vein receives tributaries from the left inferior phrenic vein, the left gonadal and the left adrenal vein (Butler et al. 2007).

The lymphatic drainage of the kidneys follows the renal arteries to nodes situated at the origin of the renal arteries in the Para-aortic region (Butler et al. 2007).

## **Nerve supply:**

The sympathetic nerves supplying the kidney arise in the renal sympathetic plexus and run along the renal vessels. Afferent fibers, including pain fibers, travel with the sympathetic fibers through the splanchnic nerves and join the dorsal roots of the 11th and 12th thoracic and the 1st and 2nd lumbar levels (Butler et al. 2007).





**Figure: 2.5 Blood Supply of Kidney.** (classes.midlandstech.edu).

## 2.2 Physiology:

The primary function of the kidneys is to maintain a stable internal environment (homeostasis) for optimal cell and tissue metabolism. They do this by separating urea, mineral salts, toxins, and other waste products from the blood. They also do the job of conserving water, salts, and electrolytes. At least one kidney must function properly for life to be maintained. (Wiki/Human Physiology 2016).

Important roles of the kidneys are: **Regulation of plasma ionic composition.** Ions such as sodium, potassium, calcium, magnesium, chloride, bicarbonate, and phosphates are regulated by the amount that the kidney excretes. **Regulation of plasma osmolarity.** The kidneys regulate osmolarity because they have direct control over how many ions and how much water a person excretes. **Regulation of plasma volume.** Your kidneys are so important they even have an effect on your blood pressure. The kidneys control plasma volume by controlling how much water a person excretes. The plasma volume has a direct effect on the total blood volume, which has a direct effect on your blood pressure. Salt (NaCl) will cause osmosis to happen; the diffusion of water into the blood **Regulation of plasma hydrogen ion concentration (pH).** The kidneys partner up with the lungs and they together control the pH. The kidneys have a major role because they control the amount of bicarbonate excreted or held onto. The kidneys help maintain the blood Phmainly by excreting hydrogen ions and reabsorbing bicarbonate ions as needed **Removal of metabolic waste products and foreign substances from the plasma.** One of the most important things the kidneys excrete is nitrogenous waste. As the liver breaks down amino acids it also releases ammonia. The liver then quickly combines that ammonia with carbon dioxide, creating urea which is the primary nitrogenous end product of

metabolism in humans. The liver turns the ammonia into urea because it is much less toxic. We can also excrete some ammonia, creatinine and uric acid. **The creatinine** comes from the metabolic breakdown of creatine phosphate (a high-energy phosphate in muscles). **Uric acid** comes from the breakdown of nucleotides. Uric acid is insoluble and too much uric acid in the blood will build up and form crystals that can collect in the joints and cause gout. **Secretion of Hormones** The endocrine system has assistance from the kidney's when releasing hormones. Renin is released by the kidneys. **Renin** leads to the secretion of aldosterone which is released from the adrenal cortex. Aldosterone promotes the kidneys to reabsorb the sodium ( $\text{Na}^+$ ) ions. The kidneys also secrete erythropoietin when the blood doesn't have the capacity to carry oxygen. Erythropoietin stimulates red blood cell production. The Vitamin D from the skin is also activated with help from the kidneys. Calcium ( $\text{Ca}^+$ ) absorption from the digestive tract is promoted by vitamin D. (wiki/Human\_Physiology2016)

## **2.3 Pathology:**

### **Kidney stones:**

Kidney stone (renal calculus or nephrolith) is formed by combination of a high level of calcium with oxalate, phosphate, urea, uric acid, and cystine. Crystals and subsequently stones are formed in the urine and collected in calyces of the kidney or in the ureter the kidney stone varies in size from a grain of sand to the size of a golf ball and produces severe colicky pain while traveling down through the ureter from the kidney to the bladder. Common signs of kidney stones include nausea and vomiting, urinary frequency and urgency, and pain during urination (Chung et al.2012).

### **Polycystic kidney disease:**

Is a genetic disorder characterized by numerous cysts filled with fluid in the kidney; the cysts can slowly replace much of normal kidney tissues, reducing kidney function and leading to kidney failure. It is caused by a failure of the collecting tubules to join a calyx, which causes dilations of the loops of Henle, resulting in progressive renal dysfunction. This kidney disease has symptoms of high blood pressure, pain in the back and side, headaches, and blood in the urine. It may be treated by hemodialysis or peritoneal dialysis and kidney transplantation (Chung et al. 2012).

### **Hydronephrosis:**

Is a fluid-filled enlargement of the renal pelvis and calyces as a result of obstruction of the ureter. It is due to an obstruction of urine flow by kidney stones in the ureter, by compression on the ureter by abnormal blood vessels, or by the developing fetus at the pelvic brim. It has symptoms of nausea and vomiting, urinary tract infection, fever, dysuria (painful or difficult urination), urinary frequency, and urinary urgency (Chung et al. 2012).

**Pelvic kidney:**

Is an ectopic kidney that occurs when kidneys fail to ascend and thus remain in the pelvis. Two pelvic kidneys may fuse to form a solid lobed organ because of fusion of the renal anlagen, called a cake (rosette) kidney (Chung et al. 2012).

**Horseshoe kidney develops:** As a result of fusion of the lower poles of two kidneys and may obstruct the urinary tract by its impingement on the ureters (Chung et al. 2012).

## **2-4 CT scanners:**

Are complex, with many different components involved in the process of creating an image. Adding to the complexity, different CT manufacturers often modify the design of various components. From a broad perspective, all makes and models of CT scanners are similar in that they consist of a scanning gantry, x-ray generator, computer system, operator's console, and physician's viewing console. Although hard-copy filming has largely been replaced by workstation viewing and electronic archiving, most CT systems still include a laser printer for transferring CT images to film (Lois, 2011).

The three major components of a CT imaging system are the operating console, the computer, and the gantry. Each of these major components has several subsystems.

Computed tomography imaging systems can be equipped with two or three consoles. One console is used by the CT radiologic technologist to operate the imaging system. Another console may be available for a technologist to post process images to annotate patient data on the image (e.g., hospital identification, name, patient number, age, gender) and to provide identification for each image (e.g., number, technique, couch position). This second monitor also allows the operator to view the resulting image before transferring it to the physician's viewing console. A third console may be available for the physician to view the images and manipulate image contrast, size, and general visual appearance. This is in addition to several remote imaging stations (Lois, 2011).

The computer is a unique subsystem of the CT imaging system. Depending on the image format, as many as 250,000 equations must be solved simultaneously; thus, a large computing capacity is required. Many CT imaging systems use an array processor instead of a microprocessor for image reconstruction. The array processor does many

calculations simultaneously and hence is significantly faster than the microprocessor (Lois, 2011).

The gantry includes the x-ray tube, the detector array, the high-voltage generator, the patient support couch, and the mechanical support for each. These subsystems receive electronic commands from the operating console and transmit data to the computer for image production and post processing tasks (Lois, 2011).

X-ray tubes produce the x-ray photons that create the CT image. Their design is a modification of a standard rotating anode tube, such as the type used in angiography. Tungsten, with an atomic number of 74, is often used for the anode target material because it produces a higher-intensity x-ray beam. CT tubes often contain more than one size of focal spot; 0.5 and 1.0 mm are common sizes. Early CT scanners used recoiling system cables to rotate the gantry frame. Current systems use electromechanical devices called slip rings. Slip rings use a brush like. Apparatus to provide continuous electrical power and electronic communication across a rotating surface. They permit the gantry frame to rotate continuously, eliminating the need to straighten twisted system cables. (Lois, 2011).

As the x-ray beam passes through the patient it is attenuated to some degree. To create an x-ray image we must collect information regarding the degree to which each anatomic structure attenuated the beam. In CT, detectors used to collect the information. The detector array comprises detector elements situated in an arc or a ring, each of which measures the intensity of transmitted x-ray radiation along a beam projected from the x-ray source to that particular detector element. Detectors can be made from different substances, each with their own advantages and disadvantages (Lois, 2011).

All new scanners possess detectors of the solid-state crystal variety.

Detectors made from xenon gas have been manufactured but have largely become obsolete as their design prevents them from use in MDCT systems (Lois, 2011).

High-frequency generators are currently used in CT. They are small enough so that they can be located within the gantry. Generators produce high voltage and transmit it to the x-ray tube. CT generators produce high kV (generally 120–140 kV) to increase the intensity of the beam, which will increase the penetrating ability of the x-ray beam and thereby reduce patient dose. In addition, a higher kV setting will help to reduce the heat load on the x-ray tube by allowing a lower MA setting. Reducing the heat load on the x-ray tube will extend the life of the tube (Lois, 2011).

The patient lies on the table (or couch, as it is referred to by some Manufacturers and is moved within the gantry for scanning. The process of moving the table by a specified measure is most commonly called incrimination, but is also referred to as feed, step, or index. Helical CT table incrimination is quantified in millimeters per second because the table continues to move throughout the scan. The degree to which a table can move horizontally is called the scan able range, and will determine the extent a patient can be scanned without repositioning. The specifications of tables vary, but all have certain weight restrictions (Lois, 2011).

On most scanners, it is possible to place the patient either head first or feet first, supine or prone. Patient position within the gantry depends on the examination being performed (Lois, 2011).



## 2.5 Previous studies:

The study was done by (**Krairittichai U et al 2011**) by ultrasound. The average lengths of left and right kidney were  $10.24 \pm 0.70$  and  $10.09 \pm 0.68$  cm, respectively. Male kidney was bigger than the female kidney. Kidney length slightly increased until the age of 50 and become smaller at the age of 60 or more. Kidney length showed no correlation with age, body weight, body mass index and body surface area.

Another study was done by (**Abdullah et al 2014**) by MRI. The study showed that the kidneys length measured for normal Sudanese subjects were  $10.08 \pm 0.46$ ,  $10.67 \pm 0.47$  and the volumes were  $101.6 \pm 12.98$ ,  $104.0 \pm 12.99$  for right and left kidneys respectively, and it differed from other population. There were significant differences between males and females measurements and the correlation was significant between kidneys length and volume with BMI, TBW and subjects height.

The study was done by (**Maaji et al 2015**) by ultrasound the mean kidney length was  $11.3 \pm 8.8$  and  $11.6 \pm 9.8$  for right and left kidney, respectively. The mean height and weight was  $1.67 \pm 0.85$  and  $70.9 \pm 11.2$ , respectively. The mean kidney width was  $4.4 \pm 0.71$  and  $5.2 \pm 5.26$  for right and left kidney respectively. The mean renal thickness was ( $4.7 \pm 0.67$ ,  $4.5 \pm 0.68$ ) for right and left kidney respectively.

Another study was done by The (**Rathore et al 2016**) by computer tomography mean length, width, thickness and volume of the left kidney were  $11.02 \pm 1.13$  cm,  $5.21 \pm 0.75$  cm,  $4.65 \pm 0.84$  cm and  $138.22 \pm 29.81$  mL, respectively, and those for the right kidney were  $10.86 \pm 1.12$  cm,  $5.13 \pm 0.77$  cm,  $4.73 \pm 0.95$  cm and  $137.54 \pm 34.48$  ML, respectively.

# **Chapter Three**

## **Materials and Methods**

# **Chapter Three**

## **Materials and Methods**

### **3.1 Materials:**

#### **3.1.1 Study sample:**

This was descriptive study and the data collected and interpreted by radiologist.

#### **3.1.2 Study area:**

Sudanese population.

#### **3.1.3 Place of study:**

1-Ibn\_ALhitham Diagnostic Center

2-Dar\_ALelaj specialized Hospital.

#### **3.1.4 Duration of study:**

From 1/9/2016 to 1/12/2016.

#### **3.1.5 Inclusion criteria:**

Normal patients.

#### **3.1.6 Exclusion criteria:**

Patients with diseases of renal.

#### **3.1.7 Equipments:**

In the present study, CT machine were used.



**Figure3.1:** Toshiba sensation 4 slices Ibn-ALhitham Diagnostic center.



**Figure 3.2:** PHILIPS sensation 4 slices Dar-ALelaj specialized Hospital

## **3.2 Methods:**

### **3.2.1 Technique used:**

- ✓ The patient should be fasting at least 8 hours prior to exam.
- ✓ The patient position is in supine (feet first), with arms over head on CT table.
- ✓ The scout from diaphragmatic dome to the symphysis pubis.
- ✓ The slice thickness is used: 4-5 mm through the kidney.
- ✓ Breathe hold: suspended expiration.
- ✓ Land mark: xiphoid tip.
- ✓ I contrast: 2-4 ml/sec, 100-150 ml.
- ✓ Scout: AP, Pre contrast scans.
- ✓ The protocol used:  
CTU triphase: Arterial phase 30Sec, nephrogram 90 Sec,  
pyelogram phase 3-5 min.

### **3.2.2 Image interpretation:**

All CT image were studied for the study group sample to measure the RT\_LT Kidneys and evaluated the texture of kidneys and Statistical analyses were performed using excel software programmed and using the SPSS software.

### **3.2.3 Method of kidneys Measurement:**

#### **– Length measurement:**

On the coronal image from upper pole to lower pole (longest area), were measured in nephrogram phase.

#### **– Width measurement:**

On the coronal image from middle to lateral area (widest area), Three measurements were taken in nephrogram phase (First at upper pole, 2 at middle pole&3 at lower pole).

– **Cortical thickness measurement:**

On the coronal image from renal capsule to the base of renal pyramids is measured in nephrogram phase.

# **Chapter four**

## **Results**

# Chapter Four

## Results

The following tables and figures represent data obtained from randomly selected sample of patients (25males and 25females) and age range between (20\_75) years that underwent CT U for other indications without evidence of renal diseases.

Table 4.1: Show Study group gender distribution:

Gender	Frequency	Percentage
Male	25	50%
Female	25	50%
Total	50	100%





Figure 4.1: Show Study group gender distribution



**Table 4.2: Study group Age distribution of Males:**


Males		
Age classes	Frequency	Percentage %
20_30	5	20
31_40	7	28
41_50	6	24
51_60	2	8
61_70	4	16
71_80	1	4
Total	25	100%



**Figure 4.2: Show age group distribution of males**

**Table 4.3: Show Study group Age distribution of females:**

Females		
Age classes	Frequency	Percentage %
20_30	6	24
31_40	4	16
41_50	7	28
51_60	3	12
61_70	4	16
71_80	1	4
Total	25	100%



**Figure 4.3: Show age group distribution of females**

**Table 4.4: Descriptive statistics of the Age (Total Sample)**

<b>Descriptive Statistics</b>					
	<b>N</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>Std. Deviation</b>
Age	50	20.	75.	44.84	14.89742

**Table 4.5 Descriptive Statistics Mean Standard deviation of Right Kidney Measurements for the total sample:**

<b>Descriptive Statistics</b>					
	<b>N</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>Std. Deviation</b>
Length	50	8.60	12.00	10.4460	.85576
Width	50	3.50	5.90	4.5520	.64404
Cortex thickness	50	.27	1.67	.8002	.30186
Texture	50	30.00	55.00	39.200	5. 53652

**Table 4.6 Descriptive Statistics Mean Standard deviation of LT Kidney Measurements for the total sample:**

<b>Descriptive Statistics</b>					
	<b>N</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>Std. Deviation</b>
Length	50	8.70	12.00	10.5700	.91879
Width	50	3.70	6.00	4.7120	.66995
Cortex thickness	50	.35	1.75	.7558	.24367
texture	50	31.00	59.00	39.2000	5.65685

**Table 4.7 Descriptive Statistics Mean, Standard deviation of the variables for (25) Males:**

<b>Descriptive Statistics</b>					
	<b>N</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>Std. Deviation</b>
Age	50	20.00	75.00	44.8400	14.89742
Length	50	8.70	12.00	10.4720	.87902
Width	50	3.60	6.00	4.5840	.66712
Cortex thickness	50	.27	1.75	.8556	.30503
Texture	50	30.00	59.00	39.3000	6.04773


**Table 4.8 Descriptive Statistics Mean, Standard deviation of the variables for (25) females:**

<b>Descriptive Statistics</b>					
	<b>N</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Mean</b>	<b>Std. Deviation</b>
Age	50	21.00	75.00	44.8400	15.16893
Length	50	8.60	12.00	10.5440	.89947
Width	50	3.50	6.00	4.6800	.65341
Cortex thickness	50	.37	1.26	.7004	.21487
Texture	50	30.00	49.00	39.1000	5. 04773

**Table 4.9 Correlation between age and right kidney texture:**

Correlations			
		Age	texture
age	Pearson Correlation	1	.732**
	Sig. (2-tailed)		.000
	N	50	50

**\*\*.** Correlation is significant at the 0.01 level (2-tailed).




**Figure 4.4:** Scatter plot diagram shows a linear relationship between the age and the texture of RT kidney and the correlation is significant at P value 0.01, as the age increase the renal texture is also increased by factor of 0.266 starting from 27.3

**Table 4.10 Correlation between age and right kidney lengths:**

Correlations			
		age	length
age	Pearson Correlation	1	-.871**
	Sig. (2-tailed)		.000
	N	50	50


**\*\*.** Correlation is significant at the 0.01 level (2-tailed).



**Figure 4.5:** Scatter plot diagram shows a linear relation between the age and the length of RT kidney and the correlation is significant at p value 0.01, as the age increase the renal length is decreased by factor of -0.048 starting from 12.6.

**Table 4.11 Correlation between age and right kidney width :**

Correlations			
		age	width
age	Pearson Correlation	1	-.862**
	Sig. (2-tailed)		.000
	N	50	50
**. Correlation is significant at the 0.01 level (2-tailed).			




**Figure 4.6:** Scatter plot diagram shows a linear relation between the age and the width of RT kidney and the correlation is significant at p value 0.01 as the age increased the renal width is decreased by factor of -0.036 starting from 6.17.

**Table 4.12 Correlation between age and Left kidney texture:**

Correlations			
		age	texture
age	Pearson Correlation	1	.719**
	Sig. (2-tailed)		.000
	N	50	50

**\*\*.** Correlation is significant at the 0.01 level (2-tailed).




**Figure 4.7:** Scatter plot diagram shows a linear relation between the age and the texture of LT kidney and the correlation is significant at p value 0.01 as the age increased the renal texture is also increased by factor of 0.273 starting from 26.95



**Table 4.13 Correlation between age and left kidney length:**

Correlations			
		age	length
age	Pearson Correlation	1	-.900**
	Sig. (2-tailed)		.000
	N	50	50


**\*\*.** Correlation is significant at the 0.01 level (2-tailed).



**Figure 4.8:** Scatter plot diagram shows a linear relation between the age and the length of RT kidney and the correlation is significant at P value 0.01, as the age increased the renal length is decreased by factor of -0.054 starting from 12.98

**Table 4.14 Correlation between age and left kidney width:**

Correlations			
		age	width
age	Pearson Correlation	1	-.884**
	Sig. (2-tailed)		.000
	N	50	50
**. Correlation is significant at the 0.01 level (2-tailed).			



**Figure 4.9:** Scatter plot diagram shows a linear relation between the age and the width of LT kidney, and the relation show and the correlation is significant at p value 0.01, as the age increased the renal width is decreased by factor of -0.038 starting from 6.438

## **Chapter five**

# **Discussion, Conclusion and Recommendations**

# Chapter five

## Discussion, Conclusion and Recommendations

### 5.1 Discussion:

The aims of this study were to measure the kidneys size and evaluated the texture for Sudanese adult using CT.

The study showed mean and STD of total samples of the right kidney length measurements were  $10.44 \pm 0.85$ , width  $4.55 \pm 0.64$ , cortical thickness  $0.80 \pm 0.30$  cm, texture  $39.2 \pm 5.53$  HU. And the LT kidney length measurements were  $10.57 \pm 0.91$ , width  $4.71 \pm 0.66$ , cortical thickness  $0.76 \pm 0.24$  cm, texture  $39.2 \pm 5.65$  HU, in tables [4.5 and 4.6].

These measurements compare to study done by (Maaji et al 2015 ) found mean of RT,LT Kidneys length was  $11.3 \pm 8.8$  ,  $11.6 \pm 9.8$  cm ,which was decrease than this study by 0,86 , 1.03cm , and RT,LT Kidneys width was  $4.4 \pm 0.71$  and  $5.2 \pm 5.26$  cm ,the RT which was increase than this study by 0.15 cm , and the LT was decrease than this study by 0.49cm .

The study showed mean and STD of The males' kidney length  $10.47 \pm 0.87$ , width  $4.58 \pm 0.66$  and the cortex thickness  $0.86 \pm 0.30$  cm, texture  $39.30 \pm 6.04$  HU. And the females' kidney length  $10.54 \pm 0.89$ , width  $4.68 \pm 0.65$ , the cortex thickness  $0.70 \pm 0.21$  and texture  $39.10 \pm 5.04$  in tables [4.7 and 4.8].

This study showed there was no difference between males and females measurements.

The correlation between the age and the texture of Kidneys, the study showed that there was significant correlation at (P\_value 0.01) .and the texture of RT and LT kidney increased by factor of 0.266 \_0.273 with age as in figure [4.4\_4.7].

The correlation between the age and the length Diameter of RT, LT kidney, the study showed that there was significant correlation at (P\_value 0.01).and the length of RT, LT Kidneys decreased by factor of -0.048 \_-0.054 with age as in figure [4.5\_4.8].

The correlation between the age and the width Diameter of RT, LT kidneys, the study showed that there was correlation at (P\_value 0.01).And the width of RT, LT Kidney decrease by factor of -0.036 \_-0.038 with age as in figure [4.5\_4.8].

## 5.2 Conclusion:

- ✓ The mean and STD for measurement of the right kidney length measurements were  $10.44 \pm 0.85$ , width  $4.55 \pm 0.64$ , cortical thickness  $0.80 \pm 0.30$  cm, texture  $39.2 \pm 5.53$  HU. And the LT kidney length measurements were  $10.57 \pm 0.91$ , width  $4.71 \pm 0.66$ , cortical thickness  $0.76 \pm 0.24$  cm, and texture  $39.2 \pm 5.65$  HU.
- ✓ The study showed that no difference between males and females subject in the kidneys measurements and kidneys texture.
- ✓ The study showed that the texture of renal increased with age.
- ✓ The study showed that the RT, LT Kidney measurements (length, width) decreased with age and this indicate that the size of kidneys decreased with age.

### **5.3 Recommendations:**

- ✓ For further assessment another study should be done using large sample of patient for more accurate results.
- ✓ Future studies should be done with several body characteristic in correlation with kidneys measurements.
- ✓ Future studies should be done with laboratory investigation in correlation with kidneys measurements.
- ✓ Futures studies should be done with other modalities (MRI, PET/CT).

# References



## References

Hilaire [http://www.hopkinsmedicine.org/healthlibrary/test\\_procedures/urology/computed\\_tomography\\_ct\\_or\\_cat\\_scan\\_of\\_the\\_kidney\\_92,p07703](http://www.hopkinsmedicine.org/healthlibrary/test_procedures/urology/computed_tomography_ct_or_cat_scan_of_the_kidney_92,p07703).  
the page appeared on 3 Dec 2016 23:51:29 GMT.

<http://classes.midlandstech.edu/carterp/Courses/bio211/chap25/chap25.htm>.

[https://en.wikibooks.org/wiki/Human\\_Physiology](https://en.wikibooks.org/wiki/Human_Physiology). the page appeared on 1 Dec 2016 04:58:57 GMT.

<https://www.studyblue.com>.

Kidneychat.com Rodger St. Hilaire 2006\_2016.

Krairitichai U, Leehacharoenkul S, Dowreang J. J Med Assoc Thai. 2011 Mar; 94 Suppl 2:S23-8.

Kyung Won Chung, Harold M. Chung 2012, BRS\_Gross Anatomy \_7<sup>th</sup> Ed, Lippincott Williams & Wilkins, a Wolters Kluwer business, Philadelphia Baltimore, New York. London, pages 215\_217.

Lois E. Romans, 2011, computed tomography for technologists: a comprehensive text, Wolters Kluwer Health/Lippincott Williams & Wilkins, Philadelphia, pages 14\_20.

Lorrie L. Kelley and Connie M. Petersen, 2007, Sectional Anatomy for Imaging Professionals, 2nd edition, Elsevier Inc, Philadelphia, page 353\_367.

Mazin Babikir Abdullah, M,C & E, Establishment of Reference Values for Renal Length and Volume for Normal Adult Sudanese using M RI Disc Summation Method,Global Journal of Medical Research, Year 2014, Volume 14 Issue 2 Version 1.0 .


Paul Butler, Adam W. M. Mitchell and Harold Ellis 2007, Applied Radiological Anatomy for Medical Students.1th edition, Cambridge University Press, New York,page47\_49.

Ranjeet S. Rathore, Nisarg Mehta, Biju S. Pillai, Mohan P. Sam, Binu Upendran, and H. Krishnamoorthy, Variations in renal morphometry: A hospital-based Indian study,Indian Journal of Urology, Jan-Mar 2016, v. 32(1): 61–64. Advanced.


Sadisu Mohammed Maaji, Odunko Daniel, Bappa Adamu, Sonographic measurement of renal dimensions of adults in northwestern Nigeria a preliminary report, Sub-Saharan African Journal of medicine, year:2015, Volume:2, Issue:3, Page:123-127.

# Appendices

## Appendix 1



**Image 1:** coronal CT image for female (27years) show site measurements of RT Kidney length, width and cortical thickness diameters.



**Image 2:** coronal CT image for Male (33 years) show site measurements of RT, LT Kidney length, width and cortex thickness diameter.

## Appendix 2

*Sudan University of Science and Technology*

*College of Graduate Studies*

### **Measurement of renal size in Sudanese adult using Computed tomography**

A thesis submitted for partial Fulfillment for the Requirement of M.Sc  
Degree in Diagnostic Radiologic technology

#### **Data collection sheet**

PT Data			RT Kidney data				LT Kidney Data			
NO	Gender	Age Year	Length cm	Width cm	Cortex thickness cm	Texture HU	Length cm	Width cm	Cortex thickness cm	Texture HU

