

**Sudan University of Science and Technology**

**College Graduate Studies**

**Measurement of Common Carotid Arteries  
Intima Media Thickness Using  
Ultrasonography in Sudanese People**

**قياس سمك البطانة الوسطى للشرايين السباتية لدى السودانيين  
بالموجات فوق الصوتية**

Thesis Submitted for Partial Fulfillment for requirement of Master  
Degree (M.Sc) in Diagnostic ultrasound

By:

MenassJadallah Mohamed Alamin

Supervisor:

Dr. Mona Ahmed Mohammed Ahmed

2016

## الايه

بسم الله الرحمن الرحيم

قال تعالى:

(وَلَيْسَ الَّذِي خَلَقَ السَّمَوَاتِ وَالْأَرْضَ بِقَابِرٍ عَلَيَّ أَنْ يَخْلُقَ مِثْلَهُمْ بَدَلًا وَهُوَ الْخَلَّاقُ الْعَلِيمُ) [81] نَمَّا أَمْرُهُ  
إِنَّا أَرَادَ شَيْئًا أَنْ يَقُولَ لَهُ كُنْ فَيَكُونُ [82]

صدق الله العظيم

سورة يس الايات (81-82)

## Dedication

To:

My family

My friends

My colleagues

My teachers

## **Acknowledgment**

First of all, I Thank Allah for helping me to complete this project.I Thank Dr. Mona Ahmed my supervisor for her help and guidance, my great fullness for my all teachers in different educational levels.

Finally I would like to thank everybody who helped me in this project.

## **ABSTRACT**

The main objective of this study was to carry the measurement of carotid artery intima-media thickness in healthy Sudanese people. This was carried in order to account the variation in respect to international index and correlate CAIMT measurement with: gender, age and body mass index as well as compare between right and left CAIMT in Khartoum state, Sudan; Department of Ultrasound in Bashayir hospital for the period from May 2016 to September 2016.

The observations of study were included 50 healthy Sudanese males and females and the data were obtained from people who were examined for right and left main common carotid arteries. Therefore, the sonographic measurement of CAIMT was made in the longitudinal plane at the point of CCA along a section of the artery 1 cm proximal to the carotid bulb.

The study we found that the right main common carotid artery IMT range between 0.04cm to 0.073, while the left main common carotid artery IMT range between 0.04 cm to 0.07cm.

The study found that there was a weak degree of correlation between CAIMT with subject age and BMI while there was a strong degree of correlation between the values of right and left CAIMT. Hence, the study concluded that the CAIMT +values were higher in female than in male. Thus, there were weak degree of correlation between CAIMT and the subject age and BMI, and hence the CAIMT had higher values in right than left.

## مستخلص

تهدف هذه الدراسة اليمعرفة وقياس وسطبطانة الشريان السباتي عند الاشخاص الاصحاء من اجل معرفة الاختلاف مع المؤشرالعالمى وقياس ارتباط وسطبطانة الشريان السباتي مع كل من العمر - مؤشرالجس مفضلاعن مقارنة ذلك بين اليمين واليسارفي السودان.

اجريت الدراسة بولاية الخرطوم –السودان-قسم الموجات فوق الصوتية بمستشفى بشائرفي الفترة بين مايو 2016 اليسبتمبر 2016.

تضمنت الدراسة 50 مشاهدة من الاشخاص الاصحاء في السودان من الاناث والذكوروقد تحصلت الدراسة علي البيانات والمعلومات من الاشخاص الذين اجري عليهم الفحص للشريان السباتي المشترك لليمين واليسار. وباستخدام قي اسالسنوغراف كاطولي عند نقطة التقييم القطري المشترك علي طولي جزء من سمب القري من الشريان السباتي لهذه النقطة.

تفيد الدراسة بوجو دار تباط ضعي فبين كل من CAIMT والعمر ومؤشر كتلة الجسم في حين انها وجدت انهنال كدرجة عالية بين الارتباط بين قيما ليمينواليسار.

خلصت الدراسة الي انقيم CAIMT كانت اعلي في الاناث من الذكور كما كانتهنا كدرجة عالية بين علاقة كل من , CAIMT والعمر ومؤشر كتلة الجسم وCAIMT والقيم العلي الليسار.

## List of content

<b>Topic</b>	<b>Page</b>
الإيه	I
Dedication	II
Acknowledgments	III
Abstract	IV
Arabic Abstract	V
List of content	VI
List of tables	VIII
List of figures	IX
List of Appendices	X
Abbreviation	XI
<b>Chapter one</b>	
1.1 introduction	2
1.2 Problem	3
1.3 Objectives	3
1.4 Over view of the study	3
<b>Chapter two:Literature review</b>	
2.1 Theoretical Background	5
2.2 Previous Studies	30
<b>Chapter three M: arterials &amp; Methods</b>	
3.1Materials	32
3.1.1 patient	32

3.1.12 instrumentation	<b>32</b>
3.2 Methods	<b>32</b>
<b>Chapter four</b>	
Result	<b>35</b>
<b>Chapter five</b>	
Discussion	<b>43</b>
Conclusion	<b>45</b>
Recommendation	<b>46</b>
Reference	<b>47</b>
Appendix	<b>49</b>

Appendix 1	Data sheet
Appendix 2	<b>Images for explanation</b>
Image 1	Image No 1 longitudinal ultrasound image show measurement of carotid artery IMT.
Image 2	Image No 2 longitudinal ultrasound image show measurement of carotid artery IMT.
	<b>Images of the research</b>
Image 1	Image NO 1 longitudinal ultrasound image show measurement of carotid artery IMT for 31 years female with subject in supine, the neck extended and the head turned in the opposite direction.
Image 2	Image NO 2 longitudinal ultrasound image show measurement of carotid artery IMT for 20 years male with subject in supine, the neck extended and the head turned in the opposite direction.
Image 3	Image NO 3 longitudinal ultrasound image show measurement of carotid artery IMT for 25 years male with subject in supine, the neck extended and the head turned in the opposite direction.



## List of tables

Table 2-1	Table (2.1): Normal IMT values—median (P50), 25th and 75th percentile (P) IMT values for men and women at different age categories, separately for right (A) and left (B) CCA (1)
Table 4-1	Tale 4.1 showing the mean, standerdiviation, minimum and maximum values according to age, body mass index (BMI), right main intima media thickness and left main intima media thickness .

## List of figures

figure	Name of figure
Figure 2-1	Figure (2.1): The common carotid artery arises directly from the aorta on the left, and as a branch of the brachiocephalic trunk on the right (Wikipedia ,2016)
Figure 2-2	Figure (2.2): carotid artery in the neck (Stephanie et al, 2004).
Figure 2-3	Figure (2.3): Arch aortogram : subtraction film . The catheter is in the aortic arch . All branches of the external carotid are shown except the occipital artery. The facial artery is not fillin g on the right (Stephanie et al,2004).
Figure 2-4	Figure(2.4)structure of arterial wall (nanobme ,2016)
Figure 2-5	Figure (2.5): The optimal position for scanning the carotid arteries (Hartshorne et al,2005).
Figure 2-6	Figure (2.6): Transverse B-mode images. A: CCA and jugular vein. B: The ICA and ECA just above the carotid bifurcation (Hartshorne et al,2005).
Figure 2-7	Figure (2.7): Longitudinal scan planes used to visualize the carotid arteries. A: Posterior. B: Lateral. C: Anterior (Hartshorne et al,2005).
Figure 2-8	Figure (2.8): B-mode image of the ECA showing the superior thyroid branch (arrow) (Hartshorne et al,2005).
Figure 2-9	Figure (2.9): Longitudinal B-mode image of the carotid bifurcation with the ICA and ECA seen in the same plane. The arrows mark where the intima-media layer can be seen (Hartshorne et al, 2005).
Figure2-10	Figure (2.10): Intima-media thickness (IMT) definition – IMT is measured as the distance between lumen-intima (yellow line) and media-adventitia (pink line) interfaces (Escardio,2015).
Figure2-11	Figure (2.11): Proper location for IMT measurement (Escardio,2015).
Figure 4-1	Figure 4.1 show age frequencies.
Figure 4-2	Figure 4.2 show account of Gender.
Figure 4-3	Figure 4.3 show distribution of Right main common carotid IMT measurement by cm for all patient.

Figure 4-4	Figure 4.4 show distribution of left main common carotid IMT measurement by cm for all subject.
Figure 4-5	Figure 4.5 show correlation between values of right main IMT measurement by cm and gender.
Figure 4-6	Figure 4.6 show correlation between values of left main IMT measurement by cm and gender.
Figure 4-7	Figure 4.7: A scatter plot diagram shows relationship between the LT main CCA IMT by cm (vertical )and the BMI (horizontal )of the subjects, $R^2=0.0428$ .
Figure 4-8	Figure 4.8: A scatter plot diagram shows relationship between the RT main CCA IMT by cm (vertical )and the BMI(horizontal ) of the subjects, $R^2=0.0231$ .
Figure 4-9	Figure 4.9: A scatter plot diagram shows relationship between the RT main CCA IMT by cm (vertical ) and the Age by year (horizontal) of the subject, $R^2=0.239$ .
Figure4-10	Figure 4.10: A scatter plot diagram shows relationship between the LT main CCA IMT by cm (vertical ) and the Age by year (horizontal) of the subject, $R^2=0.2556$ .
Figure4-11	Figure 4.11: A scatter plot diagram shows relationship between the RT main IMT by cm (vertical ) and the LT main IMT by cm (horizontal) of the subjects, $R^2=0.7468$ .

## **List of Abbreviations**

IMT	Intima media thickness
BMI	Body mass index
CCA	Common carotid artery
C	Cervical vertebra
ICA	Internal carotid artery
ECA	External carotid artery
CVD	Cardiovascular disease