

References

.

.

Alexander, .C .M. ,Landsman. P.B., Teutsch,. S.M.(2003). defined metabolic syndrome, diabetes, and prevalence of coronary heart disease among NHANES III participants age 50 years and older; *Diabetes journal* .52(1) :1210-1214.

Aguilera, C., GilCampos, M., & Canete, R.. (2008). Alterations in plasma and tissue lipids associated with obesity and metabolic syndrome. *Clinical Science*, 114(1): 183–193.

Assadi ,F.N.(2010). Hypomagnesemia: an evidence based approach to clinical cases;*Iran Journal of kidney diseases* .4(1):9-13.

Baaij,D.E. , Hoenderop,G.H., Bindels,J.G.(2012).Regulation of magnesium balance:lessons learned from human genetic disease; *Clinical Kidney Journal* .5(1):15-24..

Bishop, L. M., Edward, P., Fody .E, Schoeff, E., (2004). Clinical Chemistry :Principles, Procedures, Correlations .(5thed) (Bishop, Clinical Chemistry): Clinical correlation and analytic procedures. P,289 – 301.

Choi,J.h., Joseph,L., Pilote,L.(2013). Obesity and CRP in various populations ;*obesity review journal* .14(3),232–244.

Daniel.J.,Rader.J.,Helen.H.,(2011).disorder of lipid metabolism ;*Harrison principle of internal medicine* ,volume 1(18):333-354.

Das, U. N.(2001) .Is obesity an inflammatory condition? *Nutrition*, 17(1), 953–966. available from :[http://dx.doi.org/10.1016/s0899-9007\(01\)00672-4](http://dx.doi.org/10.1016/s0899-9007(01)00672-4)

factors in women from Kolkata, W.B, India. *Arq Bras Cardiol*,96(1),38-46.

Faraj , M., Salem , N. (2012).C - reactive protein, Blood Cell - An Overview of Studies in Hematology, Dr. Terry Moschandreu (Ed.), ISBN: 978-953-51-0753-8, InTech, DOI: 10.5772/47735. Available from: <http://www.intechopen.com/books/blood-cell-an-overview-ofstudies-in-hematology/c-reactive-protein> .

Fernandez.J.R. ,Richard.N.P.,Zimian.M.W.,David.B.A. (2003). Is percentage body fat differentially related to body mass index in Hispanic Americans, African Americans, and European Americans? *American Journal of Clinical Nutrition*, 77(1): 71–75.

Ganguli .D.E., Das .N.U., Saha. m .I., Sanapala. K., Chaudhuri .D., Ghosh. S. (2011) Association between inflammatory markers and cardiovascular risk factors in women from Kolkata, W.B, India. *Arq Bras Cardiol*,96(1),38-46

Gupta,A.B., Dilip, K.U., Satyam ,R.I., Arvind ,K.B., Arya,T.S. (2015).To study the association of high sensitivity C-reactive protein with newly diagnosed DM; *Indian Academy of clinical medicine journal* ,16(1):5-12 .

Hajer,G. R., and Visseren,W.T.(2008). Adipose tissue dysfunction in obesity, diabetes, and vascular diseases ;*European Heart Journal* , 29(24): 2959-2971.

Hanyu ,O.R, Miida,T.F, Kosuge ,K.N. (2007). Preheparin lipoprotein lipase mass is a practical marker of insulin resistance in ambulatory type 2 diabetic patients treated with oral hypoglycemic agents. *Clinical Chemistry Acta* 384(1):118-123.

Heilbronn, L. K., & Clifton, P. M. (2002). C-reactive protein and coronary artery disease: influence of obesity, caloric restriction and weight loss. *Journal of Nutritional Biochemistry*, 13(1): 316–321.

Heseker, H., Schmid,A.(2000). Epidemiology of obesity. *Ther Umsch.* 57(8):478-81.

Horakova . D.P and Stejskal .D.A.(2010). Adiposity fatty acid binding protein and C-reactive protein levels as indicators of insulin resistance , *Department of Preventive Medicine, Palacky University Olomouc*,155(4):35-59

IMayama .I., ulrich . C. M. ,Alfano .C.M .(2012) .Effects of a Caloric Restriction Weight Loss Diet and Exercise on Inflammatory Biomarkers in Overweight/Obese Postmenopausal Women, *cancer Res* : 72(9):2314-26.

Jahnen. W., Ketteler.M. (2012). Magnesium Basics; *Clinical Kidney Journal*, 5(1) 3-14.

Khan.M.Y., Haque.S.F., Quaiser.S.A ,(2011). C-Reactive Protein and TNF- α predict cardiovascular risks in obese Biomedical Research , *Kidney disease* 4(22):13-19.

Kushner .R.F., Roth .J.L.,(2003) .Assessment of the obese patient. *Endocrinol Metabo Clin N Am* ;32: 915-933.

Laurson .K .R., dam .D. ,Senchina.S. (2011).Age, sex, and ethnicity may modify the influence of obesity on inflammation. *Journal of Investigation Medicine* , 59(1),27-31

Malpuech. B.C., Nowacki .W., Daveau .M., Gueux .E., Linard .C.,Rock .E., Lebreton .J., Mazur .A., Rayssiguier.Y .(2000). Inflammatory response following acute magnesium deficiency in the rat .*Biochim Biophys* ,1501(2-3):91–98.

- Mantovani. A., Garlanda .C., Doni .A., Bottazzi. B.,** (2008). Pentraxins in Innate Immunity: from C- Reactive Protein to the Long Pentraxins PTX3, *Journal of Clinical Immunology*, Volume 28 : 1-13
- Marti A., Marcos A., & Martinez J. A.** (2001). Obesity and immune function relationships. *Obesity Reviews*, 2, 131–140.
- Micheal,L.B., Edward, P.F., Larry,E.S.**(2010) Clinical chemistry principles procedures correlations; 6th ed. Lippincou Williams and Wilkins ,p:328.
- Maria .P., and Evagelia. S.**(2009).obesity disease ;*health science journal* ,3(3),132-138 .
- Nakamura .H., Ito.H., Egami .Y .**(2008) .Waist circumference is the main determinant of elevated C-reactive protein in metabolic syndrome .*Diabetes Research Clinical Practical* ;79: 330–336.
- Nappo. F., Esposito. K., Cioffi. M., Giugliano. G., Molinari.A.M., Paolisso. G.,Marfella. R., Giugliano. D.** (2002) Postprandial endothelial activation in healthy subjects and in type 2 diabetic patients (role of fat and carbohydrate meals). *J. Am. Coll. Cardiol.* 39, 1145–1150.
- Park .H.S and park .J.Y,** (2005). Relationship of obesity and visceral adiposity with serum concentrations of CRP, TNF-[alpha] and IL-6. *Diabetes Research and Clinical Practice*, 69(1), 29–35.
- Pickup .J.C., Chusney .G.D., Mattock. M.B.** (2000).The innate immune response and type 2 diabetes: evidence that leptin is associated with a stress-related (acute-phase) reaction. *Clinical Endocrinology* 72(9):2314-2326.
- Ridker. P.M., Buring .J.E., Cook. N.R., Rifai .N.R.** (2003). C-reactive protein, the metabolic syndrome and risk of incident cardiovascular events:an 8-year follow-up of 14,719 ;*initially healthy American women Circulation* ,107(1):363-369
- Romani .A.** (2011). Cellular magnesium Homeostasis. *Archives of Biochemistry and Biophysics* ,512(1):1-23
- Sarinn .A., apakorn.V. , Wanicagool . W.** (2008). Association between hs-CRP and Hba1c in overweight type 2 diabetic female patients, *The FASEB Journal*,22(1):83-109.
- Sato. H., Miida. T., Wada .Y .**(2007). Atherosclerosis is accelerated in patients with long-term well-controlled systemic lupus erythematosus (SLE). *Clinical Chemistry Acta* ,385:35-42

Shils . M.E., Shike .C.M., Ross . A.C., Caballero . B. ,Cousins .R.J. (2006). disease *Modern nutrition in health and disease: Eleventh edition*. Wolters Kluwer Health Adis ,p 421.

Spiegel .D.M.,(2011).Magnesium in chronic kidney disease: unanswered questions. *Blood Purif* ,31(1-3), 172-176.

Swaminathan. R., (2003) .Magnesium metabolism and its disorders. *Clinical Biochemistry Review* ;24(2):47-66.

Visscher T. L. S., Seidell J. C., Molarius A., van der Kuip D., Hofman A., & Witteman J. C. (2001). A comparison of body mass index, waist-hip ratio and waist circumference as predictors of all-cause mortality among the elderly: The Rotterdam study. *International Journal of Obesity*, 25: 1730–1735.

WHO Technical Series (2000) Obesity: Preventing and managing the global epidemic. Report of a WHO con-sultation. *World Health Organization Technical Report Series*, **894**, 1-253.

Winkler.G.A., and Keszthelyi .S.K.(2003) Expression of tumor necrosis factor (TNF)-alpha protein in the subcutaneous and visceral adipose tissue in correlation with adipocyte cell volume, serum TNFalpha, soluble serum TNF-receptor-2 concentrations and C-peptide level;*European Endocrinol journal* , 149(2):129-135.

Yudkin.J.S., stehouwer.CD., Emeis.J.J.(1999).C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue; *Arterioscler Thrombo Vascular Biology*, 19(4),972-978.

Klop.B., Jukema.J.W., Rabelink.T.J., CastroCabezas.M. (2012)A physician’s guide for the management of hypertriglyceridemia: The etiology of hypertriglyceridemia determines treatment strategy. *Panminerva Medical journal* . 54, 91–103.

Bhatti MS, Akbri MZ, Shakoor M.(2001) Lipid profile in obesity ”. *Journal of Ayub Medical College Abbottabad*. 13(1):31-3.

Research questionnaire

Appendix I

**Sudan University of Science and Technology
College of Graduate Studies**

Evaluation of high sensitive C-reactive protein , lipid profile and serum magnesium level in non Diabetic Non hypertension obese Sudanese subject in Khartoum state

No ()

The name: ----- age: -----

Social status: ----- no of phone: -----

Weight: -----kg height: -----cm

BMI: -----

Blood pressure: -----

Systolic: ----- diastolic: -----

Waist circumference:-----cm hipscm

WTH ratio=.....

The disease the effect you

-There are history of diabetes in family? Yes () No ()

-Are you suffering from recurrent infection? Yes () No ()

-Are you abuse alcohol or cigarette? Yes () No ()

-There is history of obesity in family? Yes () NO ()

-Are you suffering from arthritis? Yes () No ()

Laboratory investigation:

1-hs C-reactive protein: ----- mg\l 2-HDL cholesterol -----mg\dl

3-total cholesterol:----- mg\dl 4-LDL cholesterol: -----mg\dl

5-magnesium level:-----mg\dl 6-triglyceride: (TG): ----- mg\dl