

Dedication

To the soul of my mother.....

To my dear father, brothers and
sisters.....

To my beloved wife

To my kind colleagues and friends
.....

I dedicated this work.

Acknowledgement

I would like to express my sincere thanks to my supervisor D.Munsoor Mohammed Munsoor for his efforts supervision and guidance during the course of this project. He is unlimited help and valuable comments and suggestions are highly appreciated.

My grateful thanks also extend to D.Amar Obubaker for his constant care, valuable advice and assistant.

Abstract

This is descriptive study, aimed at determining the frequency of Rh antigens (C, D, c, E, and e) and **kell1** among sickle cell patients . The study was conducted during approximately four months (from February 2010 to July 2010), an informed consent was obtained from all patients a total of forty venous blood samples were collected from sickle cell patients in a 2.5 ml EDTA containers .All samples were tested for Rh D antigen using the slide agglutination techniques, and the same samples tested for Rh C, c, E, e, and **kell1** antigens by immunodiffusion gel technique. The antigens were determined.

The results obtained showed that, the **c** antigen was most common frequency (100%), followed by **D** antigen occurred (90%), **e** antigen was (82.5%), and the least common in Rhesus system were antigen **C** (67.5%), and **E** (80%).

The **kell1** antigen was the least frequent and constitute (10%).

النتائج والنتائج

هذه دراسة وصفية أجريت فى مرضى الأنيميا المنجلية هدفت لتحديد نسب تردد فى مرضى الأنيميا المنجلية . أيضا الدراسة هدفت لتحديد (D, C, c, E,e) الأنتجينات فى مرضى الأنيميا المنجلية التى لها أهمية فى عمليات Kell1 نسبة تردد الأنتجين

. نقل الدم

لقد أستغرقت الدراسة أربعة شهور , أخذت الموافقة من الأشخاص الذين سحب منهم الدم و قد تم احاطتهم بأهداف البحث . تم تجميع العينات من أربعين مريض فى وعاء تم فحص جميع العينات . (EDTA) سعة 2.5 مل يحتوى على مادة مانعة لتجلط الدم باستخدام طريقة الشريحة . ونفس العينات فحصت بطريقة (D) لمعرفة الأنتجين Kell1 والأنتجين (C, c, E,e) تسمى جل (مانعة النفاذية) لتحديد الأنتجينات

هى الأكثر شيوعا (100%) . ويليهما c وأظهرت نتائج الدراسة أن نسبة تردد الأنتجين C و E(80%) وجدت (82.5%) و قد كانت الأنتجينات e ثم D الأنتجين هو الأقل ترددا بنسبة Kell1 هى أقل ترددا فى النظام الريبسى . الأنتجين (67.5%) (10%).

LIST OF ABBREVIATION

Item	Abbreviation
Rhesus.	Rh
Sickle cell disease.	SCD
Hemoglobin.	Hb
Kell.	K
Delayed hemolytic transfusion reaction.	DHTR
Hemolytic disease of the new born	HDN
Hemolytic transfusion reaction	HTR
Red blood cells	RBCs
Antibodies	Abs

LIST OF TABLES

Page No.	Subject	Table No
19	Comparison of nomenclature of antigen of Rh system	Table 1.1
20	The distribution of Rh phenotype.	Table 1.2
20	The frequency of Rh antigen in some Sudanese tribe.	Table 1.3
25	The racial distribution of kell phenotype	Table 1.4
25	The comparative positivity analysis of kell2	Table 1.5
26	The comparative positivity analysis of kell1	Table 1.6
26	The antigen of kell blood group and nomenclature.	Table 1.7
38	Frequency and percentage of Rh C among study population.	Table 3.1
39	Frequency and percentage of Rh c among study population.	Table 3.2
40	Frequency and percentage of Rh E among study population.	Table 3.3
41	Frequency and percentage of Rh e among study population.	Table 3.4
42	Frequency and percentage of Rh D among study population.	Table 3.5
43	Frequency and percentage of kell1 among study population.	Table 3.6
38	The count of patients	Table 3.7

LIST OF FIGURE

Page No.	Subject	Figure No
38	Frequency and percentage of Rh C among study population	Figure3-1
39	Frequency and percentage of Rh c among study population.	Figure 3-2
40	Frequency and percentage of Rh E among study population	Figure3-3
41	Frequency and percentage of Rh e among study population.	Figure 3-4
42	Frequency and percentage of Rh D among study population.	Figure 3-5
43	Frequency and percentage of kell1 among study population.	Figure 3.6

Page No.	Subject
---------------------	----------------

I	Dedication
II	Acknowledgement
III	Abstract English
IV	Abstract Arabic
V	List of abbreviation
VI	List of tables
VII	List of figure
VIII	List of Content
Chapter one	
1	1.Introduction and literature review
1	1.1.Genetic regulation:
2	.2.Pathogenesis of sickle cell disease1
3	1.3. Clinical manifestation
3	.3.1Crises 1
4	.3.2.Pain episodes1
4	.3.3.Infections
5	.3.4.Eye disease1
4	1.3.5. Neurological complications
5	1.3.6.Pulmonary complications
6	1.3.7.Renal abnormalities
6	.4.Homozygous sickle cell disease: 1
6	.5.Sickle cell trait1
7	.6.Other sickling disorders:1
7	.7. Diagnosis of sickle cell disease: 1
7	1.7.1. Tests for Hb S
7	1.7.1.1. Sickling in Whole Blood
7	1.7.1.2. Reagents
8	1.7.1.3. Method
8	1.7.2. Hb S Solubility Test
8	1.7.2.1. Principle
8	1.7.2.2. Reagents
8	1.7.2.3. Method
9	1.7.2.4. Interpretation and Comments
10	1.7.3.Cellulose Acetate Electrophoresis at Alkaline pH
10	1.7.3.1.Principle
10	1.7.3.2. Equipment
11	1.7.3.3.Reagents
11	1.7.3.4.Method
12	1.7.4. Neonatal Screening
13	1.8. Management of the sickle cell patient
14	1.8.1. Painful sickle crises

15	1.8.2. Transfusion therapy
17	1.8.3. Hydroxyurea
18	1.9.1 Nomenclature and genetic theories of Rhesus blood group system
18	1.9.1.1. The Fisher- Race theory
19	1.9.1.3 The Rosen field nomenclature
20	1.9.2. The distribution and previous studies of RH blood group antigen.
20	1.9.3. The frequency of Rh antigens in some Sudanese tribes
20	1.9.4. The Rhesus antigen and encoding gene
22	1.9.5. RHNULLE syndrome
22	1.9.6. The Rhesus antibody
23	1.9.17. Clinical significant of Rh system
23	1.10. The genetic of Kell blood group system
24	1.10.1. The antigen of Kell blood group system
25	1.10.1.1 Racial distribution of kell type
27	1.10.1.2 KEL1 and KEL2 (K and k) antigens
27	1.10.1.3 KEL3 and KEL4 (Kpa and Kpb)
27	1.10.1.4 KEL 21 (KPC)
28	1.10.1.5 KEL6 and KEL7 (JSa and JSb)
28	1.10.1.6 Common Kell phenotype
28	1.10.1.7 Null phenotype
28	1.10.1.8 McLeod syndrome
29	1.10.2. The antibody of Kell blood group system
29	1.10.3. The clinical significant of Kell blood group system
29	1.10.3.1 Transfusion reaction
29	1.10.3.2. Hemolytic disease of new born
Chapter Two	
33	2. MATERIAL AND METHODS
33	2.1. Study area
33	2.2. Study design
33	2.3. Study population
33	2.3.1. Inclusion criteria
33	2.3.2. Exclusion criteria
33	2.4. Ethical consideration
33	2.5. Data collection
33	2.6. Data processing
33	2.7. Data presentation
34	2.8. Study period
34	2.9. Sampling
34	2.10. Techniques used
34	2.10.1. RhD grouping method

34	2.10.1.1. Principle of RhD slide grouping method
34	2.10.1.2. Reagents
34	2.10.1.3. Test procedures
34	2.10.1.4. Interpretation of results
35	2.10.2. Rh C, c, E, e and Kell1 grouping method
35	2.10.2.1. General principle of Gel Immune Diffusion technique
35	2.10.2.2. Reagents
35	2.10.2.3. Equipments
36	2.10.2.4. Preparation of blood sample
36	2.10.2.5 Test procedure
	2.10.2.6. Interpretation of the result
Chapter Three	
37	3. RESULT
Chapter Four	
46	4. DISCUSSION
Chapter Five	
	5. Conclusion & Recommendations
49	5.1 Conclusion
50	5.2 Recommendations
References	
Appendixes	
52	Appendix: 1 Questionnaire