



Groundwater Quality Assessment of Some Selected Hand-dug Wells in Wuntin Dada Community of Bauchi- Northwest Nigeria

Yusuf Usman Muhammad¹, Abdelatif Mokhtar Ahmed¹, Josiah Nuhu Jabbo² and Ahmed Abubakar Sarki³

Geology Department, Sudan University of Science and Technology, Khartoum-Sudan, abdanihamid@gmail.com

Department of Geological Technology, School of Science and Technology, Abubakar Tatari Ali Polytechnic Bauchi, P.M.B. 0094, Bauchi State, Nigeria

Geology Department, Faculty of Earth and Environmental Science, Bayero University Kano, igeria.ym_usman@yahoo.com

ARTICLE INFO

ARTICLE HISTORY

Received:24/10/2019

Accepted:1/12/2019

Available

online:June2020

KEYWORDS: Wuntin Dada, potable water, hard water and disinfection

ABSTRACT

This investigation assessed the water quality of ten selected hand-dug wells which for quite sometimes serve as alternative sources of drinking water of the *Wuntin Dada* inhabitants due to inadequate public supply of *potable water*. Water from each of the selected hand-dug wells was sampled in a sterilized 500ml plastic container and taken to the laboratory for analysis. Samples were analysed for physical, chemical and microbial parameters. Physical parameters analysed include; pH, Temperature, Turbidity, Electrical Conductivity and Total Dissolved Solids. The mean values of all the physical parameters are within the standards set by the Nigerian Industrial Standard (NIS, 2015) as stipulated in the Nigerian Standards for Drinking Water Quality (NSDWQ) guidelines. Waters from the water sources investigated are therefore safe as far as the physical quality is concerned. Chemical parameters analysed include; Sodium, Calcium, Magnesium, Iron, Lead, and Arsenic. Calcium and Magnesium concentrations are the only two parameters with concentrations higher than the maximum permissible levels set by the Nigerian Standard for Drinking Water Quality (NSDWQ). Calcium mean value is 184mg/l against the maximum permissible level (mpl) of 75mg/l while that of Magnesium is 70mg/l against the maximum permissible level (mpl) of 20mg/l. The rest are within the maximum

permissible levels set by the Nigerian Industrial Standard (NIS), 2015 standards for drinking water quality. Total hardness is generally high in all the samples with range value of 154 -255mg/l. Water hardness is the traditional measure of the capacity of water to react with soap, hard water requiring considerably more soap to produce lather (WHO, 2011). The water sources investigated provide the inhabitant with *hard water*. The mean value for Faecal Coliform is 1.6cfu/100ml against the maximum permissible level of 0.00cfu/100ml was rather alarming. This is an indication that the water sources under investigation were recently contaminated and may contain pathogens responsible for water borne diseases. Periodic *disinfection* of the water sources is highly recommended.

Introduction:

Access to safe drinking-water is essential to health, a basic human right and a component of effective policy for health protection (WHO, 2008). A person requires about three liters of potable water per day to maintain the essential fluids of the body (Fetter, 2007). Water is the major constituent of the human body, since sixty percent of adult body weight is due to water (WET, 2010). Water related diseases remain an issue of major global public health concern, with the water-sanitation-hygiene risk complex globally accounting for about ten percent of the global burden of disease (Pruess et al., 2008).

The quality of water obtained from hand-dug wells must be assessed considering the reliance on such sources of water by the inhabitants for their daily consumption and other uses. This study would be helpful in the determination of whether the water sources investigated are safe and therefore suitable for human consumption or not.

The area under investigation is Wuntin Dada community situated about seven kilometers northwest of Bauchi, the State Capital of Bauchi State in the northeastern Nigeria. The study area is underlain by crystalline rocks of the Nigerian Basement Complex. There is need to investigate the quality of water people consumes in the absence of public water supply, as water for human consumption need to be free from pathogens like bacteria, toxic substance like arsenic and chemical substances in concentration large enough to affect human health.

There is a steady rise in the demand for groundwater in most hard rock areas, most of which cannot boast of any constant surface source of water supply (Adamu, 1994). Oyawoye (1972) described the hydrogeologic condition found in Bauchi; he concluded that water is found in the weathered and fractured zone within the Basement Complex. Groundwater is generally less susceptible to contamination and pollution when compared to surface water bodies.

Josiah et al, (2016); Pavendan et al., (2011); Musa et al., (1999) and Elueze et al, (2004) have conducted similar investigations at Bauchi- NE Nigeria Tiruchirappali District-South India, Peri-urban communities-Northern Sudan and Ilesha area- SW Nigeria respectively. The sole aim of such assessments is to determine the physical, chemical as well as the microbial quality of the water sources investigated. Results from such

investigations determine suitability or otherwise of a particular water source to serve intended purposes.

Materials and Methods:

For the purpose of collecting water samples in the study area, the following instruments; Global Positioning System (GPS eTrex 10), TDS/EC Meter and pH/Temperature Meter were used. The Gamin eTrex 10 GPS was used to get the coordinates (Longitudes and Latitudes) of the location of water source sampled. Total Dissolve Solid (TDS) and Electrical Conductivity of each the water samples were measured by the TDS/EC Meter in the field. Hydrogen ion concentration (pH) and Temperature of the samples were also measured using pH/Temperature Meter.

Ten water samples (S1 –S10) from ten selected Hand-Dug Wells in different parts of Wuntin Dada community were collected and analysed for the physical, chemical and microbial parameters. Water sample from each of the ten hand-dug wells was collected in 500ml sterilized plastic containers and were instantly labeled according to sample’s location. A sterilized plastic cup was also used to fetch water sample for instant measurement of some physical parameters such as pH, Temperature, Total Dissolved Solid and Electrical Conductivity. Finally, the water samples collected were preserved in the refrigerator of the laboratory before the analysis was done.

Results:

The ten water samples under investigation were analysed for physical, chemical and microbial parameters. The results of the physical, chemical and microbial parameters are as shown in tables 1 .2a, 2b and 3 respectively.

Table (1) Physical Parameters Result:

Location	pH	Temperature (0C)	Turbidity (NTU)	Electrical Conductivity (uS/cm)	Total Dissolved Solid (mg/l)
Near Wikki	8.4	32	2.8	718	376
Wuntin Dada Primary	7.9	33	2.3	490	245
Wuntin Dada Market	8.6	30	4.2	1150	538
Mato Lame	7.7	31	3.4	660	323
Malam Abdullahi	7.6	31	3.1	980	550
Kara I	8.3	32	4.2	1020	545
Kara II	8.7	31	2.7	979	534
Kara III	8.5	30	3.9	740	373
Kara IV	8.4	31	4.0	770	340
IBB Squaare	7.9	30	4.1	1035	540
Range	7.6-8.7	30-33	2.3-4.2	490-1150	245-550
Mean	8.2	31.10	3.5	854	436
NSDWQ*MPL	6.5-8.5	Ambient	5	1000	500

NSDWQ: Nigerian Standard for Drinking Water Quality MPL: *Maximum Permissible Level

Table (2a) Chemical Parameters Result

Location	Total Alkalinity (mg/l)	Total Hardness (mg/l)	Total Iron, Fe ²⁺ (mg/l)	Calcium, Ca ²⁺ (mg/l)	Magnesium, Mg ²⁺ (mg/l)	Sodium, Na ⁺ (mg/l)
Near Wikki	120	184	0.2	106	78	5.3
W/Dada	115	160	0.3	111	49	1.7
Primary						
W/Dada	160	258	0.1	179	79	0.3
Market						
Mato Lame	144	154	0.5	107	47	4.0
Malam	78	248	0.2	199	49	1.9
Abdullahi						
Kara I	224	156	0.1	117	39	3.8
Kara II	180	378	0.1	286	92	1.1
Kara III	188	288	0.2	201	87	1.6
Kara IV	136	350	0.4	239	111	0.7
IBB Square	168	370	0.1	290	80	0.3
Range	78-188	156-378	0.1-0.5	106-290	39-111	0.3-5.3
Mean	151	255	0.2	184	71	2.1
NSDWQ	100	150	0.3	75	20	200
*MPL						

NSDWQ: Nigerian Standard for Drinking Water Quality

*MPL: Maximum Permissible Limit

Table (2b) Chemical Parameters Result

Location	Lead, Pb ²⁺ (mg/l)	Arsenic, As ⁺ (mg/l)	Chloride, Cl ⁻ (mg/l)	Fluoride, F ⁻ (mg/l)	Sulphate, SO ₄ ²⁻ (mg/l)	Nitrate, NO ₃ ⁻ (mg/l)	Nitrite, NO ₂ ⁻ (mg/l)
Near Wikki	0.00	0.00	8.2	0.9	116	10	0.1
Wuntin Dada	0.001	0.00	2.6	0.7	72	14	0.1
Primary							
Wuntin Dada	0.00	0.00	0.4	1.2	107	23	0.6
Market							
Mato Lame	0.00	0.00	6.1	0.6	60	18	1.1
Malam Abdullahi	0.00	0.002	3.0	1.4	145	5	0.1
Kara I	0.00	0.00	5.8	1.1	12	22	6.4
Kara II	0.00	0.00	1.6	0.8	70	10	0.1
Kara III	0.00	0.001	2.4	0.3	131	21	8.7
Kara IV	0.00	0.00	1.0	0.0	37	17	0.36
IBB Square	0.00	0.00	0.5	0.1	69	17	0.2
Range	0.0-0.001	0.0-0.002	0.4-8.2	0.0-1.1.4	12-145	5-23	0.1-8.7
Mean	0.00	0.00	0.3	0.7	82	16	1.8
NSDWQ	0.01	0.01	250	1.5	100	50	0.2
*MPL							

NSDWQ: Nigerian Standard for Drinking Water Quality

*MPL: Maximum Permissible Limit

Table (3) Microbial Parameters Result

Location	Total Coliform (cfu/100ml)	Faecal Coliform (cfu/100ml)
Near Wikki	19	7
Wuntin Dada Primary	12	5
Wuntin Dada Market	0	0
Mato Lame	0	0
Malam Abdullahi	2	0
Kara I	0	0
Kara II	0	0
Kara III	0	0
Kara IV	13	3
IBB Square	9	1
Range	0-19	0-7
Mean	5.5	1.6
NSDWQ*MPL	10	0

NSDWQ: Nigerian Standard for Drinking Water Quality *MPL: Maximum Permissible Level

Discussion:

For the purpose of this investigation, water quality will be assessed based on the Nigerian Standard for Drinking Water Quality (NSDWQ) of the Nigerian Industrial Standard (NIS) provisions. Analysis results of the five physical parameters: pH, Temperature, Turbidity, Total Dissolved Solid (TDS) and Electrical conductivity (EC) as shown in table 1, indicated that the mean values of these parameters are within the permissible levels set by the Nigerian Standard for Drinking Water Quality (NSDWQ).

Six cations' (Sodium, Calcium, Magnesium, Iron, Lead, and Arsenic) concentrations were determined and their mean values, with the exception of Calcium and Magnesium, were below the maximum permissible levels set by the Nigerian Industrial Standard (NIS), 2015 standards for drinking water quality. Calcium mean value is 184mg/l against the maximum permissible level (mpl) of 75mg/l while that of Magnesium is 70mg/l against the maximum permissible level (mpl) of 20mg/l. The mean values for these two cations were much higher than permissible levels set by the Nigerian Standard for Drinking Water Quality (NSDWQ), consequently, water from these sources are classified as hard. Though Calcium and Magnesium are essential minerals and beneficial to human health in several respects, inadequate intake of either nutrient can result in adverse health consequences (WHO, 2011). Total Alkalinity and Total Hardness concentrations' mean values are 151mg/l and 255mg/l are higher than the maximum permissible levels (mpl) of 100mg/l and 255mg/l respectively.

On the other hand, five anions' (Chloride, Fluoride, Sulphate, Nitrate, and Nitrite) concentrations were also determined. The mean values of Chloride, Fluoride, Sulphate and Nitrate and are within the maximum permissible levels set by the Nigerian Standard for Drinking Water Quality (NSDWQ). The mean values for Nitrite is 1.8mg/l which is higher than the maximum permissible levels (mpl) of 0.2mg/l.

The mean values for Total and Faecal Coliforms as shown in table 3 are 5.5cfu/100ml and 1.6cfu/100ml respectively. While the mean value for Total Coliform was lower than maximum permissible level of 10cfu/100ml set by the Nigerian Standard for Drinking Water Quality (NSDWQ), the mean value for Faecal Coliform of 1.6cfu/100ml against the maximum permissible limit of 0.00cfu/100ml was rather high. This is an indication that the water sources under investigation were recently contaminated and may contain pathogens responsible for water borne diseases.

Conclusion:

This investigation assessed the water quality of ten selected hand-dug wells which for quite sometimes serve as alternative sources of drinking water of the Wuntin Dada inhabitants due to inadequate public supply of potable water. Water from each of the selected hand-dug wells was sampled in a sterilized plastic container and taken to the laboratory for analysis. Samples were analysed for physical, chemical and microbial parameters.

Physical parameters analysed include pH, Temperature, Turbidity, Electrical Conductivity (EC) and Total Dissolved Solids (TDS). The mean values of all the physical parameters are within the standards set by the Nigerian Industrial Standard (NIS, 2015) as stipulated in the Nigerian Standards for Drinking Water Quality (NSDWQ) guidelines.

Total hardness is generally high in all the samples with range value of 154 -255mg/l. Calcium and Magnesium concentrations were the only two cations with concentrations higher than the maximum permissible levels set by the Nigerian Standard for Drinking Water Quality (NSDWQ). Water hardness is the traditional measure of the capacity of water to react with soap, hard water requiring considerably more soap to produce lather (WHO, 2011).

The mean value for Faecal Coliform is 1.6cfu/100ml against the maximum permissible level of 0.00cfu/100ml was rather alarming. This is an indication that the water sources under investigation were recently contaminated and may contain pathogens responsible for water borne diseases.

Waters obtained from the water sources investigated are safe as far as the physical quality is concerned but periodic disinfection of the water sources is highly recommended to get rid of the microbial contamination. Toxic chemical constituents (Pb and As) were not found in all the sample analysed, this rule out presence of toxicity in the analysed samples. The water sources investigated provide the inhabitants with hard water which require more soap when washing.

Referance:

Adamu E.A.(1994) *Groundwater Development and Management in the Basement Complex Terrain in Zaria, Kaduna State.* Water Resources 4

Elueze A A, Omidran J O and Nton M E (2004) Hydrogeochemical Investigation Surface water and Groundwater around Ibokun, Ilesha area, SW Nigeria. *Journal of Mining and Geology* Volume 40(1) P57

Fetter, C. W. (2007) *Applied Hydrogeology*, Second Edition, CBS Publishers, New Delhi. P.1

Josiah N J, Francis X O, Yusuf U M, Nasir A M (2016) Water Quality Assessment of Some Selected Hand-dug Wells and a Borehole in NE part of Bauchi Metropolis, Bauchi-Nigeria, *Journal of Environment and Earth Science*

Musa H A, Shears P, Kafi Sand Elsabag S K (1999) Water Quality and Public Health in Northern Sudan: A study of Rural and Peri-urban Communities. *Journal of Applied Microbiology*. 1999

Nigeria Industrial Standard-NIS (2015) Guidelines for drinking water quality. Policy document (*Nigerian Standard for Drinking Water Quality- NSDWQ*)

Oyawoye M. O. (1972) *The Basement Complex of Nigeria.* In: Dessauvage TFJ. Whiteman A J (eds) *African Geology.* Ibadan University Press, P 67

Pavendan p, Selvan S A and Rajasekaran C S (2011) Physico Chemical and Microbial Assessment of Drinking Water From Different Water Sources of Tiruchirappali District-South India, Pelagia Research Library

Pruess et. al. (2008) Animal Waste, Water Quality and Human health. In: (WHO, 2012) **World Health Organization-WHO (2008)**, Guidelines for drinking water quality.

Water Education for Teachers -WET (2010) *Healthy Hydration*--WET Project

WHO (2011) Drinking Water Standards and Health, Advisories Office of Water, U. S. Environmental Protection Agency, Washington DC