

Reference List

- Abate, N., Chandalia, M., Cabo-Chan, A. V., Jr., Moe, O. W., and Sakhaee, K. (2004). "The metabolic syndrome and uric acid nephrolithiasis: novel features of renal manifestation of insulin resistance." *Kidney Int.*, 65(2), 386-392.
- Abboud, O. L., Osman, E. M., and Musa, A. R. (1989). "The aetiology of chronic renal failure in adult Sudanese patients." *Ann. Trop. Med. Parasitol.*, 83(4), 411-414.
- Aggour, A., Ziada, A. M., AbdelHamid, A. Z., AbdelRahman, S., and Morsi, A. (2009). "Metabolic stone composition in Egyptian children." *J. Pediatr. Urol.*, 5(2), 132-135.
- Al-Eisa, A. A., Al-Hunayyan, A., and Gupta, R. (2002). "Pediatric urolithiasis in Kuwait." *Int. Urol. Nephrol.*, 33(1), 3-6.
- Al-Hermi, B., and Abbas, B. (2003). "Cystinuria in arab countries." *Saudi. J. Kidney Dis. Transpl.*, 14(3), 358-366.
- Al-Rasheed, S., Al Jurayyan, N. A., Al Nasser, M. N., Al-Mugeiren, M. M., Al-Salloum, A. A., and Petterson, B. A. (1995). "Nephrolithiasis in children and adolescents in the South Western region of saudi arabia." *Saudi. J. Kidney Dis. Transpl.*, 6(4), 396-399.
- al-Rasheed, S. A., el-Faqih, S. R., Husain, I., Abdurrahman, M., and al-Mugeirin, M. M. (1995). "The aetiological and clinical pattern of childhood urolithiasis in Saudi Arabia." *Int. Urol. Nephrol.*, 27(4), 349-355.

- Alaya, A., Nouri, A., and Najjar, M. F. (2009). "Urolithiasis in Tunisian children: a study of 100 cases." *Saudi. J. Kidney Dis. Transpl.*, 20(6), 1096-1100.
- Ali, S. H., and Rifat, U. N. (2005). "Etiological and clinical patterns of childhood urolithiasis in Iraq." *Pediatr. Nephrol.*, 20(10), 1453-1457.
- Alon, U. S., Zimmerman, H., and Alon, M. (2004). "Evaluation and treatment of pediatric idiopathic urolithiasis-revisited." *Pediatr. Nephrol.*, 19(5), 516-520.
- Angwafo, F. F., III, Daudon, M., Wonkam, A., Kuwong, P. M., and Kropp, K. A. (2000). "Pediatric urolithiasis in sub-saharan Africa: a comparative study in two regions of Cameroon." *Eur. Urol.*, 37(1), 106-111.
- Asplin, J., Parks, J., Lingeman, J., Kahnoski, R., Mardis, H., Lacey, S., Goldfarb, D., Grasso, M., and Coe, F. (1998). "Supersaturation and stone composition in a network of dispersed treatment sites." *J. Urol.*, 159(6), 1821-1825.
- Asplin, J. R. (2002). "Hyperoxaluric calcium nephrolithiasis." *Endocrinol. Metab Clin. North Am.*, 31(4), 927-949.
- Asplin, J. R., and Coe, F. L. (2007). "Hyperoxaluria in kidney stone formers treated with modern bariatric surgery." *J. Urol.*, 177(2), 565-569.
- Baker, P. W., Coyle, P., Bais, R., and Rofe, A. M. (1993). "Influence of season, age, and sex on renal stone formation in South Australia." *Med. J. Aust.*, 159(6), 390-392.

- Balla, A. A., Salah, A. M., Khattab, A. H., Kambal, A., Bongartz, D., Hoppe, B., and Hesse, A. (1998). "Mineral composition of renal stones from the Sudan." *Urol. Int.*, 61(3), 154-156.
- Barcelo, P., Wuhl, O., Servitge, E., Rousaud, A., and Pak, C. Y. (1993). "Randomized double-blind study of potassium citrate in idiopathic hypocitraturic calcium nephrolithiasis." *J. Urol.*, 150(6), 1761-1764.
- Belhadj, R., Marrakchi, O., Zerelli, L., Mohsni, A., Chebil, M., and Abdelmoula, J. (2008). "[Cystine lithiasis: retrospective study about 25 cases]." *Tunis Med.*, 86(6), 556-559.
- Bellizzi, V., De, N. L., Minutolo, R., Russo, D., Cianciaruso, B., Andreucci, M., Conte, G., and Andreucci, V. E. (1999). "Effects of water hardness on urinary risk factors for kidney stones in patients with idiopathic nephrolithiasis." *Nephron*, 81 Suppl 1, 66-70.
- Bergsland, K. J., Kelly, J. K., Coe, B. J., and Coe, F. L. (2006). "Urine protein markers distinguish stone-forming from non-stone-forming relatives of calcium stone formers." *Am. J. Physiol Renal Physiol*, 291(3), F530-F536.
- Bisceglia, L., Purroy, J., Jimenez-Vidal, M., d'Adamo, A. P., Rousaud, F., Beccia, E., Penza, R., Rizzoni, G., Gallucci, M., Palacin, M., Gasparini, P., Nunes, V., and Zelante, L. (2001). "Cystinuria type I: identification of eight new mutations in SLC3A1." *Kidney Int.*, 59(4), 1250-1256.

- Borghi, L., Meschi, T., Amato, F., Briganti, A., Novarini, A., and Giannini, A. (1996). "Urinary volume, water and recurrences in idiopathic calcium nephrolithiasis: a 5-year randomized prospective study." *J. Urol.*, 155(3), 839-843.
- Borghi, L., Meschi, T., Guerra, A., and Novarini, A. (1993). "Randomized prospective study of a nonthiazide diuretic, indapamide, in preventing calcium stone recurrences." *J. Cardiovasc. Pharmacol.*, 22 Suppl 6, S78-S86.
- Borghi, L., Schianchi, T., Meschi, T., Guerra, A., Allegri, F., Maggiore, U., and Novarini, A. (2002). "Comparison of two diets for the prevention of recurrent stones in idiopathic hypercalciuria." *N. Engl. J. Med.*, 346(2), 77-84.
- Breslau, N. A., Brinkley, L., Hill, K. D., and Pak, C. Y. (1988). "Relationship of animal protein-rich diet to kidney stone formation and calcium metabolism." *J. Clin. Endocrinol. Metab.*, 66(1), 140-146.
- Calonge, M. J., Gasparini, P., Chillaron, J., Chillon, M., Gallucci, M., Rousaud, F., Zelante, L., Testar, X., Dallapiccola, B., Di, S. F., and . (1994). "Cystinuria caused by mutations in rBAT, a gene involved in the transport of cystine." *Nat. Genet.*, 6(4), 420-425.
- Camargo, S. M., Singer, D., Makrides, V., Huggel, K., Pos, K. M., Wagner, C. A., Kuba, K., Danilczyk, U., Skovby, F., Kleta, R., Penninger, J. M., and Verrey, F. (2009). "Tissue-specific amino acid transporter partners ACE2 and collectrin differentially interact with hartnup mutations." *Gastroenterology*, 136(3), 872-882.

- Carlisle, E. J., Donnelly, S. M., and Halperin, M. L. (1991). "Renal tubular acidosis (RTA): recognize the ammonium defect and forget the urine pH." *Pediatr. Nephrol.*, 5(2), 242-248.
- Caudarella, R., Rizzoli, E., Buffa, A., Bottura, A., and Stefoni, S. (1998). "Comparative study of the influence of 3 types of mineral water in patients with idiopathic calcium lithiasis." *J. Urol.*, 159(3), 658-663.
- Chillaron, J., Estevez, R., Samarzija, I., Waldegger, S., Testar, X., Lang, F., Zorzano, A., Busch, A., and Palacin, M. (1997). "An intracellular trafficking defect in type I cystinuria rBAT mutants M467T and M467K." *J. Biol. Chem.*, 272(14), 9543-9549.
- Chou, Y. H., Li, C. C., Wu, W. J., Juan, Y. S., Huang, S. P., Lee, Y. C., Liu, C. C., Li, W. M., Huang, C. H., and Chang, A. W. (2007). "Urinary stone analysis of 1,000 patients in southern Taiwan." *Kaohsiung J. Med. Sci.*, 23(2), 63-66.
- Coe, F. L., Evan, A., and Worcester, E. (2005). "Kidney stone disease." *J. Clin. Invest.*, 115(10), 2598-2608.
- Coe, F. L., Favus, M. J., Crockett, T., Strauss, A. L., Parks, J. H., Porat, A., Gantt, C. L., and Sherwood, L. M. (1982). "Effects of low-calcium diet on urine calcium excretion, parathyroid function and serum 1,25(OH)₂D₃ levels in patients with idiopathic hypercalciuria and in normal subjects." *Am. J. Med.*, 72(1), 25-32.
- Coe, F. L., Parks, J. H., and Asplin, J. R. (1992). "The pathogenesis and treatment of kidney stones." *N. Engl. J. Med.*, 327(16), 1141-1152.

- Coe, F. L., Parks, J. H., Bushinsky, D. A., Langman, C. B., and Favus, M. J. (1988). "Chlorthalidone promotes mineral retention in patients with idiopathic hypercalciuria." *Kidney Int.*, 33(6), 1140-1146.
- Coe, F. L., Parks, J. H., and Moore, E. S. (1979). "Familial idiopathic hypercalciuria." *N. Engl. J. Med.*, 300(7), 337-340.
- Coll, D. M., Varanelli, M. J., and Smith, R. C. (2002). "Relationship of spontaneous passage of ureteral calculi to stone size and location as revealed by unenhanced helical CT." *AJR Am. J. Roentgenol.*, 178(1), 101-103.
- Coward, R. J., Peters, C. J., Duffy, P. G., Corry, D., Kellett, M. J., Choong, S., and van't Hoff, W. G. (2003). "Epidemiology of paediatric renal stone disease in the UK." *Arch. Dis. Child*, 88(11), 962-965.
- Curhan, G. C., Willett, W. C., Rimm, E. B., Spiegelman, D., and Stampfer, M. J. (1996). "Prospective study of beverage use and the risk of kidney stones." *Am. J. Epidemiol.*, 143(3), 240-247.
- Curhan, G. C., Willett, W. C., Rimm, E. B., and Stampfer, M. J. (1993). "A prospective study of dietary calcium and other nutrients and the risk of symptomatic kidney stones." *N. Engl. J. Med.*, 328(12), 833-838.
- Curhan, G. C., Willett, W. C., Rimm, E. B., and Stampfer, M. J. (1997). "Family history and risk of kidney stones." *J. Am. Soc. Nephrol.*, 8(10), 1568-1573.

- Curhan, G. C., Willett, W. C., Speizer, F. E., and Stampfer, M. J. (2001). "Twenty-four-hour urine chemistries and the risk of kidney stones among women and men." *Kidney Int.*, 59(6), 2290-2298.
- Dahlberg, P. J., van, d. B., Kurtz, S. B., Wilson, D. M., and Smith, L. H. (1977). "Clinical features and management of cystinuria." *Mayo Clin. Proc.*, 52(9), 533-542.
- Daudon, M. (2000). "[Component analysis of urinary calculi in the etiologic diagnosis of urolithiasis in the child]." *Arch. Pediatr.*, 7(8), 855-865.
- Daudon, M., Bader, C. A., and Jungers, P. (1993). "Urinary calculi: review of classification methods and correlations with etiology." *Scanning Microsc.*, 7(3), 1081-1104.
- Daudon, M., Bounxouei, B., Santa, C. F., Leite da, S. S., Diouf, B., Angwafoo, F. F., III, Talati, J., and Desrez, G. (2004). "[Composition of renal stones currently observed in non-industrialized countries]." *Prog. Urol.*, 14(6), 1151-1161.
- Daudon, M., and Jungers, P. (2004). "Drug-induced renal calculi: epidemiology, prevention and management." *Drugs*, 64(3), 245-275.
- Daudon, M., Lacour, B., and Jungers, P. (2006). "Influence of body size on urinary stone composition in men and women." *Urol. Res.*, 34(3), 193-199.
- DeFoor, W., Asplin, J., Jackson, E., Jackson, C., Reddy, P., Sheldon, C., Erhard, M., and Minevich, E. (2006). "Urinary metabolic evaluations in normal and stone forming children." *J. Urol.*, 176(4 Pt 2), 1793-1796.

- DeFoor, W., Asplin, J., Jackson, E., Jackson, C., Reddy, P., Sheldon, C., and Minevich, E. (2005). "Results of a prospective trial to compare normal urine supersaturation in children and adults." *J. Urol.*, 174(4 Pt 2), 1708-1710.
- Dello, S. L., and Rizzoni, G. (2006). "Cystinuria." *Acta Paediatr. Suppl*, 95(452), 31-33.
- Devuyst, O., and Pirson, Y. (2007). "Genetics of hypercalciuric stone forming diseases." *Kidney Int.*, 72(9), 1065-1072.
- Djelloul, Z., Djelloul, A., Bedjaoui, A., Kaid-Omar, Z., Attar, A., Daudon, M., and Addou, A. (2006). "Urinary stones in Western Algeria: study of the composition of 1,354 urinary stones in relation to their anatomical site and the age and gender of the patients." *Prog. Urol.*, 16(3), 328-335.
- Doumi, E. B., (2008). " Vesical calculi in El Obeid Hospital, Western Sudan." *Sudan Journal of Medical Science.*, 3(1), 29-32.
- Egoshi, K. I., Akakura, K., Kodama, T., and Ito, H. (2000). "Identification of five novel SLC3A1 (rBAT) gene mutations in Japanese cystinuria." *Kidney Int.*, 57(1), 25-32.
- Ekeruo, W. O., Tan, Y. H., Young, M. D., Dahm, P., Maloney, M. E., Mathias, B. J., Albala, D. M., and Preminger, G. M. (2004). "Metabolic risk factors and the impact of medical therapy on the management of nephrolithiasis in obese patients." *J. Urol.*, 172(1), 159-163.

el-Reshaid, K., Mughal, H., and Kapoor, M. (1997). "Epidemiological profile, mineral metabolic pattern and crystallographic analysis of urolithiasis in Kuwait." *Eur. J. Epidemiol.*, 13(2), 229-234.

Elliott, D., and Opas, L. M. (1999). "Consultation with the specialist: Renal stones." *Pediatr. Rev.*, 20(8), 280-2, quiz.

Ettinger, B., Pak, C. Y., Citron, J. T., Thomas, C., ms-Huet, B., and Vangessel, A. (1997). "Potassium-magnesium citrate is an effective prophylaxis against recurrent calcium oxalate nephrolithiasis." *J. Urol.*, 158(6), 2069-2073.

Evan, A. P., Coe, F. L., Lingeman, J. E., Shao, Y., Matlaga, B. R., Kim, S. C., Bledsoe, S. B., Sommer, A. J., Grynepas, M., Phillips, C. L., and Worcester, E. M. (2006). "Renal crystal deposits and histopathology in patients with cystine stones." *Kidney Int.*, 69(12), 2227-2235.

Evan, A. P., Coe, F. L., Lingeman, J. E., Shao, Y., Sommer, A. J., Bledsoe, S. B., Anderson, J. C., and Worcester, E. M. (2007). "Mechanism of formation of human calcium oxalate renal stones on Randall's plaque." *Anat. Rec. (Hoboken.)*, 290(10), 1315-1323.

Evan, A. P., Lingeman, J. E., Coe, F. L., Parks, J. H., Bledsoe, S. B., Shao, Y., Sommer, A. J., Paterson, R. F., Kuo, R. L., and Grynepas, M. (2003). "Randall's plaque of patients with nephrolithiasis begins in basement membranes of thin loops of Henle." *J. Clin. Invest.*, 111(5), 607-616.

- Evan, A. P., Lingeman, J. E., Coe, F. L., Shao, Y., Parks, J. H., Bledsoe, S. B., Phillips, C. L., Bonsib, S., Worcester, E. M., Sommer, A. J., Kim, S. C., Tinmouth, W. W., and Grynepas, M. (2005). "Crystal-associated nephropathy in patients with brushite nephrolithiasis." *Kidney Int.*, 67(2), 576-591.
- Fielding, J. R., Steele, G., Fox, L. A., Heller, H., and Loughlin, K. R. (1997). "Spiral computerized tomography in the evaluation of acute flank pain: a replacement for excretory urography." *J. Urol.*, 157(6), 2071-2073.
- Finlayson, B. (1974). "Symposium on renal lithiasis. Renal lithiasis in review." *Urol. Clin. North Am.*, 1(2), 181-212.
- Font-Llitjos, M., Jimenez-Vidal, M., Bisceglia, L., Di, P. M., de, S. L., Rousaud, F., Zelante, L., Palacin, M., and Nunes, V. (2005). "New insights into cystinuria: 40 new mutations, genotype-phenotype correlation, and digenic inheritance causing partial phenotype." *J. Med. Genet.*, 42(1), 58-68.
- Fraser, M., Joyce, A. D., Thomas, D. F., Eardley, I., and Clark, P. B. (1999). "Minimally invasive treatment of urinary tract calculi in children." *BJU. Int.*, 84(3), 339-342.
- Gasparini, P., Calonge, M. J., Bisceglia, L., Purroy, J., Dianzani, I., Notarangelo, A., Rousaud, F., Gallucci, M., Testar, X., Ponzzone, A., and . (1995). "Molecular genetics of cystinuria: identification of four new mutations and seven polymorphisms, and evidence for genetic heterogeneity." *Am. J. Hum. Genet.*, 57(4), 781-788.

- Gault, M. H., and Chafe, L. (2000). "Relationship of frequency, age, sex, stone weight and composition in 15,624 stones: comparison of results for 1980 to 1983 and 1995 to 1998." *J. Urol.*, 164(2), 302-307.
- Gillespie, R. S., and Stapleton, F. B. (2004). "Nephrolithiasis in children." *Pediatr. Rev.*, 25(4), 131-139.
- Goldfarb, D. S. (2004). "Microorganisms and calcium oxalate stone disease." *Nephron Physiol*, 98(2), 48-54.
- Goldfarb, D. S., Fischer, M. E., Keich, Y., and Goldberg, J. (2005). "A twin study of genetic and dietary influences on nephrolithiasis: a report from the Vietnam Era Twin (VET) Registry." *Kidney Int.*, 67(3), 1053-1061.
- Griffin, D. G. (2004). "A review of the heritability of idiopathic nephrolithiasis." *J. Clin. Pathol.*, 57(8), 793-796.
- Guillen, M., Corella, D., Cabello, M. L., Gonzalez, J. I., Sabater, A., Chaves, J. F., and Hernandez-Yago, J. (2005). "Identification of novel SLC3A1 gene mutations in Spanish cystinuria families and association with clinical phenotypes." *Clin. Genet.*, 67(3), 240-251.
- Hari, P., Bagga, A., Vasudev, V., Singh, M., and Srivastava, R. N. (1995). "Aetiology of nephrolithiasis in north Indian children." *Pediatr. Nephrol.*, 9(4), 474-475.
- Henry, R. (1974) *Clinical Chemistry: principles and technics*, 2nd ed., New York, NY: Harper&Row, 723.

- Hesse, A., Muller, R., Schneider, H. J., and Taubert, F. (1978). "[Analytic experiments on the significance of the fluorine content of urinary calculi (author's transl)]." *Urologe A*, 17(4), 207-210.
- Hesse, A., Siener, R., Heynck, H., and Jahnen, A. (1993). "The influence of dietary factors on the risk of urinary stone formation." *Scanning Microsc.*, 7(3), 1119-1127.
- Hofbauer, J., Hobarth, K., Szabo, N., and Marberger, M. (1994). "Alkali citrate prophylaxis in idiopathic recurrent calcium oxalate urolithiasis--a prospective randomized study." *Br. J. Urol.*, 73(4), 362-365.
- Holmes, R. P., Goodman, H. O., Hart, L. J., and Assimos, D. G. (1993). "Relationship of protein intake to urinary oxalate and glycolate excretion." *Kidney Int.*, 44(2), 366-372.
- Hulton, S. A. (2001). "Evaluation of urinary tract calculi in children." *Arch. Dis. Child*, 84(4), 320-323.
- Hymes, L. C., and Warshaw, B. L. (1987). "Thiazide diuretics for the treatment of children with idiopathic hypercalciuria and hematuria." *J. Urol.*, 138(5), 1217-1219.
- Ibrahim, A., Zein, M., and Beleil, O. (1977). "Uric acid lithiasis in the Sudan." *J. Trop. Med. Hyg.*, 80(8), 173-176.

- Jaeger, P., Portmann, L., Jacquet, A. F., and Burckhardt, P. (1984). "Drinking water for stone formers: is the calcium content relevant?" *Eur. Urol.*, 10(1), 53-54.
- Jungers, P., Joly, D., Gagnadoux, M. F., and Daudon, M. (2001). "[Cystine lithiasis: physiopathology and medical treatment]." *Prog. Urol.*, 11(1), 122-126.
- Junge. W., Wilke B, Halabi.A, Klein G. publication in preparation
- Kamoun, A., Daudon, M., Abdelmoula, J., Hamzaoui, M., Chaouachi, B., Houissa, T., Zghal, A., Ben, A. S., Belkahia, C., and Lakhoua, R. (1999). "Urolithiasis in Tunisian children: a study of 120 cases based on stone composition." *Pediatr. Nephrol.*, 13(9), 920-925.
- Kamoun, A., Daudon, M., Kabaar, N., Dhaoui, R., Ben, A. S., Zghal, A., Zerilli, L., Chaouachi, B., Belkahia, C., and Lakhoua, R. (1995). "[Etiologic factors of urinary lithiasis in Tunisian children]." *Prog. Urol.*, 5(6), 942-945.
- Kamoun, A., Zghal, A., Daudon, M., Ben, A. S., Zerelli, L., Abdelmoula, J., Chaouachi, B., Houissa, T., Belkahia, C., and Lakhoua, R. (1997). "[Urinary calculi in children: contribution of anamnesis, biological exploration and physical analysis of calculi to the etiological diagnosis]." *Arch. Pediatr.*, 4(7), 629-638.
- Kessler, T., and Hesse, A. (2000). "Cross-over study of the influence of bicarbonate-rich mineral water on urinary composition in comparison with sodium potassium citrate in healthy male subjects." *Br. J. Nutr.*, 84(6), 865-871.

- Kheradpir, M. H., and Bodaghi, E. (1990). "Childhood urolithiasis in Iran with special reference to staghorn calculi." *Urol. Int.*, 45(2), 99-103.
- Kim, H., Jo, M. K., Kwak, C., Park, S. K., Yoo, K. Y., Kang, D., and Lee, C. (2002). "Prevalence and epidemiologic characteristics of urolithiasis in Seoul, Korea." *Urology*, 59(4), 517-521.
- Klee, L. W., Brito, C. G., and Lingeman, J. E. (1991). "The clinical implications of brushite calculi." *J. Urol.*, 145(4), 715-718.
- Koh, S. S., Li, H., Lee, Y. H., Widelitz, R. B., Chuong, C. M., and Stallcup, M. R. (2002). "Synergistic coactivator function by coactivator-associated arginine methyltransferase (CARM) 1 and beta-catenin with two different classes of DNA-binding transcriptional activators." *J. Biol. Chem.*, 277(29), 26031-26035.
- Krieger, J. N., Kronmal, R. A., Coxon, V., Wortley, P., Thompson, L., and Sherrard, D. J. (1996). "Dietary and behavioral risk factors for urolithiasis: potential implications for prevention." *Am. J. Kidney Dis.*, 28(2), 195-201.
- Kumar, P., Clark, M., (2002). "Clinical Medicine". 15th ed. United Kingdom: W.B.Saunders, 625-633
- Kumar, V., and Lieske, J. C. (2006). "Protein regulation of intrarenal crystallization." *Curr. Opin. Nephrol. Hypertens.*, 15(4), 374-380.

- Kuo, R. L., Lingeman, J. E., Evan, A. P., Paterson, R. F., Parks, J. H., Bledsoe, S. B., Munch, L. C., and Coe, F. L. (2003). "Urine calcium and volume predict coverage of renal papilla by Randall's plaque." *Kidney Int.*, 64(6), 2150-2154.
- Laerum, E., and Larsen, S. (1984). "Thiazide prophylaxis of urolithiasis. A double-blind study in general practice." *Acta Med. Scand.*, 215(4), 383-389.
- Lee, G. S., Cho, I. S., Lee, Y. H., Noh, H. J., Shin, D. W., Lee, S. G., and Lee, T. Y. (2002). "Epidemiological study of clonorchiasis and metagonimiasis along the Geum-gang (River) in Okcheon-gun (county), Korea." *Korean J. Parasitol.*, 40(1), 9-16.
- LeeLewandowski, S., Rodgers, A., and Schloss, I. (2001). "The influence of a high-oxalate/low-calcium diet on calcium oxalate renal stone risk factors in non-stone-forming black and white South African subjects." *BJU. Int.*, 87(4), 307-311.
- Lieske, J. C., Monico, C. G., Holmes, W. S., Bergstralh, E. J., Slezak, J. M., Rohlinger, A. L., Olson, J. B., and Milliner, D. S. (2005). "International registry for primary hyperoxaluria." *Am. J. Nephrol.*, 25(3), 290-296.
- Lopez, M., and Hoppe, B. (2008). "History, epidemiology and regional diversities of urolithiasis." *Pediatr. Nephrol.*
- Mandel, N. S., and Mandel, G. S. (1989). "Urinary tract stone disease in the United States veteran population. II. Geographical analysis of variations in composition." *J. Urol.*, 142(6), 1516-1521.

- Marangella, M., Bagnis, C., Bruno, M., Vitale, C., Petrarulo, M., and Ramello, A. (2004). "Crystallization inhibitors in the pathophysiology and treatment of nephrolithiasis." *Urol. Int.*, 72 Suppl 1, 6-10.
- Massey, L. K., and Whiting, S. J. (1996). "Dietary salt, urinary calcium, and bone loss." *J. Bone Miner. Res.*, 11(6), 731-736.
- Matlaga, B. R., Shah, O. D., and Assimos, D. G. (2003). "Drug-induced urinary calculi." *Rev. Urol.*, 5(4), 227-231.
- Mehrsai, A., Taghizadeh, A. A., Zohrevand, R., Djaladat, H., Steffes, H., Hesse, A., and Pourmand, G. (2004). "Evaluation of urinary calculi by infrared spectroscopy." *Urol. J.*, 1(3), 191-194.
- Melton, L. J., III, Crowson, C. S., Khosla, S., Wilson, D. M., and O'Fallon, W. M. (1998). "Fracture risk among patients with urolithiasis: a population-based cohort study." *Kidney Int.*, 53(2), 459-464.
- Milliner, D. S. (2005). "The primary hyperoxalurias: an algorithm for diagnosis." *Am. J. Nephrol.*, 25(2), 154-160.
- Milliner, D. S., and Murphy, M. E. (1993). "Urolithiasis in pediatric patients." *Mayo Clin. Proc.*, 68(3), 241-248.
- Mora, C., Chillaron, J., Calonge, M. J., Forgo, J., Testar, X., Nunes, V., Murer, H., Zorzano, A., and Palacin, M. (1996). "The rBAT gene is responsible for L-cystine

- uptake via the b₀(+)-like amino acid transport system in a "renal proximal tubular" cell line (OK cells)." *J. Biol. Chem.*, 271(18), 10569-10576.
- Mosckovitz, R., Udenfriend, S., Felix, A., Heimer, E., and Tate, S. S. (1994). "Membrane topology of the rat kidney neutral and basic amino acid transporter." *FASEB J.*, 8(13), 1069-1074.
- Mshelia, D. S., Gali, B. M., Naaya, U. H., and Habu, S. A. (2005). "Chemical composition of urinary calculi in Maiduguri, Nigeria." *Afr. J. Med. Med. Sci.*, 34(2), 185-188.
- Muldowney, F. P., Freaney, R., and Moloney, M. F. (1982). "Importance of dietary sodium in the hypercalciuria syndrome." *Kidney Int.*, 22(3), 292-296.
- Oussama, A., Kzaiber, F., Mernari, B., Semmoud, A., and Daudon, M. (2000). "[Analysis of calculi by infrared spectroscopy in children from the Moroccan mid-Atlas region]." *Ann. Urol. (Paris)*, 34(6), 384-390.
- Ozalp, I., Coskun, T., Tokol, S., Demircin, G., and Monch, E. (1990). "Inherited metabolic disorders in Turkey." *J. Inherit. Metab Dis.*, 13(5), 732-738.
- Ozokutan, B. H., Kucukaydin, M., Gunduz, Z., Kabaklioglu, M., Okur, H., and Turan, C. (2000). "Urolithiasis in childhood." *Pediatr. Surg. Int.*, 16(1-2), 60-63.
- Pak, C. Y. (1988). "Medical management of nephrolithiasis in Dallas: update 1987." *J. Urol.*, 140(3), 461-467.

- Pak, C. Y., Poindexter, J. R., Peterson, R. D., Koska, J., and Sakhaee, K. (2002). "Biochemical distinction between hyperuricosuric calcium urolithiasis and gouty diathesis." *Urology*, 60(5), 789-794.
- Pak, C. Y., Sakhaee, K., Moe, O., Preminger, G. M., Poindexter, J. R., Peterson, R. D., Pietrow, P., and Ekeruo, W. (2003). "Biochemical profile of stone-forming patients with diabetes mellitus." *Urology*, 61(3), 523-527.
- Palacin, M., Fernandez, E., Chillaron, J., and Zorzano, A. (2001). "The amino acid transport system b(o,+) and cystinuria." *Mol. Membr. Biol.*, 18(1), 21-26.
- Parks, J. H., Worcester, E. M., Coe, F. L., Evan, A. P., and Lingeman, J. E. (2004). "Clinical implications of abundant calcium phosphate in routinely analyzed kidney stones." *Kidney Int.*, 66(2), 777-785.
- Parks, J. H., Worcester, E. M., O'Connor, R. C., and Coe, F. L. (2003). "Urine stone risk factors in nephrolithiasis patients with and without bowel disease." *Kidney Int.*, 63(1), 255-265.
- Parry, E. S., and Lister, I. S. (1975). "Sunlight and hypercalciuria." *Lancet*, 1(7915), 1063-1065.
- Pietrow, P. K., and Karellas, M. E. (2006). "Medical management of common urinary calculi." *Am. Fam. Physician*, 74(1), 86-94.
- Pietrow, P. K., Pope, J. C., Adams, M. C., Shyr, Y., and Brock, J. W., III (2002). "Clinical outcome of pediatric stone disease." *J. Urol.*, 167(2 Pt 1), 670-673.

- Polito, C., La, M. A., Cioce, F., Villani, J., Nappi, B., and Di, T. R. (2000). "Clinical presentation and natural course of idiopathic hypercalciuria in children." *Pediatr. Nephrol.*, 15(3-4), 211-214.
- Porena, M., Guiggi, P., and Micheli, C. (2007). "Prevention of stone disease." *Urol. Int.*, 79 Suppl 1, 37-46.
- Ramello, A., Vitale, C., and Marangella, M. (2000). "Epidemiology of nephrolithiasis." *J. Nephrol.*, 13 Suppl 3, S45-S50.
- Rieu, P. (2005). "[Infective lithiasis]." *Ann. Urol. (Paris)*, 39(1), 16-29.
- Rizvi, S. A., Naqvi, S. A., Hussain, Z., Hashmi, A., Hussain, M., Zafar, M. N., Mehdi, H., and Khalid, R. (2002). "The management of stone disease." *BJU. Int.*, 89 Suppl 1, 62-68.
- Robertson, W. G. (2003). "Renal stones in the tropics." *Semin. Nephrol.*, 23(1), 77-87.
- Robertson, W. G., Peacock, M., Heyburn, P. J., and Hanes, F. A. (1980). "Epidemiological risk factors in calcium stone disease." *Scand. J. Urol. Nephrol. Suppl*, 53, 15-30.
- Rodgers, A. (1999). "Effect of cola consumption on urinary biochemical and physicochemical risk factors associated with calcium oxalate urolithiasis." *Urol. Res.*, 27(1), 77-81.

- Saigal, C. S., Joyce, G., and Timilsina, A. R. (2005). "Direct and indirect costs of nephrolithiasis in an employed population: opportunity for disease management?" *Kidney Int.*, 68(4), 1808-1814.
- Saita, A., Bonaccorsi, A., and Motta, M. (2007). "Stone composition: where do we stand?" *Urol. Int.*, 79 Suppl 1, 16-19.
- Sakhaee, K., Harvey, J. A., Padalino, P. K., Whitson, P., and Pak, C. Y. (1993). "The potential role of salt abuse on the risk for kidney stone formation." *J. Urol.*, 150(2 Pt 1), 310-312.
- Sakhaee, K., ms-Huet, B., Moe, O. W., and Pak, C. Y. (2002). "Pathophysiologic basis for normouricosuric uric acid nephrolithiasis." *Kidney Int.*, 62(3), 971-979.
- Saso, L., Grippa, E., Gatto, M. T., and Silvestrini, B. (2001). "Inhibition of calcium oxalate precipitation by bile salts." *Int. J. Urol.*, 8(3), 124-127.
- Schwarzenbach. G., (1955) ,"The complexones and their analytical application." *Analyst.*,80,713-729.
- Seftel, A., and Resnick, M. I. (1990). "Metabolic evaluation of urolithiasis." *Urol. Clin. North Am.*, 17(1), 159-169.
- Seltzer, M. A., Low, R. K., McDonald, M., Shami, G. S., and Stoller, M. L. (1996). "Dietary manipulation with lemonade to treat hypocitraturic calcium nephrolithiasis." *J. Urol.*, 156(3), 907-909.

- Shah, A. M., Kalmunkar, S., Punekar, S. V., Billimoria, F. R., Bapat, S. D., and Deshmukh, S. S. (1991). "Spectrum of pediatric urolithiasis in western India." *Indian J. Pediatr.*, 58(4), 543-549.
- Shuster, J., Jenkins, A., Logan, C., Barnett, T., Riehle, R., Zackson, D., Wolfe, H., Dale, R., Daley, M., Malik, I., and . (1992). "Soft drink consumption and urinary stone recurrence: a randomized prevention trial." *J. Clin. Epidemiol.*, 45(8), 911-916.
- Siener, R., and Hesse, A. (2003). "Fluid intake and epidemiology of urolithiasis." *Eur. J. Clin. Nutr.*, 57 Suppl 2, S47-S51.
- Sierakowski, R., Finlayson, B., and Landes, R. (1979). "Stone incidence as related to water hardness in different geographical regions of the United States." *Urol. Res.*, 7(3), 157-160.
- Silverberg, S. J., Shane, E., Jacobs, T. P., Siris, E., and Bilezikian, J. P. (1999). "A 10-year prospective study of primary hyperparathyroidism with or without parathyroid surgery." *N. Engl. J. Med.*, 341(17), 1249-1255.
- Smith, R. C., Rosenfield, A. T., Choe, K. A., Essenmacher, K. R., Verga, M., Glickman, M. G., and Lange, R. C. (1995). "Acute flank pain: comparison of non-contrast-enhanced CT and intravenous urography." *Radiology*, 194(3), 789-794.
- Stamatelou, K. K., Francis, M. E., Jones, C. A., Nyberg, L. M., and Curhan, G. C. (2003). "Time trends in reported prevalence of kidney stones in the United States: 1976-1994." *Kidney Int.*, 63(5), 1817-1823.

- Stapleton, F. B. (2002). "Childhood stones." *Endocrinol. Metab Clin. North Am.*, 31(4), 1001-15, ix.
- Stapleton, F. B. (2003). "Imaging studies for childhood urinary infections." *N. Engl. J. Med.*, 348(3), 251-252.
- Stapleton, F. B., Roy, S., III, Noe, H. N., and Jerkins, G. (1984). "Hypercalciuria in children with hematuria." *N. Engl. J. Med.*, 310(21), 1345-1348.
- Sternberg, K., Greenfield, S. P., Williot, P., and Wan, J. (2005). "Pediatric stone disease: an evolving experience." *J. Urol.*, 174(4 Pt 2), 1711-1714.
- Suvachittanont, O., Meksongsee, L. A., Dhanamitta, S., and Valyasevi, A. (1973). "The oxalic acid content of some vegetables in Thailand, its possible relationships with the bladder stone disease." *J. Med. Assoc. Thai.*, 56(11), 645-653.
- Tamaoku, K., Ueno, K., Akiura, K., Ohkura, Y., (1982). New water soluble hydrogen donor for the enzymatic photometric determination of hydrogen peroxide. *Chem Pharm Bull* (30) 2492-2497
- Tang, R., Nancollas, G. H., Giocondi, J. L., Hoyer, J. R., and Orme, C. A. (2006). "Dual roles of brushite crystals in calcium oxalate crystallization provide physicochemical mechanisms underlying renal stone formation." *Kidney Int.*, 70(1), 71-78.
- Taylor, E. N., and Curhan, G. C. (2007). "Oxalate intake and the risk for nephrolithiasis." *J. Am. Soc. Nephrol.*, 18(7), 2198-2204.

- Trinchieri, A. (1996). "Epidemiology of urolithiasis." *Arch. Ital. Urol. Androl*, 68(4), 203-249.
- Wagner, C. A., Lang, F., and Broer, S. (2001). "Function and structure of heterodimeric amino acid transporters." *Am. J. Physiol Cell Physiol*, 281(4), C1077-C1093.
- Whitby.L.G, Percyrobb I.W ., Smith. A. F. "Lecture notes on clinical chemistry", 3ed ed . London : Black wells scientific publications, 1984: 107-109, 278-284, 327-330.
- Worcester, E. M. (2002). "Stones from bowel disease." *Endocrinol. Metab Clin. North Am.*, 31(4), 979-999.
- Worcester, E. M., and Coe, F. L. (2008). "Nephrolithiasis." *Prim. Care*, 35(2), 369-91, vii.
- Worcester, E. M., Gillen, D. L., Evan, A. P., Parks, J. H., Wright, K., Trumbore, L., Nakagawa, Y., and Coe, F. L. (2007). "Evidence that postprandial reduction of renal calcium reabsorption mediates hypercalciuria of patients with calcium nephrolithiasis." *Am. J. Physiol Renal Physiol*, 292(1), F66-F75.
- Worcester, E. M., Parks, J. H., Evan, A. P., and Coe, F. L. (2006). "Renal function in patients with nephrolithiasis." *J. Urol.*, 176(2), 600-603.
- Yagisawa, T., Chandhoke, P. S., and Fan, J. (1998). "Metabolic risk factors in patients with first-time and recurrent stone formations as determined by comprehensive metabolic evaluation." *Urology*, 52(5), 750-755.

Yoshida, O., Terai, A., Ohkawa, T., and Okada, Y. (1999). "National trend of the incidence of urolithiasis in Japan from 1965 to 1995." *Kidney Int.*, 56(5), 1899-1904.

Yuen, Y. P., Lam, C. W., Lai, C. K., Tong, S. F., Li, P. S., Tam, S., Kwan, E. Y., Chan, S. Y., Tsang, W. K., Chan, K. Y., Mak, W. L., Cheng, C. W., and Chan, Y. W. (2006). "Heterogeneous mutations in the SLC3A1 and SLC7A9 genes in Chinese patients with cystinuria." *Kidney Int.*, 69(1), 123-128.

Zarse, C. A., Deaton, E. A., and Weiner, W. W. (2004). "Light intensity appears to be more important than an endogenous seasonal clock for regulating structural rhythms in the lateral eye of the horseshoe crab." *Biomed. Sci. Instrum.*, 40, 407-412.