

## ACKNOWLEDGMENTS

I am greatly indebted to my advisor Dr. Saif Eldin Mohamed Kheir for his helpful guidance, advice and supervision throughout the study.

I express my thanks to my co-advisor Prof. Zohair E. Elabgar for his advice and criticism.

Thanks are extended to the members of the Pesticides Alternatives Division, the Environmental and Natural Resource Research Institute, National Research center for their kindness during plants extraction and preparation period.

Special thanks are extended to the staff of National Malaria Administration, Khartoum and Malaria Center, Sinnar for their generous attitudes during the laboratory work.

Special gratitude to Dr. Insaf Sh. Mohamed and Dr. Hassan Farag, Shambat Research Station of the ARC for their assistance in B.t supplying and thesis revising, respectively.

Thanks are extended to Sinnar university Administration, Ministry of Agriculture, Khartoum state, and Dr. Rudwan M. Ahmed, Rahad Agricultural scheme Manager for kind financial support.

Thanks are due to Dr. Mahammed H. Awad- Omdorman Islamic University and Dr. Abdelaziz Omer, Khartoum University for their assistance in the statistical analysis.

Thanks are due to the staff members of Plant Protection Department, Faculty of Agricultural studies, Shambat. Special thanks extended to Dr. Awad K. Taha.

Finally I am completely obliged to my small and extended family that backed and encouraged me all the time.