

**Table (3).The effect of Combination Malathion57 and NeemAzal-T/S (50:50%) compared with Malathion57 Recommended dose and NeemAzal-T/S Recommended dose against larvae and adults of *Tribolium castaneum***

<i>Treatments</i>	<i>Mortality Percentage</i>							
	1 <sup>st</sup> day		2 <sup>nd</sup> day		3 <sup>rd</sup> day		4 <sup>th</sup> day	
	Larva	Adult	Larva	Adult	Larva	Adult	Larva	Adult
Combination of Malthion57 and NeemAzal-T/S	57a	60b	70b	80b	77b	93a	97a	100a
Malathion57 Recommended Dose	43a	87a	83a	100a	97a	100a	100a	100a
NeemAzl-T/S Recommended Dose	3.3b	0bc	3.3bc	0bc	13bc	0bc	13bc	0bc
Control	0b	0bc	0bc	0bc	0bc	0bc	0bc	0bc
SE ±	6.72	3.28	3.98	3.29	5.43	2.92	4.65	0
LSD	20.14	9.83	11.93	9.86	16.28	8.76	13.93	0
C.V%	60.2	19.1	20.8	14.1	21.1	11.5	15.9	0

**Means followed by the same letter (s) are not significantly different at p 5%**