

References

- Ahmed, E.E.** (1988). Range Resource Inventory For Evaluation (El Odaya Project – Sudan). M.Sc. Thesis, ITC. The Netherland. (unpublished).
- Ahmed, J.E. and Bonham, C.D.** (1980). Algorithm and computer program for optimum allocation in multivariate double sampling for biomass estimation. Range Science Series No. 33. Range Science Department, Colo. State Univ., Ft. Collins, Colorado.
- Ahmed, J.E. Bonham, C.D., and Laylock, W.A.** (1983). Comparison of techniques used for adjusting biomass estimates by Double sampling, J. of range manage 36 : 217 – 222.
- Andrew, M.H., Noble, I.R. and Lange R.T.** (1979). A non-destructive method for estimating the weight of forage on shrubs. Aust. Rangl. J. 1: 225 – 231.
- Andrew, M.H., Noble, I.R., Lange R.T. and Johnson, A.W.** (1981). The measurements of shrub forage weight: Three methods compared. Aust. Rangl. J. 3: 74 – 82.
- Arab Organization for Agricultural Development-AOAD** – (1996). Conservation and development of the natural resources of *El Baja* area – WHITE NILE STATE.
- Ashby, E.** (1948). The quantitative analysis of vegetation. Ann. Bot. Lon. 49: 779 – 802.
- Barnes, D.L.** (1976). A review of plant-based methods of estimating food consumption. Percentage utilization, species preferences and feeding patterns of grazing and browsing animals. Proc. Grassld. Soc. Sth. Afr. 11: 65 – 71.
- Barnes, D.L., Odendaal, J.J. and Berkes. B.H.** (1988). Use of the Dry Weight Rank method of botanical analysis in the eastern Transvaal Highveld. Proc. Grassld. Soc. Sth. Afr. 17: 79 – 82.

- Bashir, M.** (2001). Sudan Country Study on Biodiversity. The national biodiversity strategy and action plan (NBSAP), project No. SUD/97/G 31.
- Barbour, M.C., Burk, J.H. and Pitts** (1987). Terrestrial plant ecology. Benjamin Cummins Publishing Co. Menlo Park, CA. 2nd ed. Pp 191 – 193.
- Bedell, T.E. (1973).** Botanical composition of sub-clover – grass pastures As affected by single and dual grazing by cattle and sheep. Agron. J. 65: 502 – 504.
- Behnke, R.H.** (1992). New Directions in African Range Management Policy. Workshop sponsored by the Commonwealth Secretariat, Matopos, Zimbabwe.
- Blair, R.M.** (1971). Forage production after hardwoods control in a Southern Pine hardwood stand. For. Sci. 17: 279 – 284.
- Bookhout, T.A.** (1965). Feeding co-actions between snowshoe hares and white-tailed deer in northern Michigan. Trans. N. Am. Wildl. Conf. 30: 321 – 335.
- Brown, D.** (1954). Methods of Surveying and Measuring Vegetation. Bucks, England: Commonwealth Agricultural Bur.
- Bryan, W.W.** (1968). Grazing trials on Wallum of south-eastern Queensland. 1. A comparison of four pastures. Aust. J. Exp. Agric. Anim. Husb. 8: 512-520.
- Bryan, W.W. and Evans, T.R.** (1973). Effects of soils, fertilizers and Stocking rates on pastures and beef production on the Wallum of South- Eastern Queensland. 1. Botanical composition and chemical effects on plants and soils. Aust. J. Exp. Agric. Anim. Hus. 2: 137 – 141.

- Bureau. Of Land Management** (1996). Sampling vegetation attributes. Interagency Technical Reference, BLM/RS/ST/ - 96 / 002 + 1730. CEGET/CNRS, Paris. Pp. 28 – 29.
- Carpenter, L.H., Wallmo, O.C. and Moriss, M.J.** (1973). Effects of woody stems on estimating herbage weights with a capacitance meter. *J. Range Manage.* 26 (2) : 151 – 152.
- Condon, R.W. (1968).** Estimation of grazing capacity on arid grazing Lands. P. 112 – 124. In: CSIRO, Symp. Land evaluation, Canberra. MacMillan and Co.
- Cook, C.W. and T.W. Box.** (1961). A comparison of the loop and point methods of analyzing vegetation. *J. Range Mangt.* 14: 22 – 27.
- Cook, C.W. and Stubbendiek** (1986). *Range Research: Basic problems and techniques.* Society for Range Management. Denver Co. pp 51 – 63.
- Cooper, C.F.** (1957). The variable plot method for estimating shrub density. *J. Range Manage.* 10: 111 – 115.
- Cooper, C.F.** (1959). Cover vs. density. *J. Range Mangt.* 12: 215.
- Cooper, C.F., Hyder, D.N, Petersen, R. G. and Sneva F.A.**(1957). The constituent differential method of estimated species composition in mixed hay. *Agron. J.* 49: 190 – 193.
- Crocker, R.L. and Tiver, N.S.** (1948). Survey methods in grassland ecology. *J.Br. grassld. Soc.* 3: 1 –26.
- Darag, A. and Suliman, M.** (1988). Training course in Range Management and Range Improvement – Range and Pasture Administration (RPA) – Sudan – 47 pp.
- De Vries, D.M. and De Boer, T.** (1959). Methods used in botanical grassland research in the Netherlands and their application. *Herbage Abstr.* 24: 1 – 7.

- Dekker, B., Kirkman, K.P. and Plessis, sl. Du** (2001). Use of the dry – weight rank method of botanical analysis in semi – arid savannah communities. *Afr. J. of Range & Forrest Science*. 18: 67 – 68.
- Donaldson, C.H. (1967)**. The immediate effects of the 1964/65 drought on the vegetation of specific study areas in the Vrybung district. *Proc. Grassld. Soc. Sth. Afr.* 2: 137 – 141.
- Du Toit, P.F.** (1968). A preliminary report on the effect of *Acacia karroo* competition on the composition and yield of sweet grassland. *Proc. Grassld. Soc. Sth. Afr.* 3: 147 – 149.
- El Samani, M.O. et al** (1995). Pastoral policies in the Sudan, In: Pastoral sector development conference – Kosti – Min. of Agric. & Nat. Res. – WHITE NILE STATE.
- Fadlalla, B.** (1987). The dry season nutritional status of transhumant Baggara sheep, *Proc. International Conference on Animal Production in Arid Zones (ICAPAZ)*, Damascus, Syria: 843 – 844.
- Foran, B.D. Tainton, N.M. and Booyesen, P.de V.** (1978). The development of a method for assessing veld condition in three grassveld types in Natal. *Proc. Grassld. Soc. Sth. Afr.* 13 : 27 – 33.
- Francis, R.C. Baker, C.V. and Gustafson, J.D.** (1971). A study of the weigh estimation method of botanical analysis. *U.S. Grassland Biome*, Tech. Report No. 117. Ft. Collins, Colorado. 147 pp.
- Friedel, M.H. and Shaw, K.** (1987). Evaluation of methods for monitoring sparse patterned vegetation in arid rangeland. I. Herbage. *J. Environ. Manage.* 25: 297 – 308.
- Friedel, M.H., Chewing, V.H. and Bastel, G.N.** (1988). The use of comparative yield and dry weight rank techniques for monitoring arid rangelands. *J. Range manage* 41: 430 – 435.

- Frost, W.E. and E.L. Smith** (1991). Biomass productivity and range condition on range sites in southern Arizona. *J. Range Mangt.* 44: 64 – 67.
- Gaiballa A.K. et al** (1997). Surveying of natural resources in *El Baja* area, with special emphasis on levels of utilization and management indicators for system sustainability (unpublished).
- Goodall, D.W.** (1952). Some considerations in the use of point quadrates for the analysis of vegetation. *Aus. J. Sci. Res. Ser. B*, 5 : 1 – 41.
- Graetz,R.D.** (1978). The influence of grazing sheep on the structure of a saltbush (*Atriplex vesicaria*) population. *Aust. Rangl. J.* 1: 117 – 125.
- Gray, S.G.** (1970). The place of trees and shrubs as sources of forage in tropical and subtropical pastures. *Trop. Grassld.* 4: 57 –62.
- Greig-Smith, P.** (1983). *Quantitative Plant Ecology*. 3rd. Ed. Butterworths, London. 359 pp.
- Guy, D.R.** (1976). The feeding behavior of elephant (*Loxodonta africana*) in the Sengwa area, Rhodesia. *S. Afr. J. Wildl. Res.* 6: 55 – 63.
- Hacker, R.B.** (1973). Range assessment theory. In: 2nd Rangeland management conference, Carnarvon, W.A. pp 9 – 31.
- Harrison, N.M. and Jackson, J.K.** (1958). Ecological classification of vegetation of the Sudan. *Forestry Bull.* 2 new series, Khartoum.
- Hilmon, J.B.** (1959). Determination of herbage weight by double-sampling : Weight estimate and actual weight. In : *Techniques and Methods of Measuring Understorey Vegetation*. Pp 20 – 25. Proc. Symp. at Tifton, Georgia, Oct. 1958. 174 pp.
- Hodgson, J.** (1979). Nomenclature and definitions in grazing studies. *Grass and Forage Science* 34: 11 – 18.

- Holechek, J.L. Pieper, R.D. and Herbel C.H.** (1989). Range management principles and practices. Prentice Hall, New Jersey – U.S.A. 501 pp.
- Holm, A.McR and Payn, A. L.** (1980). Effect of mineral supplementation on the performance of steers grazing birdwood grass (*Cenchrus setigerus*) pastures in north western Australia. Aust. J. Exp. Agric. Anim. Husb. 20: 298-405.
- Holm, A.McR and Payn, A. L, Morgan, P.D. and Speijers, E.J.** (1981). The response of weaner cattle grazing natural pastures in north western Australia to phosphoric acid, non protein nitrogen and sulphur in their drinking water. Aust. Rangel. J. 3 (2) : 133-141.
- Holm, A.McR, Curry, P.J. and Wallace J.F.** (1984). Observer differences in transect counts, cover estimates and plant size measurement on range monitoring sites in an arid shrubland. Aust. Rangel. J. 6 (2) : 28 – 102.
- Hunting Technical Services, Ltd. (1974).** Southern Darfur land use Planning survey. Annex 3: Animal resource and range ecology. Khartoum. Min. of Agric., Food and Nat. Res. 180 p.
- Igaimi, O.** (1999). White Nile State Encyclopedia, unpublished report.
- Jones, R.M. and Hargreaves, J.N.G.** (1979). Improvements to the dry weigh rank method for measuring botanical composition. Grass and Forage Sci. 14 : 181 – 189.
- Johnson, A.** (1957). A comparison of the line interception, vertical point quadrat and loop methods as used in measuring basal area of grassland vegetation. Can. J. Pl. Sci. 37 : 34 – 42.
- Kelly, R.D. and McNeil, L.** (1980). Tests of two methods for determining herbaceous yield and botanical composition. Proc. Grassld. Soc. Sth. Afr. 15 : 167 – 171.

- Kinsinger, F.E., R.E. Eckert and P.O. Currie.** (1960). A comparison of the line-interception, variable-plot and loop methods as used to measure shrub-crown cover. *J. Range Mangt.* 13: 17 – 21.
- Le Houerou, H.N. and Hoste, C. H.** (1977). Rangeland production and Annual rainfall relations in the Mediterranean Basin and in African Sahelo-Sudanian zone. *J. Range Manage.* 30(3): 181 – 189.
- Leigh, J.H. and Mulham W.F.** (1966). Selection of diet by sheep grazing semi-arid pastures on the Riverine Plain.
1. A bladder saltbush (*Atriplex vesicaria*) – cottonbush (*Kochia aphylla*) community. *Aust. J. Exp. Agric. Anim. Husb.* 6 : 468 – 474.
- Levy, E.B. and Madden, E.A.** (1933). The point method of pasture analysis. *N.Z. J. Agric.* 46 – 47 : 267 – 279.
- Lodge, G.M. and Gleeson, A.C** (1979). The effect of sample size and plot stratification on the precision of the wheel-point method of estimating botanical composition in clustered plant communities. *Aust. Rangel. J.* 1 (4) : 346 – 350.
- Lodge, G.M. and Gleeson, A.C** (1982). The importance of plant density, plant basal area and plant mass per unit basal area as factors influencing the herbage mass of some native perennial grasses. *Aust. Rangel. J.* 4 (2) : 61 – 66.
- Mannetje, L. t. and Haydock, K.P.** (1963). The dry weight rank method for the botanical analysis of pastures. *J.Br. grassld. Soc.* 18: 268 – 275.
- Maxwell, T.J.** (1991). Diagnosis and improvement methods in range utilization system. Mac Cauley Land Use Research Institute, Bent land field, Roslin Midlothian, Great Britain.

- Mazaika, R. and Krausman, P.R.** (1991). Use of dry-weight rank multipliers for desert vegetation. *J. of range manage.* 44 (4): 409 – 411.
- Mentis, M.T.** (1981). Evaluation of the wheel-point and step-point methods of veld condition assessment. *Proc. Grassld. Soc. Sth. Afr.* 16: 89 – 94.
- Mohamed, D.E.R.** (2001). The management indicators of forested and open rangelands in semi-arid areas, west of Ed Dueim – White Nile State. M.Sc. Thesis, Dept. of Range Science, Sudan Univ. of Sci. & Tech. (Unpublished).
- Mohamed, G.O.E.** (2004). The effect of utilization levels on range condition at water points for proper range management. M.Sc. Thesis, Dept. of Range Science, Sudan Univ. of Sci. & Tech (Unpublished).
- Morris, J.W. and Muller, P.J.** (1970). Seasonal variation of grassland basal cover. *Proc. Grassld. Soc. Sth. Afr.* 5: 145 – 152.
- Morris, M.J., Johnson, K.L. and Neal, D.L.** (1976). Sampling shrub ranges with a electronic capacitance instrument. *J. Range Manage.* 29: 78 – 81.
- Mueller-Dombois, D. and Ellenberg, H.** (1974). *Aims and Methods of Vegetation Ecology.* J. Wiley and sons, N.Y., 547 pp.
- Mukhtar, H.K. and Fadlalla, B.** (1987). Effect of watering interval on desert sheep growth during the summer in Northern Kordofan (Unpublished results).
- Mukhtar, M.E. and Yousif, M.A.** (1995). Environmental impact assessment in Ed Dueim province – phase I report.
- Musnad, H.A. and Abdul mutalab, H.** (1998). Forest utilization as a fodder source. *Social forestry and environment J.* – 5th ed., pp 10 –15.

- National Academy of Sciences.** (1962). Basic Problems and Techniques in Range Research. National Academy of Sciences, Publ. No. 890, Washington, D.C. 342 pp.
- Neal, D.I. and Neal J. L.** (1973). Uses and capabilities of electronic capacitance instruments for estimating herbage. Part I: history and development. *J. Br. Grassl. Soc.* 28: 81 –89.
- Neuteboom, J.H., Lantinga, E.A. and Struik, P.C.** (1998). Evaluation of the dry weight rank method for botanical composition analysis of grassland by means of simulation. *Netherlands J. Agrical. Sci.* 46: 285 – 304.
- Noble, J.C. and Totterdell, C.J.** (1978). Assessment of ground photography for sampling an *Atriplex vesicaria* Hew. ex Benth. (Bladder saltbush) community. *Aust. Rangl. J.* 1: 191 – 193.
- Opstrup, R.A.** (1968). SYSTEMS ANALYSIS OF TWO OLD-FIELD ECOSYSTEMS. M.Sc. Thesis, Univ. of Tennessee. Knoxville. 164 pp.
- Owen, D.F. and Wiegert, R.G.** (1973). Do consumers maximize plant fitness? *Oikos* 27: 488 – 492 (cited by Rutherford, M.C. 1979 plant based techniques for determining available browse and browse utilization: a review. *The pot.* 45: 203 – 228).
- Parker, K.W.** (1950). Report on 3-step method for measuring condition and trend of forest ranges. U.S. forest service. Washington D.C.
- Parker, K.W. and Harris, R.W.** (1959). The 3-step method for measuring condition and trend of forest ranges : a resume of its history, development and use. In: Techniques and methods of measuring understory vegetation. Proc. of a symposium at Tifton, Georgia.

- Payne, G.F.** (1974). Cover-weight relationships. *J. Range Mangt.* 27: 403 – 404.
- Payne, A.L., Kubicky, A. and Wilcox, D.G.** (1974). Range conditions guides for West Kimberley Area. Dept. of Agric. W. Aust. Publ.
- Payne, A.L., Mitchell, A.A. and Holman, W.F.** (1982). An Inventory Survey of Rangelands in the Ashburton River Catchment, Western Australia (1975). Tech. Bull. 62 – Dept. of Agric. W.A.
- Pechanec, J.F. and Pickford, G.D.** (1937). A weight-estimate method for the determination of range or pasture production. *J. Of the Amer. Society for Agronomy* 29: 894 – 904.
- Plan International Sudan** (1996). Environmental impact assessment – Water supply project Ed Dueim province (main report – EDS)
- Range and pasture administration office** (1993). The role of the agricultural sector in Ed Dueim province development. W.N. State.
- Rees, M.C., Jones, R.M. and Roe, R.** (1976). Evaluation of pasture grasses and legumes grown in mixtures in South-East Queensland. *Trop. Grasslds.* 10: 65 – 78.
- Ridder, N. de, Stroosyder, L., Cisse, A.M. and Keulen, H.** (1982). Productivity of Sahelian Rangeland: Study of the soils, the vegetation and the exploitation of that natural resource. Wageningen Agricultural University, the Netherlands.
- Risser, P.G.** (1984). Methods for inventory and monitoring of vegetation, litter and soil surface condition. In : *Developing Strategies for Rangeland Management*. Ed. National Academy of Sciences, NAS, pp. 647 – 690. Westview press. Boulder and London.
- Rittenhouse, L.R. and Sneva, F.A.** (1977). A technique for estimating big sagebrush production. *J. Range manage.* 30: 68 – 70.

- Rutherford, M.C. (1979).** Plant based techniques for determining Available browse and browse utilization: a review. *The Pot.* 45: 203 – 228.
- Sandland, R.L., Alexander, J.C. and Haydock, K.P. (1982).** A statistical assessment of the dry weight rank method of pasture sampling. *Grass and Forage Sci.* 37: 263 – 273.
- Sankari, M.N. (1996).** Methods of measuring density and species composition of vegetation. In: training session in modern criteria for range and pasture development pp. 117 – 156 (Arabic).
- Schumacher, F.X. and Chapman, R.A. (1948).** Sampling methods in forestry and range management. Duke Univ. School of For. Bull. No. 7.
- Segelquist, C.A. and Permington, (1966).** Deer browse in the Quachila forest in Oklahoma. *J. Wildl. Manage.* 32: 623 – 626.
- Segelquist, C.A. and Permington, (1972).** Browse resources of the Quachila national forest in Arkansas. *For. Serv. Res. Note* 50 – 140. 4 pp.
- Shaw, N.H. and Mannetje, L.'t. (1970).** Studies in speargrass pastures in central coastal Queensland – The effect of fertilizer, stocking rate and oversowing with *Stylosanthes humilis* on beef production and botanical composition. *Trop. Grasslds.* 4 (1): 43 – 56.
- Shepherd W.O. (1968).** Range and Pasture Management. Report to The government of the Sudan. Rep. Ta. No. 2468.FAO, Rome 49 p.
- Skinner, T.E. (1976).** 2. A comparison between the effects of continuous grazing by Angora goats and Merino sheep on veld in the central lower Karroo. *Proc. Grassld. Soc. Sth. Afr.* 11: 131 – 134.
- Smith, E.L. and Despain, D.W. (1991).** The Dry Weight Rank method of estimating plant species composition. In: G.B. Ruyle (ed), Some methods for monitoring rangelands and other natural area vegetation. Uni. of Ariz.,

college of Agric., Extn. Report 9043. pp 27 – 48.

- Squires, V. R.** (1976). Walking, watering and grazing behavior of Merino sheep on two semi-arid rangelands in South-West New South Wales. *Aust. Rangel. J.* 1: 13 – 23.
- Stewart, J.H.** (1960). Land and water resources of Tripolitania. Washington, U.S. Agr. Mission to Libya. USAID 248 p.
- Stoddard, L.A, Smith, A.D. and Box, T.W.** (1975). Range Management, New York : Mc. Graw Hill.
- Tilfer, E.S. and Crains, A.** (1978). Stem breakage by Moose. *J. wildl. manage.* 42: 636 – 642.
- Tidmarsh, C.E.M. and Havenga, C.M.** (1955). The wheel-point method of survey and measurement of semi-open grasslands and Karroo vegetation in South Africa. *Botl. Surv. Sth. Afr., Memo* No. 7.
- Tothill, J.C., Hargreaves, J.N.G. and Jones, R.M.** (1978). BOTANAL – A comprehensive sampling and computer procedure for estimating pasture yield and composition. 1. Field Sampling. CSIRO Australia Div. Trop. Crops and Pastures, Trop. Agron. Tech. memo No. 8.
- Trumble, H.C. and Woodroffe, K.** (1954). Influence of climatic factors in the reaction of desert shrubs to grazing by sheep. In: “Biology of deserts”, pp. 129 – 147. Institute of Biology, London.
- Van Dyne, G.M., Beldsoe, L.J., Gustafson, J.D., Hughes, J.H. and Swift, D.M.** (1975). A study of the Dry Weight Rank Method of Botanical Analysis. IBP Grassld. Biome Tech. Report No. 297. Natural resource ecology laboratory, Colorado State Univ.
- Ulritch, J.** (1995). Analysis of participatory approaches to natural resource management in the Sudan.
- Uresk, D.W., Gilbert, R.O. and Rickford W.H.** (1977) sampling big sagebush for phytomass. *J. Range manage.* 30: 311 – 314.

- Von Broembsen, H.H.** (1965). A wheel-point apparatus for survey and measurement of open and semi-open savannah vegetation. Proc. 9th Internl. Grassld. Congr. 2: 1345 – 1348. Sao Paulo, Brazil.
- Walker, B.H.** (1970). An evaluation of eight methods of botanical analysis of grasslands in Rhodesia. J. App. Ecol. 7:403 – 416.
- Whalley, R.D.B. and Hardy, M.B.** (2000). Measuring botanical composition of grasslands. In: CAB International 2001. Field and laboratory methods for grassland and animal production research (Eds. L't Mannelje and R.M. Jones).
- Wilm, H.G., Costello, D.F. and Kipple, G.E.** (1944). Estimating forage by double-sampling method. J. Amer. Soc. Agron.36: 194 – 203.
- Willms, W., Mclean, A. and Ritcey, R.** (1976). Feeding habits of mule deer on fall, winter and spring ranges near Kamloops. British Columbia. Can. J. Anim. Sci. 56 : 531 – 542.
- Wilson, A.D.** (1982). Range assessment. In: land resource management divisional report, pp. 17 – 26.
- Woodroffe, K.** (1941). Shrub pastures under low rainfall: an experiment on grazing management. J. Aust. Inst. Agric. Sci. 7: 117 – 121.
- Zahran, B.B.H.A.** (1982). The changes in nutritive requirements due to changes in livestock population in Southern Darfur – Sudan. Preliminary M.Sc. in Natural Resource Management. Univ. of Western Australia (Unpublished).
- Zahran, B.B.H.A.** (1986). FORAGE COMPOSITION AND PRODUCTION OF ANNUAL AND BIENNIAL SPECIES ON CONTRASTING SOILS IN RANGELAND GRAZED BY SHEEP NEAR CARNARVON, WESTERN AUSTRALIA. M.Sc Thesis in Agric., Agronomy group, School of Agric., Univ. of W. Australia.

(Unpublished).

Zaroug, M.G. (1993). Rangeland improvement and development in the African Horn Countries and its role in combating desertification, pp. 328 – 363 (Arabic).

Zhou, Q. and Robson, M. (1998). Comparing methods for quantitative estimation of rangeland vegetation cover. In: People and Rangelands: Building the Future, Eldridge, D. and Freudenberger, D. (eds), VI Int. Rangel. Congress, Inc, Aitkenvale, Australia: 750 – 752.

Zorich, P.R. (1966). EVALUATION OF SPECIES COMPOSITION BY FOUR METHODS ON TWO PERENNIAL GRASS PASTURES (*Festuca arundinacea*) Schreb.and *Lolium perenna* L.) GRAZED LIGHTLY AND HEAVILY IN WESTERN OREGON. M.Sc. Thesis, Oregon State Univ., Corvallis, 87 pp (unpublished).

Web Sites

- [http:// Rangelandswest.org/az/inventorymonitoring/dryweight.html](http://Rangelandswest.org/az/inventorymonitoring/dryweight.html).
Arizona Rangelands: Inventory and Monitoring: Dry-Weight Rank Method.
- crop.scijournals.org/Hodgson.j.1979/ Nomenclature and definitions in range studies.