

REFERENCES

- Attwill, (1994).** Water river watershed land management plan 2003-2012. <http://www.state.ma.us/mdc/WRLMPch10.p161>.
- Aucamp, A. J., Smith, D.W., Howe, L.G., and B. D. Viljoen. (1973).** 'n Belowende tegniek vir die bepaling van struikbenutting. Hand. Weidingsveren. S. Afr. 8: 129-132pp.
- Austin, M.P. (1981).** Use of ordination and other multivariate descriptive methods to study succession. *Vegetation*.35:165-175pp.
- B.L.M.. (1996).** Sampling Vegetation Attributes. Interagent Technical Reference, BLM/RS/ST-96/002+1730 28-29 pp.
- Ball, D. A., and Miller, S. D. (1989).** A comparison of techniques for estimation of arable soil seed bank and their relationship to weed flora. *Weed Res.* 29:365-373pp.
- Barbour, M.G., Burk, J.H., and W.D. Pitts. (1987).** Terrestrial plant ecology. Benjamins Publishing Co., Menlo Park, CA. 2nd ed. 191-193 pp.
- Barnes, D. L. (1976).** A review of plant-based methods of estimating food consumption, percentage utilization, species preferences and feeding patterns of grazing and browsing animals. *Proc. Grassl. Soc. S. Afr.* 11:65-71.
- Barralis, G. (1972).** Evolution comparative de la flore adventice avec ou sans de's herbage chimique. *Weeds Res.* 12:115-127pp.
- Basile, J. V., and S. S. Hutchings. (1966).** Twig diameter-length-weight relations of bitter-brush. *J. Range Manage.* 19:34-38pp.
- Baskin, G.G., and Baskin, J.M. (1998).** Seeds: Ecology, Biogeography and Evolution of Dormancy and Germination. Academic Press, San Diego, 666pp.
- Blair, R.M. (1971).** Forage production after grazing
- BMELF, (1994).** Waldbericht der Bundesrepublik Deutschland, Bonn,.

- Bobek, B., and R. Dzieciolowski. (1972).** Method of browse estimation in different types of forests. *J. Wildl. Manage.* 41: 677-684pp.
- Bonham, C.D. (1989).** Measurements for terrestrial vegetation. John Wiley & Son, New York, NY.90-96pp.
- Bookhout, T. A. (1965).** Feeding coactions between snowshoe hares and white-tailed deer in northern Michigan. *Trans. N. Am. Wildl. Conf.* 30:321-335pp.
- Cavers, P.B. (1983).** Seed demography. *Can.J.Bot.*61:3578-3590pp.
- Chan, S. C. (2000).** Slopeland capability classification as a tool in land use planning. A. Taiwan. www.fftc.agnet.org/library/article/eb483a.html
- Cook, C.W., and J. Stubbendieck(eds.)(1986).** Range research. Basic problems and techniques. Society for range management, Denver, CO. 51-56pp.
- Darag, A., and Suliman, M. (1988).** Training course in Range Management and Range Improvement. Range and pasture administration (Sudan).47pp.
- De Vos, A., and H.S. Mosby. (1969).** Habitat analysis and evaluation. 135-172pp. In: **R. H. Giles (ed.).** Wildlife management techniques, 3rd ed. The Wildlife Society, Washington, D. C.
- Despain, D.W., Ogden, P.R., and E.L. Smith. (1991).** Plant frequency sampling for monitoring rangelands. In: G.B.Ruyle.(ed.). Some methods for monitoring rangelands and other natural area vegetation. University of Arizona, College of Agriculture, Extension Report 9043.7-25pp.
- Dessaint, F., Chadoeuf, R., and Barralis, G. (1991).** Spatial pattern analysis of weed seeds in the cultivated soil seed bank. *Jour. Of App. Eco.* 28:721-730pp.
- Dougall, H. W., and Bogdon, A. V. (1958).** Browse plants of Kenya, *East African Agricultural* 23:236-245pp.

- Elsiddig, E.A. (1980).** The prediction of stem and branch volume of *Acacia nilotica* in the Fung Region in the Sudan.
- F.A.O. (1975).** The ecological management of the arid and semi arid rangelands in Africa and the Near Middle East. Report of F.A.O.
- F.A.O, (1993).** The Challenge of Sustainable Forest Management-what future for the world's forests.
- F.A.O, (1993).** FAO development series I. Guidelines for landuse planning. Rome, 1993.
- Falinska, K. (1999).** Seed bank dynamics in abandoned meadows during a 20-year period in the Bialowieza National Park. *Journal of Ecology* 87:461-475pp.
- Fay, R. K., and Olson, W. A. (1978).** Techniques for separating weed seed from soil. *Weed Science* 26: 530-533pp.
- Fenner, M. (1985).** Seed ecology. Chapman and Hall, London.
- FPCS (1995).** Forest Products Consumption Survey. Sudan. FNC.
- FRA (2000).** Global Forest Resources Assessment. The situation and developments in the forest sector. Part I.
- Fries, J. (1990).** Natural forest management in semi-arid Africa status and research needs. The Sahel.
- Gaiballa, A.K. (1995).** The effect of utilization on seed bank and vegetation dynamic of indigenous plant species including selected wild food in Rawashda forest in Eastern Sudan. PhD (thesis) University of Khartoum, Sudan.
- Grandian, U., and H. Rydin (1998).** Attributes of the seed bank after a century of primary succession on islands in Lake Hjalmar, Sweden. *J. of Ecology* 86:922-933pp.
- Gray, S.G. (1970).** The place of trees and shrubs as sources of forage in tropical and subtropical pastures. *Trop.Grassl.*4:57-62pp.

- Gray, S. G. (1970).** The place of trees and shrubs as source of forage in tropical and subtropical pasture. *Trop. Grassl.* 4:37-62pp.
- Halls, L.K. and R.F. Harlow. (1971).** Weight-length relations in flowering dogwood twigs. *J. Range Manage.* 24: 236-237pp.
- Harper, J.L. (1977).** Population Biology of plants. 437-444pp. Academic press.
- Hassan, R. M. (2000).** Depletion of forest resources in Sudan: Intervention options for optimal control. University of Pretoria.
- Heermans, J. (1991).** Forests, trees and people Newsl., No.11.
- Holechek, J.L., Pieper, R.D., and C.H. Herbel. (1995).** Range management principles and practices. Prentice Hall, Englewood Cliffs, N.J. 2nd ed. 135-136pp.
- Holechek, J.L., Pieper, R.D., and Herbel, C.H. (1989).** Range Management Principles and Practices. Prentice Hall New Jersey 501pp.
- Hyatt, A., Casper, B.B. (2000).** Seed bank formation during early secondary succession in a temperate deciduous forest. *Journal of Ecology* 88: 516-527pp.
- Hyatt, L. (1999).** Differences between seed bank composition and field recruitment in a temperate zone deciduous forest. *The American Midland Naturalist.*142: 31-38pp.
- ICA (1959).** Hand book on range management. Revision of the regional ICA. Regional Range Management Course Book of 1955 for the Near South Asia Area.
- I.L.R.I. (1998).** Linking livestock and natural resource management. International Livestock Research Institute, Nairobi Kenya, Addis Ababa Ethiopia.
- ITTO' (2000).** Review of progress towards the year 2000 objectives for sustainable forest management.

- Jameson, D.A. (1963).** Responses of individual plants to harvesting. *Bot.Rev.*29:532-594pp.
- Jankowska-Blaszczuk, M., and Grubb, P.J. (1997).** Soil seed banks in primary and secondary deciduous forest in Bialowieza, Poland. *Seed Science Research* 7: 281-292pp.
- John, K. (1997).** Approaches to ecologically based forest management, partI. www.na.fs.fed.us/spfo/pubs/misc/ecoforest/p1.htm.
- John, M. K. (2004).** Ecosystem approaches and sustainable forest management. Discussion paper for the UNFF Secretariat.
- Johnston, S. K., Crowley, R. H., and Murray, P. S. (1978).** Separating seed by species with Ca Cl₂ solution. *Weed Sci.* 26(3):213-215pp.
- Joyal, R. (1976).** Winter foods of moose in La Verendrye Park, Quebec: an evaluation of two browse survey methods. *Can. J. Zool.* 54:1765-1770pp.
- Kelly, R. D. (1977).** The significance of the woody components of semi-arid savanna vegetation in relation to meat production. *Proc. Grassl. Soc. S. Afr.* 12: 105-108pp.
- King, D.R. (1975).** Estimation of yew and cedar availability and utilization. *J. Wildl. Manage.* 39: 101-107pp.
- Koniak, and Evertt (1982).** Seed bank dynamics in hayfield succession. *J.Ecology* 88.
- Knierim, P.G., K. L. Carvell, and J. D. Gill. (1971).** Browse in thinned oak and cove hardwood stands. *J.Wildl. Manage.* 35: 136-168pp.
- Korpac, Z. (1966).** Estimation of weed seed in arable soil. *Pedobiological* 6:105-128pp.
- Le Houerou, H. N. (1980).** Browse in Africa. The current state knowledge Addis Ababa: I.L.C.A.

- Leck, M.A. (1989).** Wetland seed banks. In: Leck, M.A., Parker, V.T., and Simpson, R.L.(eds.). Ecology of Soil Seed Banks. Academic Press, San Diego, 283-305pp.
- Lyon, L.J. (1968).** Estimating twig production of serviceberry from crown volumes. *J. Wildl. Manage.* 32:115-119pp.
- Major, H.E. and Pott, W.T. (1966).** Buried viable seed in two California bunch grass sites and their bearing on the definition of a flora. *Vegetation*, 13:253-282pp.
- Malone, C. R. (1976).** A rapid method for estimation of viable seed in the soil. *Weeds Res.* 15:381-382pp.
- Mark, S. and Martha, R. (2000).** Effects of grazing and cultivation on forest plant communities in Mount Elgon National Park, Uganda. East African Wild Life Society. *Afr. J. Ecol.* 38:154-162pp.
- Martin, M.H., Cox, J.R., and F.F. Ibarra. (1995).** Climate effects on buffelgrass productivity, in the Sonoran Desert. *Journal of range management* 48:60-63.
- Maxwell, T.J. (1991).** Diagnosis and improvement methods in range utilization system. Maccauely Land Use Research Institute, Bent land field, Roslin Midlothian, Great Britain.
- Maybury, k. (1999).** Seeing the forest and the trees. Ecological classification for conservation. The Nature Conservancy. Arlington, Verginia.
- McKell, M. (1980).** Multiple use of fodder trees and shrubs. A world wide perspective . 141-149pp.
- Medwecka-Kornas, A. (1967).** Estimation of primary production as a basis for studies of secondary production. 83-95pp. In: **K. Petruszewicz (ed.).** Secondary productivity of terrestrial ecosystems. Vol. 1. Krakow, Warsawa.

Michael, S.P. (1994). Measuring trees and forests. Cab International Press.

Mike (2001).

Miles, J. (1979). Vegetation dynamics. Chapman and Hall London.

Moller, P.F. (2000). The Danish strategy for natural forest- Background, realization and perspectives. University of Tartu, Estland.6-12pp.

Mueller-Dombois, D. and Ellenberg, H. (1974). Aims and methods of vegetation ecology. J.Wiley and Sons, N.Y., 547pp.

Nakagoshi, N. (1985). The population structure of vegetation. Ed. White, J. 551-570pp.

Niamir, M. (1990). Herds decision-making in natural resource management in arid and semi arid Africa F.A.O. 126pp.

O'Connor, T. G., and Pickett, G. A. (1992). The influence of grazing on seed production and seed bank of some African savanna grasses. *J. app. Ecol.* 29:247-260pp.

Owen, D. F., and R.G. Wiegert. (1976). Do consumers maximize plant fitness? *Oikos* 27: 488-492pp.

P.T. Tueller. (ed.). Vegetation science application for rangeland analysis and management. Handbook of vegetation science, Volume 14. Kluwer Academic Publishers, Dordrecht.449-467pp.

Peek, J.M. (1963). Appraisal of a moose range in southwestern Montana. *J. Range Manage.* 16: 227-231pp.

Peter, F. Kolb.(2002). The future of Montana's forests. Crafting a vision for landowners and managers of Montana's forests. www.mttreefarm.org/pdf-files/graze-pdf.

Pickett, S.T., and McDonnell, M.J. (1989). Physiology of dormancy and germination in relation to seed bank ecology. Ecology of soil seed banks. Academic Press, San Diego.123-147pp.

Pieper, R.D. (1988). Rangeland vegetation productivity and biomass. In:

Pott, A., and Humphreys, L.R. (1983). Persistence and growth of *Lotononis bainesii*, *Digitaria decumbens* pasture. 1- Sheep stocking rate. *J. Agric.* 101:1-7pp.

Plan International, (1996). Environmental impact assessment. Water supply project. Main report. Environment and Development Services (EDS).

Priestely, D.A. (1986). Seed aging: Implications for seed storage and persistence in the soil. Ithaca, NY: Cornell university press.

Rees, M. (1996). Delayed germination of seeds: a look at the effects of adult longevity, the timing of reproduction, and population age/stage structure. *American Naturalist* 144, 43-64pp.

Ridder, N. (1982). Productivity of Sahelian Range Land, study of the soil, the vegetation and the exploitation of the natural resources. Wageningen Agricultural University. Netherlands.

Roberts, H. G. (1981). Seed bank in the soil *Advances in applied biology.* 6:2-6pp. Academic Press.

Robert, H.A. and Herson, R.T. (1971). Herbicide performance and soil surface condition. *Weed Research,* 11:69-73pp.

Rutherford, M.C. (1979). Plant-based techniques for determining available browse and browse utilization: A review 45: 203-228pp.

Segelquist, C.A., and R.E. Pennington. (1968). Deer browse in the Ouachita forest in Oklahoma. *J. Wildl. Manage.* 32:623-626pp.

Segelquist, C. A., and R. E. Pennington. (1972). Browse resources of the Ouachita National forest in Arkansas. *For. Serv. Res. Note* SO-140. 4pp.

SFNSW, (2003). State Forest of New South Wales, Australia, 2003. Forest facts: Forest Types and Vegetation Patterns in New South Wales.

Shafer, E. L. (1963). The twig-count method for measuring hardwood deer browse. *J. Wildl. Manage.* 27: 428-437pp.

Silvertown, J. W. (1987). Introduction to plant population ecology. Longman London. 198pp.

Skvo-og Naturstyrelsen, (1994). Strategy for Natural Forests and Other Forest Types of High Conservation Value in Denmark. Ministry of the Environment 48pp.

Stoddart, L.A., Smith, A.D., and Box, T.W. (1975). Range Management. New York: Mc Graw Hill.

Telfer, E. S., and A. Cairns. (1978). Stem breakage by moose. *J. Wildl. Manage.* 42:639-642pp.

Telfer, E. S. (1974). Vertical distribution of cervid and snowshoe hare browsing. *J. Wildl. Manage.* 38: 944-946pp.

Thompson, K. and Grime, J.P. (1979). Seasonal variation in the seed banks of herbaceous species in the contrasting habitats. *Jour. Of Eco.* 67:893-921pp.

Thompson, K., Bakker, J.P., and Bekker, R.M. (1997). The Soil Seed Banks of North West Europe: Cambridge University Press, Cambridge, 276pp.

U.N.C.E.D, (1992). Earth Summit-Rio Declaration & Forest Principles.

Vallantine, J.F., and Philip, L.S. (1980). Range Science. Guide to information source. Gale Res. Co., Detroit, Michigan 32 I pp.

Widstrand, G. C. (1975). The rational of nomads economy. *Ambio* 4(4): 146-153pp.

Wijngaarden, W. Van. (1985). Elephants-Trees-Grass-Grazers: Relationship between climate, soils, vegetation and large herbivores in a semi-arid ecosystem (Tsavo, Kenya) ITC Publ No 4 Enschede, Netherlands.

Willie, M. L., P.Hologram, and F.Castaneda (2003). Sustainable forest management and the Ecosystem Approach:Two concepts one goal. FAO working paper FM25, Rome, Italy.

Willms, W., McLean, A., and R. Ritcey. (1976). Feeding habits of mule deer on fall, winter and spring ranges near Kamloops, British Columbia. *Can. J. Anim. Sci.* 56:531-542pp.

WORLD BANK (2002). World Bank Forest Policy and Strategy.

Wormald, T. J. (1984). The management of the natural forests in the arid and semi-arid zones of East and southern Africa (a report for O.D.A). 60pp.