

References

- 1) Addison GM, Beamish MR, Hales CN. (1972). An Immuno-radiometric Assay for Ferritin in the Serum of Normal Subjects in Patients with Iron Deficiency and Iron Overload. *J Clin Pathol*; 25: 326-9.
- 2) Allen LH. (1997). Pregnancy and Iron Deficiency: Unresolved Issues. *Nutrition Reviews*; 55: 91-101.
- 3) Andrews NC. (1999). Disorders of iron metabolism. *N. Engl. J. Med.*; 341:1986 -1995.
- 4) Atukorala T, de Silva LD, Dechering WH, Dassenaieke TS, Perera RS..(1994). Evaluation of effectiveness of iron Folate supplementation and antihelminthic therapy against anaemia in pregnancy-study in the plantation sector of Sri Lanka. *Am.J.Clin.Nutr.*;60: 286 - 292.
- 5) Bates HM. (1980). How to Detect Iron Deficiency Before Anaemia Develops. *Laboratory Pathfinder* Jan:17-22.
- 6) Bothwell TH, Charlton RW, Cook JD, Finch CA. (1979). *Iron Metabolism in Man*. Oxford Blackwell.

- 7) Brecher GM, Schneiderman M and Williams GZ. (1956). Evaluation of electronic red blood cell counter. Am J Clin Path; 26:1439-1449.
- 8) British Nutrition Foundation (1995). Iron: nutritional and physiological significance. Chapman and Hall, London. p 23 - 32.
- 9) Brittin GM, Brecher G and Johnson CA. (1969). Evaluation of the Coulter Counter . Model S. AM J Clin Path; 52: 780-783.
- 10) Brock JH, Halliday JW, Pippard MJ, Powell LW(ed). (1994). Iron Metabolism in Health and Disease. London:Saunders.
- 11) Carriaga Mt, Skikne BS, Finley B. (1991). Serum transferrin receptor for the detection of iron deficiency in pregnancy. Am.J.Clin.Nutr.; 54: 1077 - 1081.
- 12) Cazzola M, Huebers HA, MacPhail P. (1985). Transferrin saturation, plasma iron turnover and transferrin uptake in normal humans. Blood; 66: 235 - 939.

- 13) Cook JD, Skikne BS, Lynch CR. (1980). Serum Ferritin in the Evaluation of Anaemia. In: Albertin A, editor. Radioimmunoassay of Hormones, Proteins and Enzymes. Amsterdam: Excerpta Medica,;239-48.
- 14) Cook JD. (1994). Iron deficiency anaemia. Bailliere's Clin Hematol.; 7: 787 - 804.
- 15) Diejomaeoh FME, Abdulaziz A, Adekile AD. (1999). Anaemia in pregnancy. Int.J.Gynerol. Obstet.;65: 299 - 301.
- 16) Eckhoff RE. (1967). An experimental indication of the volume proportional response of the Coulter Counter for irregularly shaped particles. J Sci Inst; 44: 648-649.
- 17) England JM, Walford DM and Waters DAW. (1972). Reassessment of the reliability of the haematocrit. Brit J Haemat; 23:247-256.
- 18) Fairbanks VF, Beutler E. (1977). Iron Deficiency. Haematology. New york: McGraw Hill Book Co.: 363-87.
- 19) Fairbanks VF, Fahey JL, Beutler E. (1971). Clinical Disorders of Iron Metabolism (2nd Ed.). New York: Grune and Stratton Inc.:46-54.

- 20) Ferguson BJ, Skikne BS, Simpson KM. (1992). Serum transferrin receptor distinguishes anaemia. *J.Lab.Clin. Med*; 119: 385 - 390.
- 21) Firkin F, Chesterman C, Penington D, Rush B.(ed). (1989). Pregnancy and anaemia. In: de Gruchy,s *Clinical Hamatology in Medical Practice*. Fifth ed. Blackwell Science; 113 - 115.
- 22) Forman DT, Parker SL. (1980). The Measurement and Interpretation of Serum Ferritin. *Ann Clin Lab Sci*;10: 345-50.
- 23) Gottmann NA.(1970). Multiple haematological analysis by means of a Coulter Counter Model S. Paper presented at International Symposium of Standardization of Hematological Methods.
- 24) Grover NB, Naaman J, Ben-ason S and Dojanski F. (1972). Electrical sizing of the particles in suspension III. Rigid spheroids and red blood cells. *Biophys J*;;12: 1099-1116.
- 25) Hallberg L, Bjorn-Rasmussen E. (1972). Determination of iron absorption from whole diet. A new tool model using two radiation isotopes given as haem and non-haem iron. *Scand. J.Hematol.*; 9: 193 -197.

- 26) Hamilton PJ and Davidson RL. (1973). The interrelationships and stability of Coulter S-determined blood indices. *J Clin Path*,; 16: 700-705.
- 27) Huebers HA, Beguin Y, Pootrakul P. (1990). Intact transferrin receptors in human plasma and their relation to erythropoiesis. *Blood*,; 75: 102 - 107.
- 28) Hulten LE, Gramatkovski A, Gleerup A, Hallberg L. (1995). Iron absorption from the whole diet. Relation to meal composition, iron requirement and iron stores. *Eur. J. Clin. Nutr.*; 49: 794 - 808.
- 29) Jacobs A, Miller F, Worwood M. (1972). Ferritin in the Serum of Normal Subjects and Patients with Iron Deficiency and Iron Overload. *Br Med J*;4:206-8.
- 30) Jacobs A, Path FRC, Worwood M. (1975). Ferritin in Serum: Clinical and biochemical Implications. *N Engl J Med*;292: 951-956.
- 31) Kachel V and Ruhstroth-Bauer G. (1976). Coulter. *Micros Acta*,; 75:419-423.
- 32) Kaufer M, Casanueva E.(1990).Relation of pregnancy serum ferritin levels to haemoglobin levels throughout pregnancy .*Eur J clin Nutr* ; 44: 709- 715.

- 33) Klausner R, Rouault TA, Hartford JB. (1993).Regulating the late of mRNA: the control of cellular iron metabolism. Cells.; 72: 19 - 28.
- 34) Klinik M, Yuregri GT, Ekerbicei H. (2002). Anaemia and iron deficiency anaemia in South east Anatolia . Eur J of haematology ,96:280.
- 35) Kruse JR Stolc V. (1979). Serum ferritin and bone marrow Iron Stores. Correlation with Absence of Iron in Biopsy Specimens. Am J Clin Pathol; 72:817-20.
- 36) Kuhn LC. (1994). Molecular regulation of iron proteins. Bailliere's Clin Hematol.; 7:763 - 785.
- 37) Kuiperkramer EPA, Huisman CMS, VanRaam J.(1996). Analytical and clinical implications of soluble transferrin receptors in serum. European Journal of Clinical Chemistry and Clinical Biochemistry.;34: 645 - 649.
- 38) Letsky E. (1995). Blood volume, haematinics, anaemia. In: de Swiet M.(ed). Medical Disorders in obstetric practice, 3rd ed. Oxford: Blackwell,; 33 - 60.
- 39) Lipschitz DA, Cook JD, Finch CA. (1974). A Clinical Evaluation of Serum Ferritin as an Index of Iron Stores. N Engl J Med;290: 1213-6.

- 40) Lops VR, Hunter LP, Dixon LR. (1995). Anaemia in pregnancy *Am Fam Physician*.; 51: 1189 - 1197.
- 41) Lozoff B, Jimenez E, Wolf AW.(1992).Long-term developmental outcome of infants with iron deficiency. *N Engl. J. Med.*; 325: 687 - 694.
- 42) Miles LEM, Lipschitz DA, Bierber CP, Cook JD. (1974). Measurement of Serum Ferritin by a 2-site Immunoradiometric Assay. *Anal Biochem*;61:209-24.
- 43) Milman N, Bergholt T, Byg KE, Eriksen L, Graudal N. (1999). Iron status and iron balance during pregnancy. A critical re-appraisal of iron supplementation. *Acta Obstet. Gynecol. Scand.*; 78: 749 - 757.
- 44) Milman N, IbsenKK, Christensen JM. (1987). Serum Ferritin and Iron Status In Mother And Newborn Infants. *Acta Obstet Gynecol Scand.*; 66: 205 - 211.
- 45) Murphy JF, O Riordan J. Newcombe RJ. (1986). Relation of haemoglobin levels in first and second trimesters to outcome of pregnancy. *Lancet*; 1: 992 - 995.
- 46) Narasihns RAO, Vijaysarthy C. (1975). Fortification of common salt with iron: effect on stability and bio-availability. *Am. J. Clin. Nutr.*; 26: 1395 - 1401.

- 47) Pippard MJ, Hoffbrand AV. (2001). Iron deficiency anaemia. In: hoffbrand AV, Lewis SM, Tuddenham EGD. (ed). Postgraduate Haematology 4th ed. Arnold,; 35 - 37.
- 48) Pitrof R, Johanson R. (1996). Safe motherhood-an achievable and worthwhile aim. In :Studd J. (ed) Progress in Obstetrics and Gynecology, vol. 12. Edinburgh: Churchill Livingstone,; 47 - 57.
- 49) Prasad AS. (1996). Zinc deficiency in women, infant and children. J. Am. Coll. Nutr.; 15: 113 - 120.
- 50) Prema k, Neelak S, Ramalakshmi BA. (1981). Anaemia and adverse obstetric outcome. Nutr.Rep. Int.; 23: 637 - 643.
- 51) Preziosi P, Prual A, Galan P. (1997). Effect of iron supplementation on the iron status of pregnant women: consequences for newborns. Am.J.Clin.Nutr.;66: 1178-1182.
- 52) Prual A, Galan P, Bernis DL.(1988). Evaluation of iron status in Chadian pregnant women : consequences of maternal iron deficiency in haematopoietic status of newborns .Trop .geogr. Med., 40; 1-6.

- 53) Puolakka J, Janne O, Pakarinen A, Vihko R. (1980). Serum ferritin as a measure of stores during and after normal pregnancy with and without iron supplements. *Asta Obst Gynecol Suppl.* 95: 43 - 51.
- 54) Ross AC, Gardner EM. (1994). The function of vitamin A in cellular growth and differentiation and its role during pregnancy and lactation. *Adv. Exp. Med. Biol.*; 352: 187 - 200.
- 55) Sarin AR. (1995). Severe anaemia of pregnancy, recent experience. *Int. J. Gynecol. Obstet.*; 50 (suppl. 2): 545 - 549.
- 56) Scholl TO, Hediger ML, Fisher RL, Shearer JW. (1992). Anaemia vs. iron deficiency: increased risk of preterm delivery in a prospective study. *AmJ Clin. Nutr.*; 55: 985 - 988.
- 57) Schwartz WJ, Thurnau GR. (1995). Iron deficiency anaemia in pregnancy. *Clin. Obstet. Gynecol.*; 38: 443 - 454.
- 58) Sharma JB, Arora BS, Kumars, Goel S, Dhamija A. (2001). Helminth and protozoan intestinal infections: an important cause for anaemia in pregnant women in Delhi, India. *J. Obst. Gynecol. Ind.*: 51(6): 58 - 61.

- 59) Sharma JB. (2003). Nutritional anaemia during pregnancy in non-industrial countries. In: Studd J(ed). Progress in obstetric and gynecology, Vol.15 Churchill living stone,. pp. 103 - 122.
- 60) Skikne BS, Cook JD. (1981). Serum Ferritin in the Evaluation of Iron Status. Lab Management; 19:31-40.
- 61) Sloan NL. Jordan EA, Winikoff B. (1992). Does Iron Supplementation Make a differential ?. Mother care Project, 15 Arlington, VA, USA,.
- 62) Sood SK, Madan N, Rusia U, Sharma S. (1989). Nutritional anaemia in pregnancy and its health implications with special reference to India. Ann Natl Acad Med Sci; 41 - 50.
- 63) Sood SK, Ramachandran K, Mathur M.(1975). WHO sponsored collaborative studies on nutritional anaemia in India. The effects of supplemental oral iron administration to pregnant women. QJ Med.; 44: 241 - 258.
- 64) Stoltzfus R, Dreyfuss ML. (1998).Guidelines for the use of iron supplements to prevent and treat iron deficiency anaemia. Genera: INACG, WHO, UNICEF..
- 65) Turmen T, AbouZahr C. (1994). Safe motherhood. Int .J. Gynecol. Obstet.; 46: 145 -153.

- 66) Valberg LS.(1980). Plasma ferritin concentration: their clinical significance and relevance to patient care. *Con med assoc J*, 122: 1240.
- 67) Verhoeff FH, Brabin BJ, Chimsuki L. (1999). An analysis of the determinates of anaemia in pregnant women in rural Malawi_ abasis for action. *Ann. Trop. Med. Parasitol.*;93: 119 - 133.
- 68) Viteri FE, Alvarez E, Batres R. (1995). Fortification of sugar with NaFe EDTA improves iron status in semi-rural populations in Guatemala. *Am.J.Clin.Nutr*; 61: 1153 - 1163.
- 69) Viteri FE. (1994). The consequences of iron deficiency and anaemia in pregnancy. *Adv. Exp. Med. Biol.*; 352: 127 - 139.
- 70) Walters GO, Miller FM, Worwood M. (1973). Serum Ferritin Concentration and Iron Stores in Normal Subjects. *J Clin Pathol*; 26:770-2.
- 71) Waterman CS, Atkinson EE, Wilkins B. (1975). Improved measurement of erythrocyte volume distribution by aperture-counter signal analysis. *Clin Chem*,; 21: 1201-1211.
- 72) WHO/FRH/ MSM/96.1. Geneva: WHO, 1996.

- 73) WHO/UNICEF/UNU. (1996). Indicators for Assessing Iron Deficiency and Strategies for its prevention: WHO draft. Geneva.
- 74) World Health Organization. (1972). Report of a WHO Group of Experts on Nutritional Anaemia. Technical report series no.503. Geneva.
- 75) World Health Organization. Revised (1990). Estimates of Maternal Mortality.
- 76) World Health Organization. (1992). The Prevalence of Anaemia in women: A Tabulation of Available Information. Geneva.
- 77) World Health Organization. (1997). WHO Global Database. Geneva.
- 78) Worwood M. (2001). Serum iron concentrations in health and disease. In: Lewis SM, Bain BJ, Bates I.(ed). Dacie and Lewis Practical Haematology. 9thed.Churchill Livingstone.; 121 - 123.