

## الآية

قال تعالى:

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

قُلْ لَوْ كَانَ الْبَحْرُ مَدَادًا لِكَلَّا حَلَتْ رَبِيْ لَنِفَدَ الْبَحْرُ قَبْلَ أَنْ تَنْفَدَ كُلُّ مَاءٍ

(رَبِيْ وَلَوْ جَهْنَمْ بِمُثْلِهِ مَلَدًا ﴿١٠٩﴾)

صدق الله العظيم

سورة الكهف الآية (109)

## ***Dedication***

***To my beloved and blessed parent who did  
everything for me.....***

***To my dear all members of my family who spend  
more time encourage me.....***

***To our supervisor who stands with us step by  
step***

***To our colleague and our greatly lovely friends  
for their kindness encouragement.....***

Soaad

Sahar

Duaa

## **Acknowledgements**

First of all we would like to thank **Allah** for providing us the strength and ambition to fulfill this study.

And we would like to thank **our supervisor Mohamed Masaad Ahmed** for his support and endless valuable advice.

Our thanks are also extended to all staff member of the microbiology department for their help where the work has done.

Our poor patient to everyone who provided us with information, ideas and support.

## **Abstract**

This study had been carried out in Khartoum state during period February 2014 to April 2014. The study designed to *in vitro* evaluation of some group of antibiotics on *pseudomonas aeruginosa*, which were isolated from urine, wound swab, bed sore, sputum, blood and meningal swab.

Sixty clinical isolates were collected from Khartoum Teaching, Educational Hospital, Omdurman Military and Royal Care hospitals. The isolated bacteria were confirmed by biochemical tests and special media and then the sensitivity testing was done. *Pseudomonas aeruginosa* isolates were recovered from wound swab (48.3%), urine (36.7%), bed sore swab (6.7%), blood (5%), sputum (1.7%) and meningeal swab (1.7%).

The resistant rate of *pseudomonas aeruginosa* to Imipenem ,Meropenem, Colistin ,Streptomycin ,Amikacin ,Cefotaxime were(0%,26.7%,10%,41.7% 6.7%,95%) respectively.

## **ملخص الاطروحة**

تم تنفيذ هذه الدراسة في الفتره ما بين فبراير وابريل 2014 ،تناولت الدراسة التقييم الخارجي لبعض المضادات الحيوية على الزائفة الزنجارية المعزولة من (البول، الجروح، القروح السريرية، البلغم، الدم وغشاء المخ).

تم جمع 60 وسط غذائي لزائفة الزنجارية من المستشفيات التالية (مستشفى الخرطوم، مستشفى سوبا، مستشفى السلاح الطبي ومستشفى رويدا كير). تم التأكيد من هذه البكتيريا بإجراء المزيد من الاختبارات الخاصة بها ومن ثم إجراء اختبارات الحساسية لها.

مثلت الزائفة الزنجارية المعزولة من العينات التالية الجروح (48.3%)، البول (36.7%)، القروح السريرية (6.7%)، الدم (5%) والبلغم (1.7%) والمسحه الدماغيه (1.7%).

النتائج اشارات الى معدل المقاومة للامبينيم، الميروبينيم، الكوليستين، الاستريبيتو مايسين، الامبيكاسين والسيفوتاكسيم ( 95%, 6.7%, 10%, 26.7%, 0% ) على التوالي.

# Contents

<b>Title</b>	<b>Page</b>
الأدبية	I
Dedication	II
Acknowledgments	III
Abstract	IV
ملخص الأطروحة	V
Contents	VI
List of tables	IX
List of graphs	IX
List of color plates	IX

<b>Chapter One</b>	
1. Introduction	1
1.1 <i>Pseudomonas</i>	1
1.2 Objectives	2
Main objective	2
Specific objectives	2
<b>Chapter Two</b>	
2. Literature review	3
2.1 <i>Pseudomonas aeruginosa</i>	3
2.1.1 Definition	3
2.1.2 Normal habitat	3
2.1.3 Viability	3
2.1.4 Cell wall component	3
2.1.5 Antigenic structure	4
2.1.6 Virulence factors	4
2.1.7 Pathogenicity	5
2.1.7.1 Respiratory tract infection	5
2.1.7.2 Skin infection	6
2.1.7.3 Urinary tract infection	6
2.1.7.4 Ear infection	6

2.1.7.5 Eye infection	6
2.1.7.6 Endocarditis	7
2.1.7.7 Nosocomial infection	7
2.1.7.8 Cystic fibrosis	7
2.1.8 Laboratory diagnosis	8
2.1.8.1 Microscopy	8
2.1.8.2 Culture	8
2.1.8.3 Pigments	8
2.1.8.4 Set of sugars	9
2.1.8.5 Biochemical tests	9
2.1.8.6 Epidemiological typing methods	9
2.1.8.6.1 Serotyping	9
2.1.8.6.2 Pyocin typing	10
2.1.8.6.3 Genotyping	10
2.1.9 Treatment	10
2.2 Antimicrobials agents	12
2.2.1 Definition	12
2.2.2 Cephalosporins	12
2.2.2.1 Chemistry	12
2.2.2.2 Classification	12
2.2.2.3 Spectrum of activity	13
2.2.2.4 Toxicity	13
2.2.3 Aminoglycosides	13
2.2.3.1 Mechanism of action	13
2.2.3.2 Spectrum of activity	14
2.2.3.3 Toxicity	15
2.2.4 Polymyxin (Colistin)	15
2.2.4.1 Mechanism of action	15
2.2.4.2 Spectrum of activity	15
2.2.4.3 Toxicity	15
2.2.5 Carbapenems	15
2.2.5.1 Mechanism of action	15
2.2.5.2 Spectrum of activity	16
<b>Chapter Three</b>	
3. Materials and Methods	17
3.1 Study area	17

3.2 Study population	17
3.3 Study location	17
3.4 Study duration	17
3.5 Sample size	17
3.6 Data collection	17
3.7 Materials	17
3.7.1 Media	17
3.7.2 Glass wares	18
3.7.3 Other materials	18
3.8 Methods	18
3.8.1 Subculture	18
3.8.2 Identification	18
3.8.2.1 Colonial morphology	18
3.8.2.2 Grams stain	18
3.8.3 Biochemical tests	19
3.8.3.1 Oxidase test	19
3.8.3.2 Citrate utilization test	19
3.8.3.3 Indole test	19
3.8.3.4 Urea hydrolysis test	20
3.8.3.5 Kilger's Iron Agar	20
3.8.4 Antimicrobial sensitivity testing	20
3.8.4.1 Preparation of inoculums	20
3.8.4.2 Seeding of media	20
3.8.4.3 Application of antimicrobial disc	21
3.8.4.4 Interpretation of the zone size	21
3.8.5 Quality control	21

## **Chapter Four**

4 Results	22
-----------	----

## **Chapter Five**

5. Discussion, Conclusion and Recommendations	31
5.1 Discussion	31
5.2 Conclusion and Recommendations	32
5.2.1 Conclusion	32
5.2.2 Recommendations	32
References	34
Appendix	37

## List of tables

<b>Table</b>	<b>Page</b>
Table(1) distribution of <i>Pseudomonas aeruginosa</i> among clinical isolates	23
Table (2) susceptibility of <i>Pseudomonas aeruginosa</i> to different antibiotics	23

## List of graphs

<b>Graph</b>	<b>Page</b>
Graph (1) distribution of <i>Pseudomonas aeruginosa</i> among clinical isolates	24
Graph (2) percentage of Imipenem resistant <i>Pseudomonas aeruginosa</i>	24
Graph (3) percentage of Colistin resistant <i>Pseudomonas aeruginosa</i>	25
Graph (4) percentage of Cefotaxime resistant <i>Pseudomonas aeruginosa</i>	25
Graph (5) percentage of Streptomycin resistant <i>Pseudomonas aeruginosa</i>	26
Graph (6) percentage of Meropenem resistant <i>Pseudomonas aeruginosa</i>	26
Graph (7) percentage of Amikacin resistant <i>Pseudomonas aeruginosa</i>	27

## List of color plates

<b>Plate</b>	<b>Page</b>
Plate(1) <i>Pseudomonas aeruginosa</i> on Blood agar	28
Plate(2) <i>Pseudomonas aeruginosa</i> on MacConkey agar	28
Plate(3) <i>Pseudomonas aeruginosa</i> on Cetrimide agar	29
Plate(4) <i>Pseudomonas aeruginosa</i> Oxidase test positive	29
Plate(5) Set of Biochemical tests of <i>Pseudomonas aeruginosa</i>	30
Plate(6) <i>Pseudomonas aeruginosa</i> on Muller's Hinton agar	30