

Tables	Page No.
3-1 Texture Feature Equations	33
4-1 Normal Sudanese body characteristics (total sample)	40
4-2 Kidneys volume, and length for the total sample	41
4-3 Confusion matrix	48

List of Figures:

Figures	Page No.
2-1 Urinary system	5
2-2 Kidney anatomy	6
3-1 Kidney contouring in one slices T ₁ TSE MRI	24
3-2 Number of pixel collection.	25
3-3 Feature extraction from cortex when window size 5x5 pixels.	27
3-4 Feature extraction from medulla when window size 15x15 pixels.	27
3-5 Feature extraction from P/C system when window size 20x20 pixels.	28
4-1 Scatter plot diagram showed a linear relationship between the total body water (TBW) and RT kidney volume.	42
4-2 Scatter plot diagram showed a linear relationship between the total body water (TBW) and LT kidney volume.	42
4-3 Scatter plot diagram showed a linear relationship between the Body mass index (BMI) and RT kidney volume.	43
4-4 Scatter plot diagram showed a linear relationship between the Body mass index (BMI) and RT kidney volume.	43
4-5 Scatter plot diagram showed a linear relationship between the Height and RT kidney volume	44
4-6 Scatter plot diagram showed a linear relationship between the Height and RT kidney volume.	44
4-7 Scatter plot diagram showed a linear relationship between the Height and RT kidney length.	45
4-8 Scatter plot diagram showed a linear relationship between the Height and LT kidney length.	45
4-9 Scatter plot diagram showed a linear relationship between the LT kidney volume and RT kidney volume.	46
4-10 Classification accuracy versus window size.	47
4-11 Scatter plot portrayed the class centers with the member of the class around it versus the discriminate function.	47
4-12 Error bar plot of the feature vector for the three classes.	49

Abbreviations:

MRI.....	Magnetic Resonance Image
ROI.....	Region of interest
SGLD.....	Spatial Gary Level Depended Matrix
DSM.....	Disk Summation Method
TSE.....	Turbo Spine Echo
BMI.....	Body Mass Index
TBW.....	Total Body Water
BSA.....	Body Surface Area
C.Sr.....	Serum Ceriatinine
CrCl.....	Ceriatinine Clearance
CG-GFR.....	CookcraftGault- Glomerular Filtration Rate

Reference:

Al-Khader AA, Tamim H, Sulaiman MH, Jondeby MS, Taher S, Hejaili FF, et al. what is the most appropriate formula to use in estimating glomerular filtration rate in adult Arabs without kidney disease? *Ren Fail* 2008;30:205-8.

B. Chevaillier, B. Ponvianne, J. L. Collette, D. Mandry, M. Claudon, & O. Pietquin, (2008). Functional semi-automated segmentation of renal DCE-MRI sequences. In *Proceedings of the IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)* (pp. 525-528).

B. van Ginneken, S. Katsuragawa, and B. ter Harr Romeny B et al., "Automatic detection of abnormalities in chest radiographs using local texture analysis", *IEEE Trans Med Imag.*, Vol.21(2), 2002, pp.139-149.

Brandt TD, Neiman HL, Dragowski MJ, Bulawa W, Claykamp GU
Ultrasound assessment of normal renal dimensions *Journal Ultrasound Med.* 1982 Mar, L(2):49-52.

C.A. Binkert, U. Hoffman, D.A. Leung, H.G. Matter, M. Schmidt, J.F. Debatin, A.
. Characterization of renal artery stenoses based on magnetic resonance renal flow and volume measurements. *Kidney Int* 56:1846-1854, 1999.

C.U. McRae, F.T. Shannon, Utley WLF. Effect on renal growth of reimplantation of refluxing ureter. *Lancet* 1974; 1:1310-1312.

Cheong Raja Muthupillai, F. Mario Rubin, D Scott Flamm Normal Values for Renal Length and Volume as Measured by Magnetic Resonance Imaging *CJASN* January 2007 vol.2 no.138-45.

Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron* 1976;16:31-41.

D.Miletic , Z.Fuckar, A.Sustic,V.Mozetic,D.Stimac, G.Zauhar:
Sonographic measurement of absolute and relative renal length in adults.
*JClin Ultrasound*26 : 185-189,1998.

D.Miletic , Z.Fuckar, A.Sustic,V.Mozetic,D.Stimac, G.Zauhar:
Sonographic measurement of absolute and relative renal length in adults.
*JClin Ultrasound*26 : 185-189,1998.

D.-Y. Kim and J.-W. Park, "Computer-aided detection of kidney tumor on abdominal computed tomography scans", *Acta Radiol.* Vol.45, 2004, pp.791-795.

Emamian SA, Nielsen MB,Pedersen JF.:Intraobserver and Interobserver variations in sonographic measurements of kidney size in adult volunteers. *Acta Radiol* 36:399-401,1995.

G.N. Lee, X. Zhang, and M. Kanematsu et al.,"Classification of cirrhotic liver on MR images using texture analysis", *Int'l J. CARS*, Vol.1, Supp. 1, 2006, pp.379-381.

Gehan EA, George SL: Estimation of human body surface area from height and weight. Cancer Chemother Rep 54: 225–235, 1970.

Glodny B, Unterholzner V, Taferner B, et al. (2009). "Normal kidney size and its influencing factors - a 64-slice MDCT study of 1.040 asymptomatic patients". *BMC Urology* 9: 19. doi:10.1186/1471-2490-9-19. PMC 2813848.

H. K. Koh, S. W. Shen, B. Shuter, & A. A. Kassim, (2006). Segmentation of Kidney Cortex in MRI Studies using a constrained Morphological 3D H-maxima Transform. In *Proceedings of the 9th International Conference on Control, Automation, Robotics and Vision, (ICARCV)*, (pp. 1-5)

H. Yoshida, D. Casalino, and B. Keserci et al., "Wavelet packet based texture analysis for differentiation between benign and malignant liver tumors in ultrasound images", *Phys. Med. Biol.* Vol. 48, 2003, pp.3735-3753.

Heymsfield SB, Fulenwider T, Nordlinger B, Barlow R, Sones P, Kutner M.: Accurate measurement of liver, kidney, and spleen volume and mass by computerized axial tomography. *Ann intern Med* 90:185-187, 1979.

Ho Sik Shin, Byung Ha Chung, Sang Eun Lee, Woo Jin Kim, Hong Il Ha, and Chul Woo Yang, Measurement of kidney volume with MDCT in young Korean.

H-P. Chan, B. Sahiner, and N. Petrick et. al. "Computerized classification of malignant and benign microcalcifications on mammograms: texture analysis using an artificial neural network", *Phys. Med. Biol.*, Vol. 42, 1997, pp.549-567.

J. Xie, Y. Jiang, & H. Tsui (2005). Segmentation of kidney from ultrasound images based on texture and shape priors. *IEEE Trans. on Medical Imaging*, (Vol. 24).

J. Bakker, M. Olree, R. Kaatee, E.E. de Lange, K.G. Moons, J.J. Beutler, F.J. Beek: Renal volume measurements: Accuracy and repeatability of US compared with that of MR imaging. *Radiology* 211 :623-628, 1999.

J.R. Wilkie, M.L. Giger, and M.R. Chinander et al., "Comparison of radiographic texture analysis from computed radiography and bone densitometry systems", *Med. Phys.* Vol. 31, 2004, pp.882-891.

Joel Neugarten, Bertam Kasiske, Sharon R, Silbiger, Jens R. Nyengaard
Effects of sex on Renal Structure, *Nephron* 2002;90:139-144.

Kiw-Yong Kang, Young Joon Lee, Soon Chul Woo, Yong-Soo Kim, In Sung Moon, Yong Bok Koh, Byung Kee Bang and Bum Soon Choi. A comparative study of methods of estimating kidney length in kidney transplantation donors. *Oxford Journals, Medicine, Nephrology Dialysis Transplantation*, Volume 22, Issue 8, Pp. 2322-2327.

L. Lan, M.L. Giger, and J. R. Wilkie et al., "Automated selection of region of interest for radiographic texture analysis", *Proc SPIE Medical Imaging*, Vol. 6514, 2007, pp. 651436.

M. Marcuzzo, P. R. Masiero, & J. Scharcanski, (2007). *Quantitative Parameters for the Assessment of Renal Scintigraphic Images*, Engineering in Medicine and Biology Society, 2007.

M. Zhang, M.L. Giger, and K. Doi, "Mammographic texture analysis for the detection of spiculated lesions", In: *3rd Int'l Workshop on Digital Mammography* (eds. K. Doi, M. L. Giger, R. M. Nishikawa, R. A. Schmidt), Elsevier Science, Amsterdam, pp. 347-350, 1996.

M. Absy, C. Metreweli, C. Matthews, A. Al Khader. Changes in transplanted kidney volume measured by ultrasound. *Br J Radiol* 1987; 60:525-529.

M.R. Chinander, M.L. Giger, and R.D. Shah et al., "Investigation of using bone texture analysis on bone densitometry images", *Proc SPIE*, Vol. 4684, 2002, pp. 860-863, 2002.

Mahajan S, Mukhiya GK, Singh R, Tiwari SC, Kalra V, Bhowmik DM, et al. Assessing glomerular filtration rate in healthy Indian adults : A comparison of various prediction equations. *J Nephrol* 2005 ; 18:257-61.

N.R. Mudigonda, R.M. Rangayyan and J.E.L. Desautels, "Gradient and texture analysis for the classification of mammographic masses", *IEEE Trans. Med. Imag.*, Vol. 19(10), 2000, pp. 1032-1043.

P.Allan :The normal kidney. In :Clinical Ultrasound : A Comprehensive Text, 2nd Ed., edited by Meire H , Cosgrove D,Dewbury K,Farrant P ,NewYork, Churchill Livingstone,2001,pp513-528.

R. Gupta and P.E. Undrill, "The use of texture analysis to delineate suspicious masses in mammography", Phys. Med Biol. Vol.40, 1995, pp.835-855.

R. M Haralick,. 1979. Statistical and structural approaches to texture. Proceedings on the IEEE (67) 5: 786-803.

R. M Haralick., K. Shanmugam & I. Dinstein. 1973. Textural feature for image classification. IEEE Trans. Man, Cybern, 6:610-621.

R.H. Dean, R.W .Kieffer, B.M. Smith, et al. Reno vascular hypertension, anatomic and renal function changes during drug therapy. Arch Surg 1981; 116:1408-1415.

R.H.Breau, E.Clark ,B.Bruner,P,Cervini,T.Atwell,G.Knoll,B.C.Leibovich Simple method to staminate renal volume from CT .Can Urol Assoc J,2013 May-jun;7(5-6) :189-92.doi:10.5489/cuaj.1338.

R.J. Bartrum, E.H .Smith, C.J. D'Orsi, J .Dantonio. The ultrasonic determination of renal transplant volume . JCU 1974; 2:281-285.

R.S.Snell ,M.R.C.,L.R.C.P.,MB,BS,MD,PhD, Clinical Anatomy By System ,Lippincott Williams &Wilkins 2007.

Robert Barkow, MD ., Mark H, Beers M,D .MARK MANUAL of medical information 2009.

S .Troell, U. Berg, B. Johansson, I .Wikstad. Ultrasonographic renal parenchymal volume related to kidney function and renal parenchymal area in chlden with recurrent urinary tract infections and asymptomatic bacteriuria. Acta Radiol 1984;25:411-416

S. Poonguzhali, and G. Ravindran (2008), Automatic Classification of Focal Lesions in Ultrasound Liver Images Using Combined Texture Features. Information Technology Journal.

S. Sahiner, H.-P. Chan, and N. Petrick et al., "Computerized characterization of masses on mammograms: the rubber band straightening transform and texture analysis", Med. Phys. Vol.25(4), 1998, pp.516-526.

S.A Emamian, M.B Nielsen, J.F Pedersen, L. Yite:Kidney dimensions at sonography:Correlation with age, sex, and habitus in 665 adult volunteers. AJR Am J Roentgenol 1601 :83-86,1993.

Samuel Johnson , Rahul Rishi,Andreea And one, Wassim Khawandi , Gafar Al-Said,Nana Gletsu-Miller, Edward Lin ,Deborah A.Baumgarten , and W, Charles O'Neil Determination and Functional Significance of Renal parenchymal Volume in Adults Cin J Am Soc Nephrol.2011 January;6(1):70-76.

Sargent MA ,Long G,Karmali M,Cheng SM:.Interobserver variation in the sonographic estimation of renal volume in children. Pediatr Radiol 27:663-666, 1997.

T.B. Jones, L.R. Riddick, M.D. Harpen, R.L .Dubuisson, D. Samuel. Ultrasonographic determination of renal mass and renal volume .J Ultrasound Med 1983;2:151-4.

T.D Brandt, H.L.Neiman, M.J.Dragowski, W.Bulawa,G.Claycamp : Ultrasound assessment of normal renal dimensions.J Ultrasound Medl :49-52,1982.

V.thakur. T.Walkins,K.McCarthy, T.Beidl,N.Underwood,K.Barnes,E.M. Cook: Is kidney length a good predictor of kidney volume ?Am J Med Sci313 :85-89,1997.

Watson PE, Watson ID, Batt RD: Total body volumes for adult males and females estimated from simple anthropometric measurements. Am J Clin Nutr 33: 27-39, 1980.

Werner S Harmse Normal variance in Renal size in relation to body habits,South African Journal of Radiology,Vol 15,Noa(2011).

Widjaja E, Oxtoby JW,Hale TL, Jones PW, Harden PN, McCall IW. Ultrasound measured renal length versus low dose CT volume in predicting single kidney glomerular filtration rate .Br J Radiol 2004;77:759-64.

X. Chen, K. Doi, and S. Katsuragawa et al., "Automated selection of regions of interest for quantitative analysis of lung textures in digital chest radiographs", Med. Phys., Vol.20, 1993, pp.975-982.