

قال تعالى :

(وأنزل الله عليك الكتاب والحكمة وعلمك ما لم تكن تعلم

وكان فضل الله عليك عظيما)

سورة النساء الاية (113)

Dedication

To doses of the cup blank to give me a drop of love

To those of the fingers to give us a moment of happiness

To reap the thorns out of my way for me to pave the way science

To the heart as pure whiteness my mother

To heart the great my father

TO whom breastfed of love and healing balm

To all my family, to my Mon greenta and to my friends

Acknowledgements

First and foremost, I would like to express my deepest gratitude to Dr. Elsafi Ahmed Abdullah, for his help and guidance. I also would like to thanks Dr. Mohammed Elfadel. Without his help this work could not have been accomplished.

Abstract

In fact diabetic is the most common cause for renal failure. The purpose of the study is to characterize diabetic patient kidneys using ultrasound.

The study were done for 50 diabetic patient (30 male and 20 female) all of them have diabetic mellitus type2, their age between 45-55 year were included, and the patient with renal congenital anomalies, renal tract obstruction , malignant tumor and renal failure were excluded.

In this study ultrasound was performed for all patients by measuring the size of the kidney and evaluated the Echogensity of the right kidney compared with liver Echogensity, while the Echogensity of left kidney is compared with Echogensity of the spleen and corticomedullary ratio is also measured. The ultrasound scan is done by using TA curve linear transducer 3.5 MHZ.

The value of the study characterize that there is decrease in kidney length , decrease in C/M ratio and no deference in Echogensity.

ملخص الدراسة

يعتبر داء السكري من الأسباب الشائعة للفشل الكلوي والهدف من هذه الدراسة توصيف الكلى لدى مرض السكري بواسطة الموجات فوق الصوتية .

اجريت هذه الدراسة في 50 مريض بالسكري 30 ذكور و 20 ايناث كلهم مصابون بالسكري النوع الثاني . تتراوح اعمارهم م بين 45-55 عام وقد تم استبعاد مرض السكري المصابون بعيوب خلقية في الكلى او انسداد في المجرى البولي او اورام في الكلى.

اعتمدت الدراسة علي قياس حجم الكلى وقياس النسبة بين قشرة ونخاع الكلى بالاضافه لمعدل الصدى الراجع مقارنة بالصدى الراجع من الكبد بالنسبة للكلى اليمين والصدى الراجع من الطحال بالنسبة للكلى اليسار. كذلك قياس النسبة بين طبقة القشرة والنخاع في الكلى. ولقد اجري الفحص باستخدام المسبار البطني بواسطة جهاز موجات فوق صوتيه بتردد 3.5 هيرتز.

اوضحت النتائج وجود نقص في حجم الكلى ونقص ايضا في النسبة بين طبقة القشرة والنخاع في الكلى بينما لا يوجد تغي في معدل الصدى الراجع .

Contents

Items	Page NO.
الايه	I
Dedication	II
Acknowledgements	III
Abstract (English)	IV
Abstract (العربية)	V
Contents	VI
List of tables	IX
List of figures	X
List of abbreviation	XII
Chapter one : Introduction	
1.1 Introduction	1
1.2 Problem of study	4
1.3 Objective of study	4
1.4 significance of the study	4
1.5 Material and methods	5
1.5.1 Material	5
1.5.2 Methodology	5
1.5.3 study area	5
1.5.4 duration of the study	6
1.6 overview of the study	6
Chapter two :literature review	
2.1 embryology of the kidney	7
2.1.1 phases of development	7
2.2 anatomy of the kidney	9
2.3 physiology	17
2.4 ultrasound technique of the kidney	19
2.5 normal sonographic appearance of the kidney	19
2.6 normal renal measurement	21
2.6.1 Age related change in adult	21

2-7 D.N	21
2-6-1 Pathophysiology	22
2-6-2 signs and symptoms	23
2-6-3 treatment	23
2-6-4 prognosis	24
2-6-5 complication	24
2-7 previous study	25
Chapter three : Material and Method	
3.1 Material	31
3.1.1 machines	31
3.1.2 study population	32
3.1.3 study design	32
3.1.4 study area	32
3.1.5 duration of the study	33
3.1.6 data sheet	33
3.1.7 data analysis	33
3.1.8 study variables	33
3.2 Method	32
3.2.1 method of evaluation of volume	32
3.2.2 method of evaluation of C/M ratio	32
3.2.3 method of evaluation of echogenicity	32
Chapter four : Results	
4.1 Results	34
Chapter five : Discussion and Conclusion and Recommendation	
5.1 Discussion	40
5.2 Conclusion	43
5.3 Recommendation	44
5.4 References	45

List of tables

Table	Item	Page NO.
4.1	The mean and standard deviation of the study variables	34
4.2	Echogenicity group and their frequency	34
4.3	Cortico-medullary differentiation and their frequency	35
4.4	Gender group and their frequency distribution	35
4.5	Medication status and their frequency	36
4.6	Paired T-test result between the length, width and thickness of both kidneys.	39
4.7	Independent sample test result of medication status effect on length, width and thickness of both kidney	39

List of figures

Figure	Item	Page NO.
2.1	Developmental process of the kidney	9
2.2	Anatomy of the kidney	10
2.3	Vascular supply of the kidney	11
2.4	Superficial and juxtamedullary nephron	13
2.5	Glomerulus and many capillary loops	15
2.6	P2 receptor signaling along the nephron	18
2.7	Component of juxtamedullary apparatus	17
2.8	Sonographic appearance of normal right kidney	20
2.9	Sonographic appearance of normal left kidney	20
4.1	Echogenicity pie chart	34
4.2	CMD ratio bar chart	35
4.3	Gender distribution bar chart	36
4.4	Medication status bar chart	36
4.5	Scatter plot show correlation between duration of diabetes and length of the right kidney .	37
4.6	Scatter plot show correlation between duration of diabetes and length of the left kidney .	37
4.7	Scatter plot show correlation between duration of diabetes and CMT of the right kidney .	38
4.8	Scatter plot show correlation between duration of diabetes and CMT of the left kidney .	38

List of abbreviation	
DM.	Diabetes mellitus
IDDM.	Insulin dependent diabetes mellitus
NIDDM	Non -insulin dependent diabetes mellitus
CKD	Chronic kidney disease
DN	Diabetic nephropathy
C/M ratio	Cortico-medullary ratio
CMT	Cortico-medullary thickness
TA.	Transabdominal
K-DOQI	Kidney Disease outcomes quality
BUN	Blood urea nitrogen
ACE inhibitor	Angiotensin converting enzyme inhibitor