الآيــة

(أَفْرَأَيْتُمْ مَا تَحْرُثُونَ * أَأَنْتُمْ تَزْرَعُونَهُ أَمْ نَحْنُ الزَّارِعُونَ)

صدق الله العظيم (الواقعة 63، 64)

Dedication

To my kind Father and Mother

To the soul of professor Daw Elbait Abd Allah

To my wife and lovely daughter Omnia

And to my Brothers and Sisters

Acknowledgement

I wish to thank the Merciful God Who offered me strength and high spirit to perform this study in satisfactory form.

I am honored to express my deep gratitude indebtedness to my supervisor Ustaz Luai Osman Hashim for the valuable support criticism and close supervision Special thanks are due to Dr Gwaher Dongla S.R.C for her assistance and support for availing the land for experiments.

Thanks to Ustas Ashwag Ahmed Mohamed for her help and making photographs

And thanks are due to my colleagues in faculty of agriculture Department of plant protection in Sudan University

List of contents

Tit	es	
	i	
	Dedicationii	
	Acknowledgementiii	
	List of contentiv	
	List of tablesvi	
	List of figuresvii	
	List of platesviii	
	English Abstractix	
	Arabic Abstractx	
	Chapter one: Introduction1	
1-1	Tomato	
	1-1-1 Tomato Production in Sudan	
	1-1-2 Nutritive Value of tomato	
	1-1-3 Botanical characteristics	
	1-1-4 Classification	
	1-1-5 Pest and Diseases	
	1-1-5-1 Insects	
	1-1-5-2 Diseases	
	1-1-5-2-1 Fungal Disease	
	1-1-5-2-1-1 Powdery Mildew	
	1-1-5-2-1-2 Late Blight5	
	1-1-5-2-1-3 Fusarium Wilt5	
	1-1-5-2-1-4 Damping off Disease5	
	1-1-5-2-1-5 Dark Brown5	
	1-1-5-2-1-6 Early Blight Disease5	
	1-1-5-2-2 Bacterial Diseases6	
	1-1-5-2-2-1 Bacterial Wilt6	
	1-1-5-2-2 Bacterial Canker 6	
	1-1-5-2-3 Viral Diseases6	
	1-1-5-2-4 Nematode Diseases6	
	1-1-5-2-5 Parasitic Flowering Plants6	
	1-2 This Research	
	1-3 Objectives of the Research	
	Chapter Two: Literature Review8	
	2-1 Alternaria8	
	2-1-1 History of Early Blight Diseases	
	2-1-2 Alternaria Diseases9	
	2-1-3 Geographical Distribution	

References	39
Chapter Five : Discussion and Conclusion	
Chapter Four: The Results	
3-8 Harvesting	
3-7 Data Collection	
3-6 Treatment	
3-5 Fertilization	
3-4 Weeding	
3-3 Irrigation and Thinning	
3-2 Sowing	
3-1 Land Preparation	20
Chapter Three: Material and Methods	20
2-3-2-3 Uses	
2-3-2-2 Active Ingredient	
2-3-2-1 Mode of Action	
2-3-2 Thiovit 80 WG	
2-3-1-2 Uses	
2-3-1-1 Active Ingredient	
2-3-1 Bayleton 25% WP	18
2-3 The Role of Chemical Products on Diseases	
2-2-2 Ginger (Zingiber officinale, L)	
2-2-1-3 Biological Activities of Neem Products	17
2-2-1-2 Chemistry of Neem Tree	
2-2-1-1 Origin and Characteristics of Neem Tree	
2-2-1 Neem (<i>Azodrichta indica A . Juss</i>)	
2-2 The Role of Natural Products on Diseases	
2-1-10-2 Diological Control	
2-1-10-1 Cultural Control	
2-1-10 Control Measures 2-1-10-1 Cultural Control	
2-1-9 The Occurrence of Early Blight	
2-1-8 Etiology	
2-1-7 Host Plants	
2-1-6 Damage	
2-1-5 Disease Cycle	
2-1-4 Infection and Symptoms	

List of Table

Table (1): Effects of different Treatment on Early Blight disease incidence
Alternaria solani (on Leaves) Post Spray One25
Table (2): Effects of different Treatment on Early Blight disease incidence
Alternaria solani (on Leaves) Post Spray Two28
Table (3): Percentage of damage in number of Fruits by Alternaria solani
31
Table (4): Percentage of damage on Fruits weight infected by Alternaria
solani34

List of Figures

Figure (1): Effects of different Treatment on Early Blight disease incidence
Alternaria solani (on Leaves) Post Spray One26
Figure (2): Effects of different Treatment on Early Blight disease incidence
Alternaria solani (on Leaves) Post Spray Two29
Figure (3): Percentage of damage in number of Fruits by Alternaria solan
32
Figure (4): Percentage of damage on Fruits weight infected by Alternaria
solani35

List of Plates

Plate (1): Early Blight diseases symptoms on infec	cted Tomato Leaves
	27
Plate (2): Healthy Tomato Leaves	27
Plate (3): Early Blight disease symptoms on infec	ted Tomato Plant.
	30
Plate (4): Healthy Tomato Plant	30
Plate (5): Early Blight disease symptoms on Tomato F	Fruit33
Plate (6): Healthy Tomato Fruit	33
Plate (7): Weight Fruit	33
Plate (8): Alternaria solani Spores	36
Plate (9): Alternaria alternata Spores	36

ABSTRACT

On a study carried out at Shambat horticultural research station, to compare between the effect of some Natural products (Neem, Ginger) and fungicides (Bayleton, Thiovit). In disease incidence early blight on tomato caused by *Alternaria solani*. The effects of Neem and Ginger was high as compared to the effects of fungicides on fruits.

Both Neem and Ginger gave significant reduction of the disease in the number of infected leaves as compared to control while their effect was not significant on number of infected fruits. The effect of chemical was significant and clearly observed on the number of infected leaves compared to the control but their effect on fruit was not significantly differents the effect of the Mixture(Neem and Ginger) was the best in the reduction of the number of infected fruits . where there was not significantly differences between control

.