

5.0 References

- Abdel Hadi, A.M.** frequencies of ABO and Rhesuse system blood group among Al Shuluk tribe ethnic group approved in Sudan university of science and technology Khartoum. ,(2005).
- Abeer, A.A,** frequencies of ABO and Rhesuse system blood group among Beni Amir ethnic group approved in Sudan university of science and technology Khartoum . ,(2005)
- Allen FH Jr,** Krabbe SM, Corcoran PA. A new phenotype (McLeod) in the Kell blood-group system. *Vox Sang.* 1961 Sep;6:555-60. [PMID 13477267](#)
- Allen, F. H. &** Rosenfield, R. E. (1972). Review of Rh serology: eight new antigens in nine years. *Haematologia*, 6:113-120. . .
- Arce, M. A.;** Thompson, E. S.; Wagner, S.; Coyne, K. E.; Ferdman, B. A.; Lublin, D. M. : Molecular cloning of RhD cDNA derived from a gene present in RhD-positive, but not RhD-negative individuals. *Blood* 82: 651-655, 1993.
- Avent, N. D. &** Reid, M. E. (2000). The Rh blood group system: a review. *Blood*; 95:1-13.
- Avent, N. D.,** Liu, W., Warner, K. M., Mawby, W. J., Jones, J. W., Ridgwell, K. & Tanner, M. J. (1996). Immunochemical analysis of the human erythrocyte Rh polypeptides. *J Biol Chem.* 271:14,233-14,239.
- Avent, N. D.,** Martin, P. G., Armstrong-Fisher, S. S., Liu, W., Finning,, K. M., Maddocks, D. & Urbaniak, S. J. (1997). Evidence of genetic diversity underlying Rh D^r, weak D (D^u), and partial D phenotypes as determined by multiplex polymerase chain reaction analysis of the *RhD* gene. *Blood*.89:2568-2577.
- Bennett, P. R.;** Le Van Kim, C.; Colin, Y.; Warwick, R. M.; Cherif-Zahar, B.; Fisk, N. M.; Cartron, J.-P. : Prenatal determination of fetal RhD type by DNA amplification. *New Eng. J. Med.* 329: 607-610, 1993.

- Blancher, A.** & Socha, W. W. (1997). The Rhesus system. In: Blancher A, Klein J, Socha WW, eds. *Molecular Biology and Evolution of Blood Group and MHC*
- Blunt, T.,** Daniels, G., & Carritt, B. (1994). Serotype switching in a partially deleted RHD gene. *Vox Sang.*;67:397-401.
- Bowman, J. M.** : RhD hemolytic disease of the newborn. (Editorial) *New Eng. J. Med.* 339: 1775-1777, 1998.
- Cartron, J.-P.** : Defining the Rh blood group antigens: biochemistry and molecular genetics. *Blood Rev.* 8: 199-212, 1994.
- Castilho, L.,** Rios, M. & Pellegrino, J. Jr. (1999). Molecular analysis of the RHD gene shows a high incidence of partial D, D^{IIIa} among sickle cell disease patients. *Ser. Monogr Esc Bras Hematol*, 6 (Suppl):17^a.
- Cherif-Zahar, B.,** Raynal, V.; Gane, P.; Mattei, M.-G.; Bailly, P.; Gibbs, B.; Colin, Y.; Cartron, J.-P. : Candidate gene acting as a suppressor of the RH locus in most cases of Rh-deficiency. *Nature Genet.* 12: 168-173, 1996.
- Cherif-Zahar, B.,** Raynal, V.; Le Van Kim, C.; D'Ambrosio, A. M.; Bailly, P.; Cartron, J. P.; Colin, Y. : Structure and expression of the RH locus in the Rh-deficiency syndrome. *Blood* 82: 656-662, 1993.
- Colin, Y.,** Cherif-Zahar, B.; Le Van Kim, C.; Raynal, V.; Van Huffel, V.; Cartron, J.-P. : Genetic basis of the RhD-positive and RhD-negative blood group polymorphism as determined by Southern analysis. *Blood* 78: 2747-2752, 1991.
- Coombs RRA,** Mourant AE, Race RR. *A new test for the detection of weak and "incomplete" Rh agglutinins.* *Brit J Exp Path* 1945;26:255-66.
- Dacie** (2001)., Sir John v, Dacie, and Lewis. *Practical Hematology*. 8th edn. Churchill Livingstone, Edinburgh, 2001:424-29.
- Daniels G,** Finning K, Martin P, Summers J. "Fetal blood group genotyping: present and future" *Ann N Y Acad Sci* 1999. 1075: 88-95

Daniels, G. L., Anstee, D. J., Cartron, J. P., Dahr, W., Issitt, P. D., Jorgensen, J., Kornstad, L., Levene, C., Lomas-Francis, C. & Lubenko. (1995). Blood group terminology: ISBT working party on terminology for red cell surface antigens. *Vox Sang*, 69:265-279.

Diamond, L. K.; Blackfan, K. D.; Baty, J. M. :Erythroblastosis fetalis and its association with universal edema of the fetus, icterus gravis neonatorum and anemia of the newborn. *J. Pediat.* 1: 269-309, 1932.

Ernest Beutler,(2001). William's Hematology, 6th edn. The Mc Graw,Hill Companies, Sanfrancisco, 2001:1840-53.

Ernest Beutler,(2001). William's Hematology, 6th edn. The Mc Graw,Hill Companies, Sanfrancisco, 2001:1840-53.

Eyers, S. A., Ridgwell, K., Mawby, W. J. & Tanner, M. J. (1994). Topology and organization of human Rh (rhesus) blood group-related polypeptides. *J Biol Chem*, 269:6417-6423.

Faas, B. H. W., Beckers, E. A. M. & Simsek, S. (1996). Involvement of Ser103 of the Rh polypeptides in G epitope formation. *Transfusion*, 36:506-511.

Faas, B. H. W., Beckers, E. A. M., Wildoer, P., Smart, E., Maaskant-van Wijk, P. A., Avent, N. D., Zondervan, H. A., von dem Borne, A. E. & van der Schoot, C. E. (1997). Molecular background of VS and weak C expression in blacks. *Transfusion*, 37:38-44.

Hameed, A., Wajahat, H., Janbaz, A., Fazil-Rai. & Javed, A. Q. (2002). Prevalence of phenotypes and genes of ABO and RH Rhesus blood group in Faisalabad Pakistan. *Paki J boil scie*, 5 (6):722-724.

Hoffbrand, A. V. & Pettit, A. E. (2001). Post graduate hematology. 4th edition, British library, London.:330-370.

Huang, C. H., Chen, Y. & Reid, M. (1997). Human D^{IIIa} erythrocytes: RhD protein is associated with multiple dispersed amino acid variations. *Am J Hematol*, 55:139-145

- Huang, C.-H.;** Reid, M. E.; Chen, Y.; Coghlan, G.; Okubo, Y. : Molecular definition of red cell Rh haplotypes by tightly linked SphI RFLPs. *Am. J. Hum. Genet.* 1996 58: 133-142,.
- Hyland, C. A.;** Wolter, L. C.; Liew, Y. W.; Saul, A.. A Southern analysis of Rh blood group genes: association between restriction fragment length polymorphism patterns and Rh serotypes. *Blood* 1994 83: 566-572,.
- Innan, H. :** A two-locus gene conversion model with selection and its application to the human RHCE and RhD genes. *Proc. Nat. Acad. Sci.* 2003. 100: 8793-8798,.
- International Society of Blood Transfusion** [Table of blood group systems](#). (October 2006). Retrieved on [2006-11-14](#).
- Ishimori, T.;** Hasekura, H. : A Japanese with no detectable Rh blood group antigens due to silent Rh alleles or deleted chromosomes. *Transfusion* 7: 84-87, 1967.
- Issitt, P. D.** An invited review: the Rh antigen e, its variants, and some closely related serological observations. *Immunohematology*, (1991). 7:29-36.
- Issitt, P. D. :** The Rh blood group system, 1988: eight new antigens in nine years and some observations on the biochemistry and genetics of the system. *Transfusion Med. Rev.* 1989 3: 1-12,.
- Jaccard, P.** Etude comparative de la distribution florale dans une portion des Alpes et des jura. *Bulletin de la Societe Vaudoise des sciences Naturelles*, (1991). 37: 547-479.
- Jeremih z a,**Odumody C 2005 Rh antigens and phenotype frequency of theIbibo, EFink,and Ibo ethnic group nationality in Calabar,Nigeria
- Kolmakova, G. N. & Kononov, L. L.** (1999). The prevalence of ABO blood groups among perons of native nationality in Buryatia. *Sud. Med. Expert*, 42:15-16.

Lacey, P. A., Caskey, C. R., Werner DJ, Moulds, J.J. (1983). Fatal hemolytic disease of a newborn due to anti-D in an Rh-positive Du variant mother. *Transfusion*;23:91-94.

Land Steiner K, Wiener AS. *An agglutinable factor in human blood recognized by immune sera for rhesus blood.* Proc Soc Exp Biol Med 1940;43:223-224. .

Landsteiner K. *Zur Kenntnis der antifermentativen, lytischen und agglutinierenden Wirkungen des Blutserums und der Lymphe.* Zentralblatt Bakteriologie 1900;27:357-62.

Le Van Kim, C.; Cherif-Zahar, B.; Raynal, V.; Mouro, I.; Lopez, M.; Cartron, J. P.; Colin, Y. : Multiple Rh messenger RNA isoforms are produced by alternative splicing. *Blood* 80: 1074-1078, 1992.

Le Van Kim, C.; Mouro, I.; Cherif-Zahar, B.; Raynal, V.; Cherrier, C.; Cartron, J.-P.; Colin, Y. : Molecular cloning and primary structure of the human blood group RhD polypeptide. *Proc. Nat. Acad. Sci.* 89: 10925-10929, 1992.

Legler, T. J., Maas, J.H., Blaschke, V., Malekan, M., Ohto, H., Lynen, R., Bustami, N., Schwartz, D.W., Mayr, W. R., Kohler, M. & Panzer, S. (1998). *RHD* genotyping in weak D phenotypes by multiple polymerase chain reactions. *Transfusion*, 38:434-440.

Levine, P., Celano, M. J., Wallace, J. & Sanger, R. (1963). A human 'D-like' antibody. *Nature*. 198:596-597.

Lo, Y. M. D.; Hjelm, N. M.; Fidler, C.; Sargent, I. L.; Murphy, M. F.; Chamberlain, P. F.; Poon, P. M. K.; Redman, C. W. G.; Wainscoat, J. S. : Prenatal diagnosis of fetal RhD status by molecular analysis of maternal plasma. *New Eng. J. Med.* 339: 1734-1738, 1998.

- Lögberg, I.**, Reid, M. E. & Miller, J. L. (2002). Cloning and genetic characterization of blood group carrier molecules and antigens. *Transf Med Rev*, 16:1-10.
- Lomas, C.**, Tippet, P., Thompson, K. M., Melamed, M. D. & Hughes-Jones, N. C. (1989). Demonstration of seven epitopes on the Rh antigen D using human monoclonal anti-D antibodies and red cells from D categories. *Vox Sang*, 57:261-264.
- Mahmoud, L.A-N.** Ibrahim, A.A-K. Ghonem, H.R.. Jouvenceaux ,A (1987). Human Blood Groups in Dakahlya, Egypt (14) 487 - 493
- Malik, H .I.**,Khalid,EAG.,Musa,AM,and ELagi,A.H(2007) Rhesus blood group system alleles, haplotype and most probable genotype in major Sudanese population approved in Sudan university of science and technology Khartoum.
- Miller LH**, Mason SJ, Clyde DF, McGinniss MH. "The resistance factor to Plasmodium vivax in blacks. The Duffy-blood-group genotype, FyFy." *N Engl J Med*. 1976 Aug 5;295(6):302-4
- Mollison, P. L.**, Engelfriet, C. P. & Marcelac. (1997). Blood transfusion in clinical medicine. 10th edition, Oxford Blackwell scientific publications.
- Moss,W.L.**(1910). Studies on iso-agglutinins and iso-hemolycins. Bulletin Johns Hopkins Hospital,**21**:1090-1095.
- Mouro, I.**, Colin, Y., Chérif-Zahar, B., Cartron, J. P. & Le Van Kim C. (1993). Molecular genetic basis of the human Rhesus blood group system. *Nature Genet*,5:62-65.
- Mouro, I.**, Le Van Kim, C., Rouillac, C., Van Rhenen, D. J., Le Pennec, Y. & Bailly, P. (1994). Rearrangements of the blood group RhD gene associated with the D^{VI} category phenotype. *Blood*, 83:1129-1135.
- Nash, R.; Shojan, A. M. :** Hematological aspect of Rh deficiency syndrome: a case report and a review of the literature. *Am. J. Hemat.* 24: 267-275, 1987.

Neville, J. B. (1994). An introduction to immunohematology. 3rd edition, Toronto: Congress L.

Noizat-Pirenne, F., Mouro, I., Gane, P., Okubo, Y., Hori, Y., Rouger, P., Le Pennec, P. Y. & Cartron, J. P. (1998). Heterogeneity of blood group RhE variants revealed by serological analysis and molecular alteration of the *RHCE* gene and transcript. *Br J Haematol*, 103:429-436.

Okuda, H.; Kawano, M.; Iwamoto, S.; Tanaka, M.; Seno, T.; Okubo, Y.; Kajii, E. : The RHD gene is highly detectable in RhD-negative Japanese donors. *J. Clin. Invest.* 100: 373-379, 1997.

Olsson, M. L., Irshaid, N. M., Hosseini-Maaf, B. & Hellberg, A. (2001). Genomic analysis of clinical samples with serologic ABO blood grouping discrepancies: identification of 15 novel A and B subgroup alleles. *Blood*, 98:1585-93.

Omotade, O.O., Adeyemo, A.A., Kayde, C.M., Falade, S.L., & IKpeme, S. (1999): gene frequencies of ABO and Rh(D) blood group alleles in health infant population in Ibadan , Nigeria. *west Africa medical journal* 1999;18: 294- 297.

Perez-Perez, C.; Taliano, V.; Mouro, I.; Huet, M.; Salat-Marti, A.; Martinez, A.; Rouger, P.; Cartron, J.-P. :Spanish Rh-null family caused by a silent Rh gene: hematological, serological, and biochemical studies. *Am. J. Hemat.* 40: 306-312, 1992.

Petz, L. D. & Garratty, G. (1980). *Acquired Immune Hemolytic Anemias*. New York: Churchill Livingstone.

Race, R. R. : An 'incomplete' antibody in human serum. Letter) *Nature* 153: 771-772, 1944.

Reid, M. E. (2003). Applications of DNA-based assays in blood group antigen and antibody identification. *Transfusion*, 43: 1.748-1.757.

Reid, M. E., Halverson, G. R., Roubinet, F., Apoil, P. A. & Blancher, A. (1998). Use of LOR-15C9 monoclonal anti-D to differentiate erythrocytes with the partial D^{VI} antigen from those with other partial D antigens or weak D antigens. *Immunohematology*, 14:89-93.

- Reid, M. E.,** Rios, M. & Powell, D. (2000). DNA from blood samples can be used to genotype patients who have recently received a transfusion. *Transfusion*, 40:48-53.
- Ridgwell, K.,** Spurr, N. K., Laguda, B., MacGeoch, C., Avent, N. D. & Tanner M. J. (1992). Isolation of cDNA clones for a 50 kDa glycoprotein of the human erythrocyte membrane associated with Rh (rhesus) blood-group antigen
- Rios, M.,** Hue-Roye, K. & Lee, A. H. (2001). DNA analysis for the Dombrock polymorphism. *Transfusion*, 41:1.143-6.
- Rios, M.,** Reid, M. E. & Naime, D. (1997). Importance of GATA box analysis in genotyping for the Duffy blood group system. *Transfusion*, 37(S):101Sa.
- Rosenfield, R. E.,** Allen, F. H. Jr. & Rubinstein, P. (1973). Genetic model for the Rh blood-group system. *Proc Natl Acad Sci U S A*, 70:1303-1307.
- Rosenfield, R. E.,** Allen, F. H. Jr., Swisher, S. N. & Kochwa, S. (1962). A review of Rh serology and presentation of a new terminology. *Transfusion*, 2:287-312.
- Rouillac, C.,** Gane, P., Cartron, J., Le Pennec, P. Y., Cartron, J. P. & Colin, Y. (1996). Molecular basis of the altered antigenic expression of RhD in weak D (D^u) and RhC/e in R^N phenotypes. *Blood*, 87:4853-4861.
- Rouillac, C.,** Le Van Kim, C., Beolet, M., Cartron, J. P. & Colin, Y. (1995). Leu110Pro substitution in the RhD polypeptide is responsible for the D^{VII} category blood group phenotype. *Am J Hematol*, 49:87-88.
- Salah Abd Elgadir** Ibrahim(2005) frequency of ABO and Rhesuse system blood group among Niwer ethnic group Sudan university of science and technology Khartoum .
- Simsek, S.,** de Jong, C. A., Cuijpers, H. T., Bleeker, P. M., Westers, T. M., Overbeeke, M. A., Goldschmeding, R., van der Schoot, C. E. & von dem Borne, A. E. (1994). Sequence analysis of cDNA derived from reticulocyte mRNAs coding for Rh polypeptides and demonstration of of E/e and C/c polymorphisms. *Vox Sang*, 67:203.

Sloan, S. R. & Silberstein, L. E. (1998). Transfusion in the face of autoantibodies. In: Reid ME, Nance SJ, eds. Red Cell Transfusion: A Practical Guide. Totowa, NJ: Humana Press, :55-70.

Smythe, J. S.; Avent, N. D.; Judson, P. A.; Parsons, S. F.; Martin, P. G.; Anstee, D. J. : Expression of RHD and RHCE gene products using retroviral transduction of K562 cells establishes the molecular basis of Rh blood group antigens. *Blood* 87: 2968-2973, 1996.

Steers, F., Wallace, M., Johnson, P., Carritt, B. & Daniels, G. (1996). Denaturing gradient gel electrophoresis: a novel method for determining Rh phenotype from genomic DNA. *Br J Haematol*, 94:417-421.

Tippett, P., Lomas-Francis, C. & Wallace, M. (1996). The Rh antigen D: partial D antigens and associated low incidence antigens. *Vox Sang*, 70:123-131.

Vengelen-Tyler, V. (1999). Technical Manual. Bethesda, MD: American Association of Blood Banks.

Wagner F, Kasulik F, Kerowgan M, Figel WA. Frequency of ABO, Rhesus, D category vi, and clinically relevant high frequency antigens in Germany. *Transfusion* 1995;22:285-90

Wagner, F. F. & Flegel, W. A. (2004). The molecular basis of Rh blood group phenotypes (review). *Immunohematology*, 20:23-36.

Wagner, F. F., Gassner, C., Müller, T. H., Schönlitz, D., Schunter, F. & Flegel, W. A. (1999). Molecular basis of weak D phenotypes. *Blood*, 93:385-393.

Westhoff, C. M., Silberstein, L. E. & Sipherd, B. (1998). Altered "e" antigen expression associated with ¹⁶Cys in exon 1 of the Rhce gene [abstract]. *Transfusion*, 38(Suppl):64S.

Wiener, A. S. : The Rh series of allelic genes. *Science* 100: 595-597, 1944.

Wu, G. G., Jin, Z. H. & Deng, Z. H. (2001). Polymerase chain reaction with sequence-specific primers-based genotyping of the human Dombrock blood group DO1 and DO2 alleles and the DO gene frequencies in Chinese blood donors. *Vox Sang*, :81:49-51.