

.Table 5.7 Summary results of hemispheric activation differences in 5 tests on 7 subjects for combination method

(Occipital(O1,O2					(Parietal(P3,P4					(Central (C3,C4					
β_h	β_l	α	θ	δ	β_h	β_l	α	θ	δ	β_h	β_l	A	θ	Δ	Freq Band
Composite Differences															Task
-	-	-	-	-	-	0.0264	0.0501	0.0212	-	-	0.0137-	0.019	-	-	1
0.30	0.28	0.151	0.0	0.07	0.0463				0.0275	0.2		7	0.049	0.2	
98	89	1	810	94						48			2	238	
										9					
-	-	-	-	-	-	0.0264	0.0501	0.0212	-	-	0.0137-	0.019	-	-	2
0.30	0.28	0.151	0.0	0.07	0.0463				0.0275	0.2		7	0.049	0.2	
98	89	1	810	94						48			2	238	
										9					
-	-	-	-	-	-	0.0264	0.0501	0.0212	-	-	0.0137-	0.019	-	-	3
0.30	0.28	0.151	0.0	0.07	0.0463				0.0275	0.2		7	0.049	0.2	
98	89	1	810	94						48			2	238	
										9					
-	-	-	-	-	-	0.0264	0.0501	0.0212	-	-	0.0137-	0.019	-	-	4
0.30	0.28	0.151	0.0	0.07	0.0463				0.0275	0.2		7	0.049	0.2	
98	89	1	810	94						48			2	238	
										9					

-	-	-	-	-	-	0.0264	0.0501	0.0212	-	-	0.0137-	0.019	-	-	5
0.30	0.28	0.151	0.0	0.07	0.0463				0.0275	0.2		7	0.049	0.2	
98	89	1	810	94						48			2	238	
										9					