

.Table 5.5 Summary results of hemispheric activation differences in 5 tests on 7 subjects for FFT technique removing method

(Occipital(O1,O2					(Parietal(P3,P4					(Central (C3,C4					
β_h	β_l	α	θ	δ	β_h	β_l	α	θ	δ	β_h	β_l	A	θ	Δ	Freq Band
Composite Differences															Task
0.0285	0.0174	0.0484	0.0153	-0.0059	-0.0270	-0.0169	0.0124	-0.000398	-0.0059	-0.0613	-0.0229	0.0161	-0.00085	-0.0136	1
0.0459	0.076	0.0209	0.0160	0.0014	0.0166	-0.0736	-0.0559	0.0041	0.0054	-0.0036	-0.0360	-0.0384	0.0081	0.0068	2
0.0580	0.0490	0.0698	0.0268	0.0068	0.0046	-0.0231	-0.0139	0.0100-	-0.0051	-0.0179	-0.0349	-0.0411	0.0131-	0.144-	3
0.0374	0.0126	0.0330	0.0206	0.0201	0.0032	-0.0487	-0.0143	0.0056	-0.0046	-0.0038	-0.0263	-0.0249	0.0111	-0.0138	4

0.009 2	0.021 7	0.042 4	0.019 6	0.002 7	- 0.00 62	- 0.037 4	- 0.130	0.0066	0.001 9	- 0.023 5	- 0.026 8	- 0.01 84	0.0014	- 0.0048	5
------------	------------	------------	------------	------------	-----------------	-----------------	------------	--------	------------	-----------------	-----------------	-----------------	--------	-------------	---