

## **ACKNOWLEDGMENTS**

First of all my thanks and praise are due to my Almighty Allah, the Beneficent, the Merciful for giving me the health and strength to accomplish this work.

My thanks are due to Professor Osman Saad Ali, Dean of the College of Technology and Human Development; former Dean of College of Veterinary Medicine and Animal Production for providing this specialization and help to establish it.

I wish to express my indebtedness and sincere thankfulness to my supervisor Dr Bushra Hussein Ahmed, College of Radiological Sciences and Nuclear Medicine. National Rabat University and my co-supervisor Dr Amel Omer Bakhiet, College of Veterinary Medicine and Animal Production, Sudan University of Science and Technology (SUST), for their keen guidance, valuable assistance, advice and encouragement. My special gratitude to the Sudan University of Science and Technology (SUST), who sponsored this work.

My grateful thanks are due to the Police Dog Administrator, Ministry of Interior for providing the ultrasound machine; especial thanks are due to Dr Ali Hassan Ali.

Especial thanks are due to the Ministry of Agriculture, Animal Resources and Irrigation, Khartoum State, for providing the experimental animals; Saanen goats. Especial thanks are due to Dr Tageldin Osman Saeed and Dr Haleema Salih Ahmed.

I am also indebted to Professor Ahmed Abdelrahim Gameel, Head of the Department of pathology, Faculty of Veterinary Medicine, University of Khartoum, and Dr Shadia Abdel Atti Omer, College of Veterinary Medicine and animal Production, Sudan University of Science and Technology for their endless help and guidance.

I am grateful to Dr. Alsafi Ahmed Abdallah College of Medical Radiological Sciences, Sudan University of Science and Technology

I am also indebted to Dr Salah Kargweel and Alsir Ali Saeed, Ribat University Teaching Hospital for their keen support and assistance.

I am also grateful to Prof Abdelgadir Musa Homeida, King Faisal University, Kingdom of Saudi Arabia for doing the Progesterone assay.

My grate thanks are due to Dr Mohamed Tageldin Ibrahim for doing the statistical analysis (SPSS).

I am also indebted to my sincere friends Rabab and Somaia for their endless help. I would like to thank Mohamed Abdulrahim for preparing the images. My thanks are due to the staff of Sudan University of Science and Technology web page, especial thanks to Bakri Mohamed. My thanks are also due to my student Makki and Yousif (Sudan Heart Center) for preparing the ultrasound images.

Thanks are due to all those whom I forgot to mention.

عن النبي صلى الله عليه وسلم قال:-  
الغنم بركة والإبل عز لأهلها والخيل معقود بنواصيها الخير إلى يوم  
القيامة.

( اخرجه البزار )

حديث شريف

وعنه صلى الله عليه وسلم قال:-  
الغنم من دواب الجنة فامسحوا رغامها وصلوا في مراتبها.  
(أخرجه الخطيب البغدادي)  
حديث شريف

وعنه صلى الله عليه وسلم قال:-  
الفخر والخيلاء في أهل الإبل والسكينة والوقار في أهل الغنم.  
(اخرجه أحمد)  
حديث شريف

وعنه صلى الله عليه وسلم قال:-  
ما بعث الله من نبي إلا ورعى الغنم.  
حديث شريف

# LIST OF CONTENTS

Dedication	
Acknowledgements	I
List of Contents	iv
List of Figures	vii
List of Tables	viii
List of Abbreviations	ix
Abstract	xii
ملخص الاطروحة	xiv

## CHAPTER ONE INTRODUCTION 1-3

INTRODUCTION	1
The Objectives	3

## CHAPTER TWO LITERATURE REVIEW 4-19

2.1.	Goats in Sudan	4
2.1.1	Saanen goats	4
2.2.	Estrus cycle	4
2.3.	Gestational period and placentation	5
2.4.	Methods of pregnancy diagnosis	5
2.4.1.	Non-return to estrus	5
2.4.2.	Abdominal ballottement	6
2.4.3.	Laboratory methods	6
2.4.3.1.	Progesterone (P4) assay	6
2.4.3.2.	Early pregnancy factor (EPF)	7
2.4.3.3.	Pregnancy specific protein B (PSPB)	8
2.4.4.	Radiography	8
2.4.5.	Ultrasound	8
2.4.5.1.	Sound waves definition	9
2.4.5.2.	Spectrum of sound waves	9
2.4.5.3.	Production and detection of sound waves	9
2.4.5.3.1.	Transducers	9
2.4.5.3.2.	Transducers arrays	9
2.4.5.4.	Different ultrasonic methods for pregnancy diagnosis	10
2.4.5.4.1	Amplitude modulation (A- Mode)	10
2.4.5.4.2.	Metadata ultrasonic pregnancy detectors	10
2.4.5.4.3.	Brightness Modulation (B-Mode)	11
2.4.5.4.3.1.	Early detection of pregnancy, embryo proper	11

	with its heart beats	
2.4.5.4.3.2.	Fetal number (multiplicity)	12
2.4.5.4.3.3.	Fetal sexing	13
2.4.5.4.3.4.	Estimation of gestational age	14
2.4. 5.4.3.4.1.	Crown Rump length (CRL)	14
1. 5. 5.4. 3. 4. 2.	Bi Parietal Diameter (BPD)	14
2.4.5.4.3.4.3.	Abdominal circumference (AC)	16
2.4.5.4. 3.4.3.	Femur length (FL)	16
2.4.3.3.3.4.5.	Placentomes size and shape	16
2.4. 5.3. 3. 4.6.	Fetal Heart Rate (FHR)	17
2.4.5.4.3.5.	Fetal anomalies	17

### **CHAPTER THREE MATERIALS AND METHODS 20-29**

3.1.	Experimental Animals and Housing	20
3.2.	Experimental techniques	20
3.2.1.	Abdominal ballottement	20
3.2.2.	Hormonal assay	20
3.2.3.	Ultrasound	21
3.2.3.1.	Animal preparation	21
3.2.3. 2.	Animal positioning	21
3.2.3.3.	Area of scanning and scanning technique	22
3.2.3.4	Ultrasound machine	22
3.2.3.5.	Hard copy imager printer	22
3.3.	Experimental design	22
3.3.1.	Experiment (1)	22
3.3.2.	Experiment (2)	23
3.3.2.1.	CRL	23
3.3.2.2.	BPD	23
3.4.	Statistical analysis	24
3.4.1.	Experiment 1	24
3.4.2.	Experiment 2	24

### **CHAPTER FOUR RESULTS 30-59**

4.1.	Experiment 1	30
4.1.1	Abdominal ballottement	30
4.1.2	Progesterone (P4) assay	30
4.1.3	Ultrasound findings	30
4.2.	Experiment 2	41
4.2.1	Ultrasound	41

4.2.1.1 Fetometry	41
4.2.1.1.1 Crown Rump Length (CRL)	41
4.2.1.1.2. Bi Parietal Diameter (BPD)	44
4.2.1.1.3 Placentomes size and shape	47
4.2.1.1.4 Other fetal parameters	51
4.2.1.1.5. Fetal sex	56
4.2.1.1.6. Fetal anomalies	56

## **CHAPTER FIVE DISCUSSION 60-64**

Conclusion and Recommendations	63
--------------------------------	----

## **REFERENCES 65-71**

## **LIST OF FIGURES**

<b>No.</b>	<b>Figure</b>	<b>Page</b>
1	Animal Positioning (Dorsal Recumbency)	25
2	Ultrasound machine	26
3	CRL at day 53. Measuring 6.41cm	27
4	CRL at day 60. Measuring 9.76cm	28
5	BPD complete line.	29
6	Gestational Sac (GS) at day 26 of gestation	34
7	Two gestational sacs (GS) at day 28 of gestation.	35
8	Cotyledons (C) at day 87 of gestation	36
9	Fetal head (H) with spine	37
10	Fetal Rib Cage	38
11	Twins at day 38 of gestation	39
12	Twins at day 60 of gestation	40
13	The relationship between the CRL (cm) and gestational age (days)	43
14	The relationship between the BPD (cm) and gestational age (days)	46
15	Cotyledons(C) at day 46 of gestation	48

16	Cotyledons at day 60 of gestation	49
17	Cotyledons (C) at day 88 of gestation	50
18	Fetal heart rate (FHR) at day 97 of gestation measuring 195 beats/min	52
19	Femur length measuring 4.4 cm at Day 97 of gestation	53
20	Cross section of the UC	54
21	Longitudinal section of the umbilical cord (UC)	55
22	Male fetus (penis)	57
23	Male fetus (scrotum)	58
24	Female fetus	59

## LIST OF TABLES

No.	Table Title	Page
Table 1.	Pregnancy diagnosis using progesterone concentration and ultrasound	32
Table 2	Fetal number using ultrasound compared with delivery	33
Table 3.	Measurements of (CRL) in Saanen goats	42
Table 4.	Measurements of (BPD) in Saanen goats	45

## **LIST OF ABBREVIATIONS**

AC	Abdominal Circumference
AI	Artificial Insemination
BPD	Bi Parietal Diameter
Bpm	Beats per minute
CL	Corpus Luetium
CRL	Crown Rump Length
CSP	Cavum Septum Pellucidum
EGA	Estimated Gestational Age
EV	Embryonic Vesicle
FC	Falx Cerebri Midline
FHR	Fetal Heart Rate
FL	Femur Length
GS	Gestational Sac
HL	Humerus Length
MHz	Megahertz
OCF	Occipito-Frontal
OvPAG	Ovine Pregnancy Associated Glycoprotein
P4	progesterone
PAG	Pregnancy Associated Glycoprotein

PD	Pregnancy Diagnosis
PSPB	Pregnancy Specific Protein B
RIA	Radio Immune Assay
Se	Sensitivity
Sp	Specificity
TD	Trunk Diameter
UC	Umbilical Cord

### **ABSTRACT**

Early pregnancy detection in goats was not practicable in the country until now. The present study evaluated the use of ultrasonography in detecting pregnancy as early as 24 days, determining fetal number, viability and establishing gestational equations in a flock of Saanen goats under Sudan conditions.

31 dairy Saanen does aged between 7-31 months were used in the study. They were housed and managed at the main center of goat improvement project, (*Hilat Kuku*), Khartoum state. Estrus was carefully observed and each doe was naturally mated twice on the 1<sup>st</sup> and 2<sup>nd</sup> day of estrus, hence the first day of estrus was designated as day 0 of gestation.

Real-time B- mode ultrasonography was performed with 3.5MHz transabdominal curvilinear transducer. The accuracy of Ultrasound (US) in detecting pregnancy was 100% in both positive (29) cases and negative (2) cases. The accuracy of detecting single feti, twins and triplets was 88.2%, 77.7% and 50% respectively. Blood sample was taken from each animal at the time of Ultrasound scanning for progesterone assay. Contrary to the ultrasound scanning the progesterone assay detected all pregnant animals but failed to detect non pregnant ones, (accuracy 93.5%) or fetal number.

Four nulliparous does of the same age, weight and date of natural mating were scanned at weekly intervals from day 46-67 and 46-95 for measuring (CRL) and (BPD) respectively and Gestational equations were established as follows:

$$\mathbf{CRL = 0.464 x - 17.767}$$

$$\mathbf{BPD = 0.055 x - 1.431}$$

Where  $x$  is the Gestational Age (GA). The correlation coefficient between the (GA) and each of the two parameters was found to be highly significant ( $P < 0.0001$ )

The (CRL) was some times difficult to measure because of the very fast movements of the fetus and the measurements should be taken when the fetus is fully extended which is very difficult to occur. The BPD became increasingly difficult as pregnancy advanced beyond day 95 of gestation. Out of four fetuses two were diagnosed as females and one as male which was confirmed at delivery.

## ملخص الاطروحة

كشف الحمل المبكر فى الاغنام لا يمكن إجراؤه عمليا إلى الآن فى السودان, الدراسة الحالية أجرت تقييم لإستخدام الموجات الصوتية لكشف الحمل المبكر فى عمر 24 يوم, لتحديد عدد الاجنة, بقائها حية و من ثم تكوين معادلة للحمل فى قطيع من أغنام السعانيين تحت ظروف السودان.

عدد 31 من أغنام اللبن سلالة السعانيين تتراوح اعمارها بين 7-31 شهر إستخدمت فى إجراء التجربة, هذه الأغنام موجودة فى مشروع تحسين نسل الماعز المركز الرئيسى /حلة كوكو /ولاية الخرطوم.

تمت المراقبة الدقيقة للشبق وكل انثى تم تلقيحها طبيعيا مرتين فى اليوم الاول والثانى للشبق لذلك أعتبر اليوم الأول للشبق هو يوم صفر فى الحمل.

اجريت الموجات الصوتية الحية من نوع (ب) بإستخدام محول ذو تردد 3.5 ميگاهرتز, كانت نسبة الدقة للموجات الصوتية لكشف الحمل 100% حيث كانت هنالك 29 حالة موجبة للحمل و 2 حالة سالبة . كانت نسبة الدقة لتحديد عدد الأجنة المفرد والمثنى والثلاثى 88.2, 77.7 و 50% على التوالي . أخذت عينة دم من كل حيوان فى نفس الوقت الذى أجريت فيه الموجات الصوتية لفحص هلامون البروجسترون . أظهر فحص هرمون البروجسترون فى الدم وجود 31 حالة موجبة للحمل وذلك بوجود حالتين موجبة كاذبة أى ان نسبة الدقة كانت 93.5% , لم يتمكن هرمون البروجسترون من تحديد عدد الاجنة.

عدد 4 اغنام بكر متساوية فى العمر, الوزن, وتاريخ التلقيح الطبيعى تم الكشف عليها بالموجات الصوتية اسبوعيا إبتداء من يوم 46-67 ومن يوم 46-95 من الحمل لقياس الطول الكلى للجنين والمسافة بين العظمين الجداريين على الترتيب لعمل معادلة للحمل و تم عمل معادلة للحمل كانت كالاتى

الطول الكلى للجنين = 464. س - 17.767

المسافة بين العظمين الجداريين = 0.055 س - 1.431

حيث أن س هي عمر الحمل وكان معامل الارتباط بين عمر الحمل وأى من القياسين قوى جدا

كان قياس الطول الكلى للجنين صعبا فى بعض الاحيان وذلك لسرعة حركة الجنين حيث أن القياس يتم فى حالة الأستقامة التامة للجنين والتي يصعب تحقيقها وكان قياس المسافة بين العظميين الجداريين يزداد صعوبة مع تقدم الحمل بعد يوم 95 عدد أربعة اجنة تم تحديد الجنس لعدد 2 انثى واخر ذكر وتم تاكيد النتائج بعد الولادة.