

References

- [1] Ravinder Nath. Ph.D, Yale University school of Medicine, Department of therapeutic radiology, physical properties and clinical uses of brachytherapy radionuclide. New Haven CT.
- [2] Khan F. M; "Physics of Radiation Therapy", Lippincott Williams & Wilkins, Newyork, London, 429,470,490,491, 1994.
- [3] Ervin B.Podgorsak, MC Gill, Review of Radiation Oncology Physics: A hand book for teacher and students, IAEA, Vienna, Austria, 374-380, May 2003.
- [4] International Commission on Radiation Units and Measurements. Dose and volume specification for reporting intracavitary in gynecology , ICRU report 38 , Bethesda, USA 1995.
- [5] IAEA – TEC Doc -1274, calibration of photon and beta ray sources in brachytherapy , March 2002.
- [6] Lin FJ, Tu CP, Chen CS: Evaluation of quality assurance program in high dose rate remote after-loading Brach therapy. *J Formos Med Assoc Feb*;89(2):139-43(1990).
- [7]DeWerd LA, Jursinic P, Kitchen R, Thomadsen BR: Quality assurance tool for high dose rate Brach therapy. *Med Phys Apr*; 22(4):435-40(1995).
- [8] Li Z, Mitchell TP, Palta JR, Liu C: A quality assurance test tool for high dose-rate remote afterloading brachytherapy units. *Med Phys Feb*;25(2):232-5(1998).
- [9] Goetsch SJ, Attix FH, Pearson DW, Thomadsen BR: Calibration of ¹⁹²Ir high-dose-rate afterloading systems. *Med Phys May-Jun*; 18(3):462-7(1991).
- [10] Henschke.UK "afterloading applicator for radiation therapy of carcinoma of the uterus". *Radiology* 74, 834-839, 1960.
- [11] Walstam R, Remote – controlled afterloading radiotherapy apparatus, physics in *Medicine and Biology*, 7, 225-228, 1962.
- [12] A .A. Martinez, CG. Orton , R.F . Mould, brachytherapy meeting – Remote afterloading, Lippicott Williams &Wilkins, Newyork, London: state of art 4-6 May 1989, Dearborn,
- [13] O, Connell D, Joslin CAF, Howard N, Ramsey Nw& Liversage WE, Te treatment of uterine carcinoma using the Cathetron, *Brit .J.Radiology*, 40, 882-889, 1967.

- [14] Joslin CA, Smith CW & Mallik A, The treatment of cervix cancer using high activity cobalt-60 sources, Brit J. Radiology, 45, 257-270, 1972.
- [15] Bates TD & Berry RJ(Eds), High dose rate afterloading in the treatment of cancer of the uterus, Brit .J. Radiology special report No17, 1978.
- [16] Haybittle, J L and Mitchell, J,S. "simple afterloading technique for treatment of cancer of the cervix". Br. J. Radiol, 48, 295-298, 1975.
- [17] Jack Venslaar, Jose\ Pe\ rez –Calatayud, A practical Guide to Quality control of brachytherapy equipment, European guide lines for quality assurance in radiotherapy, booklet No 8. 3, 13-19, 73-93.
- [18] Dr. SAUERWEIN GMBH -Isotopen –Technik, User Manual –GammaMed 12i.
- [19] William F.Hanson, Ph.D. The university of Texas MD.Anderson cancer centre Houston, TX, brachytherapy source strength: Quantities, Units and Standards.
- [20] Comite\ Francais pour La Me\ sure des Rayonnerments Ionisants(CFMRI)(1983). Recommendation pour La De\ termination des Doses Absorbies en curietherapie,Rapp.C.F.M.R.I No 1CFMRI, Paris.
- [21] International Commission on Radiation Units and Measurements (ICRU)(1985). Dose and volume specification for reporting intracavitary therapy in gynecology ICRU report No 38, ICRU, Washington, DC.
- [22] IAEA –TEC DOC- 1257, Implementation of micro source high dose rate brachytherapy in developing countries, November 2001.
- [23] International Commission on Radiation Units and Measurements. Dose and volume specification for reporting interstitial therapy, ICRU report 58, Bethesda MD 1997.
- [24] Johns, R; Cunningham; "The physics of Radiology" Charles CT Thomas, U.S.A,458, 459, 1983.